

POHL, F., GOUA, M., BERMANO, G., RUSSELL, W.R., MACIEL, P. and KONG THOO LIN, P. 2016. Study into the polyphenol content and antioxidant activity of rapeseed pomace extracts. Presented at *2016 Nutrition Society's Scottish section meeting: phytochemicals and health: new perspectives on plant based nutrition*, 21-22 March 2016, Edinburgh, UK.

Study into the polyphenol content and antioxidant activity of rapeseed pomace extracts.

POHL, F., GOUA, M., BERMANO, G., RUSSELL, W.R., MACIEL, P. and
KONG THOO LIN, P.

2016

Study into the polyphenol content and antioxidant activity of rapeseed pomace extracts

By F. Pohl¹, M. Goua¹, G. Bermano², W. R. Russell³, P. Maciel⁴ and P. Kong Thoo Lin¹,

¹School of Pharmacy and Life Science, Robert Gordon University, Aberdeen, AB10 1GJ, ²Centre for Obesity Research and Education (CORE) Robert Gordon University, Aberdeen, AB10 1GJ, ³Rowett Institute of Nutrition and Health, University of Aberdeen, AB21 9SB, ⁴Life and Health Science Institute (ICVS), Universidade do Minho, Braga, Portugal


Rowett Institute of Nutrition and Health University of Aberdeen

ICVS Life and Health Sciences Research Institute Instituto de Investigação em Ciências da Vida e Saúde

ROBERT GORDON UNIVERSITY ABERDEEN

Introduction

Rapeseed (*Brassica napus*), is a plant used for oil production and naturally grown in Scotland. During the oil production it provides a by-product called rapeseed pomace or rapeseed meal/cake.




Several bioactive compounds have been found in rapeseed, such as tocopherols, phytosterols, phospholipids and phenolic compounds¹. Most abundant phenolics in rapeseed are ferulic acid, salicylic acid, cinnamic acid, p-hydroxybenzoic acid and sinapic acid². Several phenolics have been observed to act as antioxidants and radical scavengers, potentially counteracting oxidative stress³. Due to these properties they could play an important role in the prevention/treatment of oxidative stress related disease.

Aims

The aims of this study was it to determine the polyphenol content, radical scavenging activity, ferric iron reducing antioxidant power and the oxygen-radical absorbance capacity assay, of RSP extracts, with regard to their potential application in the treatment and/or prevention of neurodegenerative diseases.

Methods

Rapeseed pomace obtained from Mackintosh of Glendaveny (Peterhead, Aberdeenshire), was subjected to the following procedures:



^a 2,2-diphenyl-1-picrylhydrazyl assay, ^b liquid chromatography-mass spectrometry, ^c Folin-Ciocalteu assay, ^d Ferric Ion Reducing Antioxidant Power assay, ^e Oxygen Radical Absorbance Capacity assay

Statistically significant differences were analysed via two-way ANOVA and Bonferroni's multiple comparisons test between the two years for each extraction technique ($p < 0.05^*$, $p < 0.01^{**}$, $p < 0.001^{***}$, $p < 0.0001^{****}$, ns-not significant).


Results

- The extraction yield (Table 1) show accelerated solvent extraction (ASE) to be the most efficient, followed by ultrasonic assisted (UAE) and soxhlet extraction (SOX)

Table 1: Extraction yields obtained from rapeseed pomace, for three different extraction techniques and two variant harvests (2012, 2014)

Extraction	Year	Yield [mg/1g pomace]	Yield (%)
SOX	2012	79.55	7.96
	2014	75.59	7.56
UAE	2012	158.59	15.86
	2014	175.46	17.55
ASE	2012	362.31	36.23
	2014	344.67	34.47

- Most abundant phenolics found in the extracts analysed by LC-MS/MS⁴ profiling are listed in Table 2, with sinapic acid (Figure 1) being at the most abundant in all the extracts



Sinapic acid has previously been shown to have anti-inflammatory properties⁵ and attenuated memory impairments in amyloid-B (AB)₁₋₄₂ protein induced mouse model.⁶

Figure 1: Most abundant phenolic sinapic acid (4-hydroxy-3,5-dimethoxycinnamic acid)

Table 2: Most abundant phenolics found in the rapeseed pomace extracts (mean \pm std)

extraction	Year	Most common phenolics (conc. g/kg)		
SOX	2012	sinapic acid (7.497 \pm 0.199)	ferulic acid (0.227 \pm 0.008)	4-hydroxyphenylpyruvic acid (0.172 \pm 0.043)
	2014	sinapic acid (4.897 \pm 0.282)	syringic acid (0.224 \pm 0.017)	ferulic acid (0.183 \pm 0.010)
UAE	2012	sinapic acid (4.610 \pm 0.062)	4-hydroxyphenylpyruvic acid (0.160 \pm 0.015)	ferulic acid (0.126 \pm 0.007)
	2014	sinapic acid (2.769 \pm 0.080)	4-hydroxyphenylpyruvic acid (0.129 \pm 0.005)	syringic acid (0.113 \pm 0.009)
ASE	2012	sinapic acid (2.682 \pm 0.058)	4-hydroxyphenylpyruvic acid (0.1118 \pm 0.008)	ferulic acid (0.071 \pm 0.002)
	2014	sinapic acid (1.923 \pm 0.018)	4-hydroxyphenylpyruvic acid (0.161 \pm 0.014)	cinnamic acid (0.085 \pm 0.003)

- Results obtained for the antioxidant properties of the extracts are shown below:




Figure 2: FC assay




Figure 3: DPPH assay




Figure 4: FRAP assay




Figure 5: ORAC assay

Conclusions

- Investigation into the phenolic content and antioxidant activity showed:
 - Phenolic content in all three extracts is qualitatively similar
 - Sinapic acid is the most abundant phenolic measured in all extracts
 - All extracts show *in vitro* antioxidant activity (FRAP, ORAC and DPPH)
- Extraction technique has a higher significant difference than harvest year
- SOX extracts contained the highest amount of phenolics and the greatest antioxidant capacity for all the *in vitro* analysis
- Next steps to investigate the potential use of rapeseed pomace extracts for the prevention/treatment of oxidative stress related disease are:
 - Antioxidant activity/protective effect in neuroblastoma cell line SH SY-5Y
 - Bioactivity and bioavailability analysis in neurodegeneration related disease models of *C.elegans*

References

- Szydłowska-Czerniak, A. Rapeseed and its products--sources of bioactive compounds: a review of their characteristics and analysis. *Crit. Rev. Food Sci. Nutr.* 53, 307-30 (2013)
- Naczek, M., Amarowicz, R., Sullivan, A. & Shahidi, F. Current research developments on polyphenolics of rapeseed/canola: a review. *Food Chem.* 62, 489-502 (1998)
- Upadhyay, S. & Dixit, M. Role of Polyphenols and Other Phytochemicals on Molecular Signaling. *Oxid. Med. Cell. Longev.* 2015, 15 (2015)
- Russell, W. R. et al. High-protein, reduced-carbohydrate weight-loss diets promote metabolite profiles likely to be detrimental to colonic health. *Am. J. Clin. Nutr.* 93, 1062-1072 (2011)
- Yun, K. J. et al. Anti-inflammatory effects of sinapic acid through the suppression of inducible nitric oxide synthase, cyclooxygenase-2, and proinflammatory cytokines expressions via nuclear factor- κ B inactivation. *J. Agric. Food Chem.* 56, 10265-10272 (2008)
- Lee, H. E. et al. Neuroprotective effect of sinapic acid in a mouse model of amyloid B(1-42) protein-induced Alzheimer's disease. *Pharmacol. Biochem. Behav.* 103, 260-6 (2012)

Acknowledgements

We would like to acknowledge Macintosh of Glendaveny for providing the rapeseed pomace for this project.

We gratefully acknowledge financial support from Tenovus Scotland.

