# Manual patient handling in the healthcare setting: a scoping review.

JOHNSON, K., SWINTON, P., PAVLOVA, A. and COOPER, K.

2023

© 2023 The Authors. Published by Elsevier Ltd on behalf of Chartered Society of Physiotherapy. Supplementary materials are appended after the main text of this document.







Physiotherapy

Physiotherapy 120 (2023) 60-77

# Manual patient handling in the healthcare setting: a scoping review



Katharine Johnson<sup>a,\*</sup>, Paul Swinton<sup>a</sup>, Anastasia Pavlova<sup>a,b</sup>, Kay Cooper<sup>a,b</sup>

<sup>a</sup> School of Health Sciences, Robert Gordon University, Garthdee Road, Aberdeen AB10 7QE, UK

#### **Abstract**

**Background** Manual patient handling is the most frequently reported risk factor for work related musculoskeletal disorders in healthcare. Patient handling tasks are routinely performed manually without assistive devices and can create awkward postures and high loads for nurses and allied health professionals (AHPs). However, AHPs, notably physiotherapists, also utilize therapeutic handling to facilitate patient movement during rehabilitation.

**Objectives** To comprehensively map the literature surrounding manual patient handling (without assistive devices) by healthcare practitioners. **Methods** AMED, CINAHL, MEDLINE, SPORTDiscus, and EMBASE databases were searched. Grey literature was sourced from Google Scholar, EThOS, Open Grey, Health and Safety Executive, National Institute for Occupational Safety and Health and Work Safe Australia. Literature published in English between 2002 and 2021 was included.

**Results** Forty-nine records were included: 36 primary research studies, 1 systematic review and 12 'other' including narrative and government reports. Primary research was predominantly observational cross-sectional (n = 21). The most common settings included laboratories (n = 13) and hospitals (n = 13). Seven research questions were identified, with patient handling practices (n = 13) the most common. Nurses formed the largest practitioner population (n = 13) and patients were often simulated (n = 12). Common outcomes included tasks performed (n = 13) and physical demands during patient handling (n = 13).

Conclusion and implications of key findings This comprehensive scoping review identified that most research was observational, investigating nurses in hospitals or laboratories. More research on manual patient handling by AHPs and investigation of the biomechanics involved in therapeutic handling is needed. Further qualitative research would allow for greater understanding of manual patient handling practices within healthcare.

#### Contribution of the paper

- Much of the research remains observational cross-sectional, investigating patient handling tasks, movement and loading in laboratory settings. Although, the number of studies being performed in hospital settings has increased recently to equal the number performed in controlled laboratory environments.
- Manual handling has been identified as a significant biomechanical factor in development of WRMSD, however, this scoping review identified a lack of literature investigating the relationship between the effect of staff training on incidence of WRMSD.
- This review identified a need for more qualitative research related to moving and handling of patients, to facilitate a deeper understanding
  of the topic.

© 2023 The Authors. Published by Elsevier Ltd on behalf of Chartered Society of Physiotherapy. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

Keywords: Allied health personnel; Delivery of health care; Lifting; Moving and lifting patients; Musculoskeletal diseases; Posture

#### **Background**

Healthcare practitioners routinely assist patients through manual handling across tasks including transferring from bed to chair, personal care tasks, repositioning or rolling in

E-mail address: k.johnson6@rgu.ac.uk (K. Johnson).

<sup>&</sup>lt;sup>b</sup> Scottish Centre for Evidence-based, Multi-professional Practice: A JBI Centre of Excellence, UK

<sup>\*</sup> Corresponding author.

bed, and walking [1]. Nursing staff, support workers, physiotherapists and occupational therapists are among the healthcare practitioners (HCPs) involved with assisting patients within healthcare [1-3]. Work-related musculoskeletal disorders (WRMSDs) are multifactorial, they have a whole range of personal, psychosocial, environmental and biomechanical contributing factors. Psychological behaviours have been highlighted as an important aspects to consider together with biomechanical factors in relation to pain. Unclear messages relating to movement, as a result of uncertainty in the literature, are suggested to result in damaging behaviours such as hypervigilance, fearavoidance and catastrophising [4–6]. Biomechanically, patient handling is frequently documented as the largest risk factor for WRMSDs within HCPs, with low back pain one of the most common complaints [4].

In the UK it is reported that 9.5 million working days are lost to WRMSDs each year with a financial burden of £400 million per annum to the United Kingdom's (UK) National Health Service (NHS) [7]. The United States (US) reported the cost of healthcare workers' related compensation losses as \$2 billion per annum [8], with Australia reporting the cost of serious workplace injuries (1 week or more off work) as A\$14 million within healthcare settings [9]. Investigation of the prevalence of WRMSD within HCPs internationally has reported rates ranging from 28% to 96% in nursing and allied health populations [9,10]. One-year prevalence of WRMSDs within physiotherapists has been reported as 58-67%, with therapeutic handling identified as an associated risk factor [11,12]. However, there is a lack of evidence investigating the risk of injury related to a flexed lifting posture [13].

Legislation on manual patient handling and associated training varies across geographical locations. Staff involved in patient handling within the NHS are required to complete annual online theory and in-person training sessions to learn recommended moving and handling principles to reduce their risk of injury [3]. Within the US, safe patient handling and mobility (SPHM) programs have been implemented in 11 states. These programs aim to improve safety of HCPs by encouraging the use of assistive devices during patient handling [14]. The majority of states, however, have not implemented SPHM programs [14].

Lifting aids and equipment can be used to improve safety of manual patient handling through decreasing the loading experienced by HCPs [14]. There are situations, however, that require nurses and AHPs to manually facilitate patient movement [1]. For example, assistive devices often require use of a sling where placement and removal can be a physically demanding task, involving moving and rolling of the patient. These patient movement tasks are completed manually by HCPs as there are no mechanical devices to aid with these steps [2]. In addition, AHPs, notably physiotherapists use therapeutic handling to aid a patient's rehabilitation. Therapeutic handling is when therapists use their own body in "guiding, facilitating, manipulating or providing resistance" to the patient [3]. The

therapist will manually move and handle the patient to achieve therapeutic benefit [3].

Given the growing body of literature investigating manual patient handling undertaken without the use of assistive devices, this scoping review aimed to map that literature and identify gaps for future research. Further research could benefit HCPs by providing a more comprehensive understanding of manual patient handling. Patient handling techniques could then be improved to reduce risk of injury to HCPs. Manual patient handling in this scoping review was defined as any patient handling task that was completed without the use of an assistive aid including assisting with transfers, moving patients for care tasks or dressing, and placing of slings or sheets under patients. This scoping review forms the first step in a programme of research on manual patient handling.

A preliminary search of MEDLINE, CINAHL, JBI Evidence Synthesis, Open Science Framework, Cochrane library and PROSPERO did not identify any published or in-progress scoping or systematic reviews on the topic.

#### **Review questions**

The objective of this scoping review was to map what information sources were available relating to manual patient handling in healthcare without assistive devices. Two broad review questions related to the primary research identified by the scoping review were: 1) What is the current evidence-base on moving and handling of patients by healthcare practitioners? and 2) What primary research has been conducted on moving and handling of patients by healthcare practitioners? In particular we wanted to answer the following further sub-questions: what questions has the research addressed?; which populations has the research been conducted on?; which settings has the research been conducted in?; which aspects of patient moving and handling have been explored?; and which outcome measures/ techniques/technologies have been used?

#### Methods

This scoping review was conducted in accordance with JBI guidance for scoping reviews [15] and followed an a priori registered open access protocol [16]. The full inclusion and exclusion criteria for the scoping review are outlined in Table 1. Qualified and unqualified staff were included as in some geographical locations (e.g., UK) both staff groups are required to undertake manual handling training and perform patient handling. Literature from any setting was included where it involved HCPs manually assisting patients (real or simulated by healthy volunteers) for tasks or transfers, including in laboratory settings. Guidelines for moving and handling differ across geographical locations. We included literature from any of the

Table 1 Inclusion criteria for scoping review.

	Inclusion Criteria	Exclusion Criteria
Participants	- Qualified or unqualified healthcare staff	Patient handling performed by:
	- Healthcare staff who are involved with manual patient handling	- Students
	tasks (e.g. nurses, AHPs and support staff within nursing and allied	-Healthy volunteers
	health)	(considered where these participants formed less than 50% of
	-Over 18 years of age	sample size and results presented separately from staff)
Concept	Manual patient handling by HCPs in:	Literature solely focused on using mechanical devices (e.g.
	<ul> <li>Healthcare settings</li> </ul>	hoists, standaids)
	<ul> <li>Laboratory settings</li> </ul>	Literature solely investigating epidemiology or prevalence of
	Legislation for correct patient handling	WRMSD
Context	Any healthcare setting where manual patient handling is performed	
	Laboratory settings included where the research objectives could be	
	addressed	
	Literature from highly developed nations (as per HDI)	
Type of Source	Primary research of any type:	Protocols only
	<ul> <li>quantitative, qualitative, mixed methods</li> </ul>	
	Literature reviews of any type:	
	<ul> <li>Narrative texts, systematic reviews, scoping reviews</li> </ul>	
	Conference abstracts if research questions could be answered from	
	abstract	
	Guidelines/legislation around manual patient handling	
	Opinion and text pieces	
	Educational resources for HCPs	

62 very highly developed nations as defined by the Human development Index (HDI) [17] to ensure the findings would be relevant to the UK context.

#### Search strategy

The search strategy aimed to locate published and unpublished primary literature, systematic reviews, text and opinion articles and educational resources. An initial limited search of AMED (EBSCOhost), CINAHL (EBSCOhost) and MEDLINE (PubMed) was undertaken using the keywords (TX moving and handling OR TX manual handling) AND (MH nurse OR TX nurs\* OR TX physiotherap\* OR TX allied health\*). The text words contained in the titles and abstracts of relevant articles, and the index terms used to describe the articles were used to develop a full search strategy. The search strategy, including all identified keywords and index terms, was adapted for each included information source and a second comprehensive search was undertaken on 12th August 2020, and updated on 10th November 2021. Literature published in English from 2002 was included, as the influential Manual Handling Operations regulations 1992 was amended in 2002 [18]. The databases that were searched included: AMED, CI-NAHL, MEDLINE, SPORTDiscus (all via EBSCOhost) and EMBASE (via Ovid). Sources of unpublished and grey literature included: Google Scholar, EThOS, Open Grey, Health and Safety Executive (HSE), National Institute for Occupational Safety and Health (NIOSH), Safe Work Australia, Canadian Centre for Occupational Health and Safety, and Worksafe New Zealand. The full search strategy is provided in supplementary material.

#### Source of evidence selection

Following the searches, all identified records were collated and uploaded into Covidence (v2477; Veritas Health Innovation, Melbourne, Australia) and duplicates removed. Titles and abstracts were initially screened independently by three reviewers (KJ, KC, AP) against the inclusion criteria for the scoping review. Good agreement was found after 10% title and abstract screening. As this scoping review formed part of a doctoral research programme, after good agreement was established, one researcher (KJ) conducted the remainder of title and abstract screening. Three reviewers (KJ, KC, AP) were initially involved in full-text screening, however after good agreement was found following 30% of the screening, one reviewer (KJ) completed full-text screening. Regular review and discussion with the review team was maintained throughout evidence selection. Full-text records that did not meet the inclusion criteria were excluded, and reasons for exclusion are provided in supplementary material.

#### Data extraction

Following JBI methodological guidance, a charting table was created to record key information from the included records. The information extracted is presented in the supplementary material. Data were initially extracted independently from 10% of the included articles by two reviewers (KJ, KC), extractions were consistent, therefore, one reviewer (KJ) completed the remaining data extraction with regular team consultation. Authors of articles were

contacted to request missing or additional data where required; at the time of writing no authors have responded.

Different study types of the included records were charted. The study types included observational cross-sectional, observational cohort, pilot, and qualitative. Observational cross-sectional research included data collected at one point in time, with observational cohort data collected over a period of time. Pilot studies were defined as such by study authors and included feasibility studies and a number of studies where the type was not clearly defined (pilot-undefined).

#### Data analysis and presentation

Search results and included records are summarized in a PRISMA flow diagram [19] (Fig. 1). Summaries of data from all included literature are displayed in Table 2. Previously reported research and populations investigated are grouped into umbrella terms and displayed in tables. Research questions, settings and outcome measures are displayed graphically. A narrative summary accompanies each of the displayed results. The findings are reported aligned to the two research questions. The first broad review question included all the literature (n = 49), the second broad review question included primary research only (n = 36).

#### Results

#### Study inclusion

Initial screening of databases retrieved 8638 records, with an additional 31 records identified from grey sources. Following removal of duplicates, 6956 records remained for title and abstract screening. Of these, 430 records proceeded to full text screening. Forty-nine records met the inclusion criteria and are included in the review (Fig. 1).

#### Characteristics of included studies

A summary of general characteristics of all included records are reported in Table 3. The scoping review included 49 records; one systematic review, 36 primary research studies and 12 other evidence types. The systematic review, published in the US in 2010, included 19 articles and aimed to investigate the effectiveness of health and safety interventions on WRMSDs [18]. The most frequent primary research study design found in this scoping review was observational (n = 24) [2,21–41]. Most of the primary research was published in the US (n = 12) [2,20,24,28,34,41,42,44,46,49,52,53] followed by Canada.




Fig. 1. Search results and study selection and inclusion process [16].

musculoskeletal.
1
MSK
Not reported,
Z
ا م
Ż
literature Kev: NI
litera
Summary of included literature Kev: NF

Author, Year, Country of origin	Aims/purpose	Study type	Setting	Participant profession	Participant age, gender, years experience, description and sample size	Patient population	Outcomes: Domain. Measurement tool
Systematic reviews [17] Tullar, D 2010, he US he ef	ews  Do occupational safety and health interventions in health care settings have an effect on musculoskeletal health status?	Systematic review	Y X	N.A.	NA, 19 studies included	V V	Effects of interventions in healthcare settings on MSK health. Evidence quality appraisal and data extraction and synthesis
OSSIvational [2] Baptiste, 2011, US	To objectively determine the physical demands of patient transfer tasks performed by nurses	Observational cross-sectional	Laboratory	Caregiver	age NR, all male, years exp NR, caregiver represented by 1 male (approx. 6 feet tall and 2001bs), sample size 1	3 mannequins of 3 different weights	Peak force and total impulse. tri-axial load cells
[18] Brusco, 2007, Australia	To develop and implement an allied health OH&S package, based on a risk assessment model, which incorporated clinicians' clinical judgement to minimize manual handling risks, while maximizing the therapeutic benefits for the patient	Observational - cohort	Hospital	5 Physiotherapists, 5 Occupational therapists, manual handling coordinator, allied health assistant	age NR, gender NR, years experience NR, description NA, 200	All health/social care patients	Training completion, MSK injury incidence rate. Staff attendance rates, staff evaluation, incidence rate of injury
[19] Cantarella, 2020, Italy	To validate the effectiveness of MAPO method (Movement and Assistance of Hospital Patient) after the introduction of some changes to improve assessment objectivity	Observational cohort, multi-centre study	Hospital	Health and safety professionals, caregivers	<ul> <li>&lt;35 = 141,</li> <li>35 - 44 = 593,</li> <li>44 - 54 = 801,</li> <li>&gt;55 = 463, majority female, 0-9 = 287,</li> <li>10-19 = 697,</li> <li>20-29 = 642,</li> <li>&gt;30 = 372, 1998</li> </ul>	Various inpatient	Incidence of back pain, risk exposure. BORG scale, staff training. MAPO risk assessment
[20] Carneiro, 2015, Portugal	To identify the main risk factors of WRMSDs for home care nurses and to perform an objective assessment of the risk for these professionals	Observational cross- sectional	Other	Home care nurses	participants age NR, gender NR, years experience NR, description NA, 5	Home care patients	Risk of WRMSDs. Video footage, photographs
[21] Chen, 2014, US	How are nursing work activities distributed over a 12 h day shift? And how does heart rate level differ across nursing work activities?	Observational cross-sectional	Hospital	Nurses	43.4 (SD 8.8), gender NR, 10.6 (SD 6.1), staff nurses, 8	Telemetry unit patients	Heart rate, nursing activities. Heart rate monitor, observation

Continued)
7
able

Author, Year, Country of origin	Aims/purpose	Study type	Setting	Participant profession	Participant age, gender, Patient population years experience, description and sample size	Patient population	Outcomes: Domain. Measurement tool
[22] Hodder, 2010, Canada	To chronicle trunk posture and work tasks of long-term healthcare professionals	Observational cross- sectional	Care home	Personal support workers	46.7 (SD 8.6), All female, all 19.5 (SD 9.3) observed 13.3 (SD 8.6), description NA. 27	Long term care residents	Trunk kinematics and tasks performed. Inclinometer and observation
[23] Hodder, 2010, Canada	To quantify the postural changes that occur with Back Injury Prevention Program (BIPP)	Observational cross- sectional	Laboratory	Nurses, untrained volunteers	41.6 (SD 10.2), All female, 11.3 (SD 9.5), Completed BIPP training in last 2 years, 12 untrained 10 nurses	Healthy volunteer patient (175 cm 81 kg, varying levels of passiveness, weight bearing and following of verbal cues	Muscle activity, 3D thoracolumbar kinematics. EMG, lumbar motion monitor
[24] Holmes, 2010, Canada	To evaluate peak and cumulative lumbar spine loads experienced by personal support workers	Observational cross- sectional	Care home	Personal support workers	47.2 (SD 9.4), All female, 19.6 (SD 9.3), description NA, 20	Long term care patients	Trunk posture, activities completed by each PSW. Observation, inclinometer
[25] Howard, 2013, US	1. To compare the muscle activity of 5 muscle groups of the back, shoulder, and upper extremity between 4 bed-to-wheelchair transfer types. Comparisons are made across the transfer as a whole and between the common components of the transfers. 2. To compare the duration of the components	Observational cross-sectional	Laboratory	1 Occupational therapist, 1 physiotherapist	55 (SD1), All female, 19.5 (SD 13.5), description NA, 2	Volunteer patients simulating varied levels of physical ability	Bilateral muscular effort, duration of task. EMG, video recording
[26] Hye- Knudsen, 2004, Denmark	(tasks) or each transier type. To investigate the kinematics of the thoracolumbar spine during commonly used patient handling tasks.	Observational cross- sectional	Laboratory	Health care workers	43, All female, 19, description NA, 10	Stroke patient, left sided paralysis	Kinematic data, muscle activity. Lumbar motion monitor, EMG, triaxial electrogoniometer
[27] Jordan, 2011, Germany	To perform a detailed investigation on the load of the lumbar spine during manual patient handling	Observational cross- sectional	Laboratory	Health care workers	age NR, All female, Years experience NR, professionally experienced healthcare	Two healthcare workers alternating as simulated partially cooperating patient	Load at IVD L5-S1, considering for posture and action force data. OPTOTRAK

Table 2 (Continued)

Author, Year, Country of origin	Aims/purpose	Study type	Setting	Participant profession	Participant age, gender, years experience, description and sample size	Patient population	Outcomes: Domain. Measurement tool
[28] Kang, 2013, Republic of Korea	To investigate the effects of the application of postural taping on the kinematics of the lumbar spine, pelvis, and hips, EMG activity of the erector spinae, and RPE in the low back during patient transfer in physical	Observational cross-sectional	Laboratory	Physiotherapists	30.68 (SD 4.23), All male, years experience NR, chronic LBP, 19	Healthy volunteer simulated patients	Peak angle and ROM of lumbo-pelvic-hip complex, muscle activity, RPE. VICON motion capture system, EMG, Borg scale
[29] Kim, 2014, Republic of Korea	To analyze, through ergonomic analyzes, through motions most used by radiological technologists that cause musculoskeletal disorders	Observational cross- sectional	Hospital	Radiological technologist	age NR, gender NR, > 5 years experience, description NA, 7	N N	Working postures. Video footage, REBA, RULA, NLE, SI
[30] Kjellberg, 2003, Sweden	To explore the work technique applied by nursing personnel in patient transfer tasks and to determine whether different personal factors were associated with work technique	Observational cross-sectional	Hospital	Nurses, enrolled nurses	35 (SD 10), Majority female, 11 (SD 8.7), mix of participants with and without low-back, neck, shoulder pain, 102	Three healthy women simulated patients	Work technique score, personal factors. Video recordings, observation, questionnaire
[31] Kurowski, 2014, US	To obtain a comprehensive analysis of the physical workload of clinical staff in long-term care facilities, before and after a safe resident handling program (SRHP)	Observational cohort	Care home	Nurse, nursing assistants	age NR, Majority female, 3.03–6.44, description NA, 58–123	Nursing home residents	Average compressive forces on the spine, body postures, manual handling frequencies, biomechanical index.  Observation, physical workload index
[32] Kyriakidis, 2021, Denmark	The purpose of our study was to investigate which organizational levels and factors determine the number of resident handlings in eldercare	Observational cross-sectional	Care home	Eldercare workers	Day shift = 44.4 (SD 10.8), night shift = 47.3 (SD 11), majority female, 15.1 (SD 11.1), 619 staff	Nursing home residents	Factors affecting number of patient handling. Staff seniority level, personal factors, number of patient handling incidences per shift

Table 2 (Continued)

Author, Year, Country of origin	Aims/purpose	Study type	Setting	Participant profession	Participant age, gender, years experience, description and sample size	Patient population	Outcomes: Domain. Measurement tool
[33] Larouche, 2019, Canada	To create an overall risk index that takes account of several aspects of risk, such as awkward postures recorded by a dosimeter, a lifting index, perceived exertion and duration of the task, and to compare the risk associated with patient transfers in total assistance mode observed in real work situations and assigned to three families of transfers	Observational cross-sectional	Other	Paramedics	35 (SD 10), Majority male, 11 (SD 11), description NA, 45	Various	3D angular movements of their back, lifting index, overall risk index. Camcorder, observation, posture dosimeter
[34] Larouche, 2019, Canada	To identify factors that may favor or inhibit the application of safe patient handling principles by paramedics performing fullbody transfers of patients from a chair to a stretcher	Observational cross-sectional	Other	Paramedics	31 (SD 9), Majority male, 8 (SD 9), description NA, 32	Various	Work activity analysis, difficulty of various tasks in the intervention, patient handling methods.  Observation, video footage, semi-structured interview
[35] Maekawa, 2009, Japan		Observational cross-sectional	Laboratory	Nurses	30 s, All female years experience NR, description NA, 2	Nursing staff volunteers	Twist angle of lumbar spine, muscle activity. Goniometer, EMG
[36] Skotte, 2008, Denmark	To investigate the low back load during repositioning of patients in bed and to assess the influence of patient's weight and disability	Observational cross- sectional	Laboratory	Health care workers	46 (SD 9), All female, 5-30 years, description NA, 9	Volunteer patients, I paraplegic with the rest otherwise healthy volunteers simulating hemiplegia, paraplegia and near paralysis	Net torque, compression and shear forces at L4/5, ground reaction forces, reaction force of thighs on the bed.  Digitized video, force platforms, force transducers
[37] Skotte, 2002, Denmark	To investigate the low back-loading during common patient-handling tasks	Observational cross-sectional	Laboratory	Health care workers	43 (SD 8.7), All female, 19 (range 6–26), no special education of training on patient handling technique, 10	Stroke patient, male, 53 years old, 88 kg, left sided weakness	LA/L.5 net moment, compression, shear forces, muscle activity, RPE, ground and bed reaction forces. EMG, Borg scale, video recording, force platforms, force transducers

Table 2 (Continued)

Author, Year, Country of origin	Aims/purpose	Study type	Setting	Participant profession	Participant age, gender, years experience, description and sample size	Patient population	Outcomes: Domain. Measurement tool
[38] Stringer, 2014, US	To investigate the influence that experience in performing manual patient transfers has on the ability to rate the assistance level required during a patient transfer	Observational cross- sectional	Laboratory	Occupational therapist, physical therapist, occupational therapy students, physiotherapy students	50.5 (range 40–56), Majority female, 26.1 (range 15–32), description NA, 23	Volunteer patients	Ground reaction forces of participant and patient, perceived level of assistance the patient required after each picot transfer. Force
[39] Theis, 2014, US	To evaluate the effectiveness of a safe patient handling program (STEPS) at an impatient rehabilitation unit in reducing injury due to patient transfers	Observational cohort	Hospital	Nurses, therapy staff	62.7 (SD 16.4), Equal split, years experience NR, description NA, 55	Inpatient rehabilitation patients	Injury rates pre and post training. Assessment forms
[40] Vieira, 2009, Canada	To quantify physical demands of frequent nursing tasks and provide evidence-based recommendations to increase low back safety	Observational cross-sectional	Laboratory	Nurses	Orthopaedic 35 (SD 7) Intensive care 34 (SD 9), All female, years experience NR, description NA, 36	Same nursing participants simulating patients	Lumbar ROM, motion during nursing tasks, L5/S1 compression and shear forces estimated, sufficient torso strength estimated. Electrogoniometer, perpendicular marker photogrammetry
Pilot studies [41] Arias, 2017, US	To characterize the physical load of trunk flexion and physical activity of patient care unit (PCA) workers	Pilot - undefined	Hospital	Nurses, patient care assistants	42 (SD 13), Majority female, Years experience NA, description NA, 50	Thoracic intensive care, orthopedic, burn and trauma, cardiac and cardiac step down	Physical activity, trunk flexion. Accelerometer, tri- axial accelerometer
[42] Fiedler, 2012, Canada	To determine the feasibility of documenting all jobrelated nursing tasks performed during a typical shift in a hospital setting using video	Pilot - feasibility	Hospital	Nurses	40.6 (SD 13.3), 13.6 (SD 11.2), description NA, 10	Intensive care unit, inpatient rehabilitation, complex continuing care unit, acute care unit, outpatient surgery, ambulatory care ambulatory care	Patient handling tasks. Camcorder, observation
[43] Fragala, 2011, US	To quantify and objectively measure the risk reduction achieved with the gravity assist feature	Pilot - undefined	Laboratory	Caregivers	age NR, gender NR, years experience NR, description NR, cannele NR	200 lb mannequin	Total force, peak force, total work, slope of the force curve. Force transducer
[44] Garzillo, 2020, Italy	To propose an MPH training model involving interdisciplinary aspects	Pilot - undefined	Hospital	Healthcare workers	men 49.4 (SD 7.2) women 45.9 (SD 8.8), equal split of gender, 24.6 (SD 8.1), 52 staff members	Various inpatient	Risk assessing tasks, staff training, effort required for tasks. Questionnaire, multidisciplinary training program, BORG scale

Table 2 (Continued)

Author, Year, Country of origin	Aims/purpose	Study type	Setting	Participant profession	Participant age, gender, years experience, description and sample size	Patient population	Outcomes: Domain. Measurement tool
[45] Newton, 2020, Australia	To ascertain the incidence of Australian private practice sonographers moving patients unassisted and determine what training these sonographers have in order to appropriately perform these procedures	Pilot - undefined	Other - sonography clinics	Sonographers	age NR, gender NR, <5 = 24, 6-10 = 6, 11-20 = 4, > 21 = 1, 35 respondents	Outpatient	Incidences of manual assistance of patients, level of training. Survey
Qualitative [46] de Ruiter, 2011 US	To identify patient-handling practices in clinical practice	Qualitative	Hospital	Registered nurse, nursing assistant	age: $<24 = 7$ $24-36 = 13 \ 40-59 = 9$ 60 + = 3, Majority female, $< 1 = 3 \ 1-3 = 9$ $4-10 = 10 \ 11-20 = 7$ 21 + = 3	Neurology and rehabilitation patients	Patient handling practices and nurses' judgment of these practices, observation of caregivers and interviews
[47] Osborne, 2021, Australia	To investigate emergency nurses' beliefs and experiences with patient handling in the emergency department	Qualitative - phenomenological	Hospital	Nurses	age NR, majority female, mean 3.9 years (range 1-27), 40 participants	ED patients	Experiences of patient handling. Focus group interviews
[48] Wangbad, 2009, Sweden	To describe nurses aids' experiences of physical strain during person transfer tasks as dementia care units	Qualitative	Hospital	Nurses aids	43 (range 26–64), All female, 15 (range 2–31), description NA, 16	Dementia patients	Experiences, apprehensions, person transfer tasks. Focus groups
Survey [49] [49] [2007, US	To describe patient-handling demands in inpatient units during a 24-hour period at a military health care facility	Survey	Hospital	Nurses, licensed practical nurses, nursing assistants, and nursing students	35 (SD 10.7), Majority female, Median 6 (range 0.5–38), description NA, 283	Military inpatients	Nature and impact of patient- handling tasks relative to a variety of nursing care units, patient characteristics, and transfer equipment. 24-hour population survey
Collections abstract [50] Lavender, To 2016, an US shirt pat	To measure the compression and shear loads on the spine that are experienced as slings are (1) placed under patients prior to lifting, and (2) removed at the completion of the transfer	Conference abstract - repeated measures study	Laboratory	Nurses	age NR, gender NR, years experience NR, description NA, 12	simulated patients, 54 kg and 100 kg	Muscle activity, lumbar spine compression, shear forces. EMG, digitized video
[51] Nikolajsen, 2015, Denmark	To document and describe how manual patient handling may be carried out as part of everyday practice	Conference abstract - Mixed methods	Other	Health care providers	age NR, gender NR, years experience NR, description NA, 56	Various	Manual patient handling activities. Field notes and observation.

Table 2 (Continued)

Author, Year, Country of origin	Aims/purpose	Study type	Setting	Participant profession	Participant age, gender, years experience, description and sample size	Patient population	Outcomes: Domain. Measurement tool
[52] Wade, 2017, UK	To investigate healthcare staff perceptions surrounding manual handling and patient handling related injuries	Conference abstract - ethnographic study	Community	Physiotherapists, occupational therapists	age NR, gender NR, years experience NR, 8 participants for observation, 6 for interviews	Community	Manual patient handling factors, challenges/benefits, perceptions of training. Observation, focus groups
[53] Apple, 2021, US	To review published research and describe the ergonomic challenges of working in the OR	Other, narrative review	N A	NA	all NA	Peri-operative patients	Ergonomic challenges, safety recommendations. Data from literature
[54] Haney, 2003, US	To review various physical stressors as they relate to the capacity of the human body during handling (lifting, transferring, moving, and walking) of residents	Other, narrative review	NA A	NA	all NA	N A	Force and repetition,dynamic lifting, posture, risk management, push/pull forces, philosophy of care. Data from literature
[55] Hignett, 2007, UK	Transport of the control of EU Health and Safety Directive on Manual Handling (90/269/EEC) for patient handling in 9 European countries and gather expert opinion on the residual problems (barriers) to safer parient handling.	Other, Narrative summary of government statistics and expert opinion from a panel of manual handling experts	Υ V	NA A	all NA	All health/social care patients	Implementation of the EU manual handling directive, expert opinions. European of National Government statistics, expert opinion.
[56] Johnstone, 2020, UK	potential interactions of the control of the contro	Other, narrative review	N A	NA	all NA	All health/social care patients	WRMSD in nursing related to moving and handling. Information from legislation and oridance materials
[57] Rinds, 2008, UK	To consider the importance of safe manual handling, risk assessment, the law and useful equipment designed to aid the care worker	Other, Narrative summary of law and guidance in manual handling in healthcare	V.	Υ <sub>Z</sub>	all NA	All health/social care patients	Safety during manual patient handling. Information and evidence from laws and guidance.
[58] Tofts, 2012, UK	To explain the legislation related to moving and handling, with particular application to community nurses	Other, Narrative summary of legislation surrounding moving and handling in the community	Z A	N.A	all NA	All community patients	Legislation surrounding moving and handling. Information from health and safety publications

Table 2 (Continued)

Author, Year, Country of origin	Aims/purpose	Study type	Setting	Participant profession	Participant age, gender, years experience, description and sample size	Patient population	Outcomes: Domain. Measurement tool
[59] Vatwani, 2017, US	To provide practical information and instruction for caregivers assisting individuals experiencing difficulty performing bed mobility tasks.	Other, Educational short article	NA	NA	all NA	All health/social care patients	Educational resources.
[60] Waters, 2007, US	To describe high-risk patient handling tasks performed frequently in critical care units, delineate the physical demands associated with each task, identify technologic solutions, and outline useful tips for making each task safer	Other, Ergonomic task force identified tasks with high risk for MSK disorders	Hospital	Nurses, nurse managers	all NR	NR T	Critical care tasks that are high force, awkward posture, repetitive loading. Observation.
[61] Waters, 2011, US	To determine the best practices for safe lateral patient transfers	Other, Ergonomic tool for positioning	NA	NA	all NA	All health/social care patients	Ergonomic tool development. Risk assessment tools.
[62] Waters, 2011, US	To determine the best practices for safe positioning and repositioning of the supine patient	Other, Ergonomic tool for positioning	NA	٩	all NA	All health/social care patients	Safety during supine patient handling. Risk assessment tools.
[63] Weiner, 2015, Israel	To present current research about the risk factors, prevention strategies, and assistive devices that could reduce work-related musculoskeletal disorders caused by repositioning patients in bed	Other, Narrative summary	ę Z	٩	AN IIs	All health/social care patients	Association between WRMSDs and repositioning patients in bed, risk factors for WRMSD during patient repositioning in bed.  Published literature and data.
[64] Weinmeyer, 2016, US	Laws and programs to address the problem of nursing-specific musculoskeletal injuries.	Other, Narrative summary of laws and programs for safe patient handling for health care workers	NA	V.A	all NA	All health/social care patients	Working conditions leading to injuries, patient handling related injuries, barriers to safe patient handling. Published laws, programs.

Table 3 Overview of reports included in scoping review (n = 49) by study type, year of publication, country of origin and sample size of number of studies included presented as a percentage of total included reports.

Research - Type of stury included (n = 37)		1		
Observational cross-sectional   Sectional   Sectiona		tudy		ture
Pilot – undefined         4 (8%)         Task force         1 (2%)           Conference abstract         3 (6%)         Educational page         1 (2%)           Qualitative         3 (6%)         Educational page         1 (2%)           Pilot – feasibility         1 (2%)         Systematic review         1 (2%)           Systematic review         1 (2%)         Year of publication           2017–2021         9 (18%)         2017–2021         3 (6%)           2012–2016         11 (22%)         2012–2016         3 (6%)           2007–2011         14 (29%)         2007–2011         5 (10%)           2002–2006         3 (6%)         2002–2006         1 (2%)           Country of origin         Country of origin           United States         12 (24%)         United States         7 (14%)           Canada         7 (14%)         United Kingdom         4 (8%)           Sweden         2 (4%)         Israel         1 (2%)           Australia         3 (6%)         3 (6%)           Sweden         2 (4%)         Israel         1 (2%)           Germany         1 (2%)         Israel         1 (2%)           Sample size and number of included studies         < 10         2 (2%)	Observational cross-	20 (41%)	Narrative summaries of legislation or	8 (16%)
Conference abstract 3 (6%) Educational page 1 (2%) Qualitative 3 (6%) Pilot – feasibility 1 (2%) Survey 1 (2%) Systematic review 1 (2%) Year of publication 2017–2021 9 (18%) 2017–2021 3 (6%) 2012–2016 11 (22%) 2012–2016 3 (6%) 2007–2011 14 (29%) 2007–2011 5 (10%) 2002–2006 3 (6%) 2002–2006 1 (2%) Country of origin United States 12 (24%) United States 7 (14%) Canada 7 (14%) United Kingdom 4 (8%) Denmark 5 (10%) Israel 1 (2%) Sweden 2 (4%) Republic of Korea 2 (4%) Republic of Korea 1 (2%) United Kingdom 1 (2%) Japan 1 (2%) Germany 1 (2%) Germany 1 (2%) Sample size and number of included studies <10 10 (20%) 10–19 9 (18%) 20–29 2 (4%) 30–39 4 (8%) 40–49 4 (8%) >50 9 (18%) Range 1–1998 Median 23 Interquartile range 41.5	Observational cohort	4 (8%)	Ergonomic tool	2 (4%)
Qualitative 3 (6%) Pilot – feasibility 1 (2%) Survey 1 (2%) Systematic review 1 (2%)  Year of publication 2017–2021 9 (18%) 2017–2021 3 (6%) 2007–2011 14 (29%) 2007–2011 5 (10%) 2002–2006 3 (6%) 2002–2006 1 (2%)  Country of origin United States 12 (24%) United States 7 (14%) Denmark 5 (10%) Israel 1 (2%) Sweden 2 (4%) Republic of Korea 2 (4%) Republic of Korea 2 (4%) Italy 2 (4%) Portugal 1 (2%) Japan 1 (2%) Germany 1 (2%) Germany 1 (2%) United Kingdom 1 (2%) Sample size and number of included studies < 10 10 (20%) United Kingdom 1 (2%) Sample size and number of included studies < 10 10 (20%) 10–19 9 (18%) 20–29 2 (4%) 30–39 4 (8%) 40–49 4 (8%) > 50 9 (18%) Range 1–1998 Median 23 Interquartile range 41.5	Pilot - undefined	4 (8%)	Task force	1 (2%)
Pilot – feasibility       1 (2%)         Survey       1 (2%)         Systematic review       1 (2%)         Year of publication       Year of publication         2017–2021       9 (18%)       2017–2021       3 (6%)         2007–2016       11 (22%)       2012–2016       3 (6%)         2007–2011       14 (29%)       2007–2011       5 (10%)         2002–2006       3 (6%)       2002–2006       1 (2%)         Country of origin       Country of origin         United States       12 (24%)       United States       7 (14%)         Canada       7 (14%)       United Kingdom       4 (8%)         Denmark       5 (10%)       Israel       1 (2%)         Australia       3 (6%)       Sweden       2 (4%)         Republic of Korea       2 (4%)       Israel       1 (2%)         Bepublic of Korea       2 (4%)       Italy       2 (4%)         Portugal       1 (2%)       Israel       1 (2%)         Germany       1 (2%)       Israel       1 (2%)         Sample size and number of included studies       < 10	Conference abstract	3 (6%)	Educational page	1 (2%)
Survey       1 (2%)         Systematic review       1 (2%)         Year of publication       Year of publication         2017–2021       9 (18%)       2017–2021       3 (6%)         2007–2016       11 (22%)       2012–2016       3 (6%)         2007–2011       14 (29%)       2007–2011       5 (10%)         2002–2006       3 (6%)       2002–2006       1 (2%)         Country of origin       Country of origin         United States       12 (24%)       United States       7 (14%)         Canada       7 (14%)       United Kingdom       4 (8%)         Denmark       5 (10%)       Israel       1 (2%)         Australia       3 (6%)       Sweden       2 (4%)         Republic of Korea       2 (4%)       Israel       1 (2%)         Portugal       1 (2%)       Israel       1 (2%)         Japan       1 (2%)       Israel       1 (2%)         United Kingdom       1 (2%)       Israel       1 (2%)         Sample size and number of included studies       < 10	Qualitative	3 (6%)		
Systematic review         1 (2%)           Year of publication         Year of publication           2017–2021         9 (18%)         2017–2021         3 (6%)           2012–2016         11 (22%)         2012–2016         3 (6%)           2007–2011         14 (29%)         2007–2011         5 (10%)           2002–2006         3 (6%)         2002–2006         1 (2%)           Country of origin         Country of origin         United States         7 (14%)           Canada         7 (14%)         United Kingdom         4 (8%)           Denmark         5 (10%)         Israel         1 (2%)           Australia         3 (6%)         Sweden         2 (4%)           Republic of Korea         2 (4%)         Israel         1 (2%)           Republic of Korea         2 (4%)         Israel         1 (2%)           Japan         1 (2%)         Israel         1 (2%)           United Kingdom         1 (2%)         Israel         1 (2%)           Sample size and number of included studies         < 10	Pilot - feasibility	1 (2%)		
Year of publication         Year of publication           2017–2021         9 (18%)         2017–2021         3 (6%)           2012–2016         11 (22%)         2012–2016         3 (6%)           2007–2011         14 (29%)         2007–2011         5 (10%)           2002–2006         3 (6%)         2002–2006         1 (2%)           Country of origin         Country of origin           United States         12 (24%)         United States         7 (14%)           Canada         7 (14%)         United Kingdom         4 (8%)           Denmark         5 (10%)         Israel         1 (2%)           Australia         3 (6%)         Sweden         2 (4%)           Republic of Korea         2 (4%)         Israel         1 (2%)           Portugal         1 (2%)         Israel         1 (2%)           Japan         1 (2%)         Israel         1 (2%)           Germany         1 (2%)         Israel         1 (2%)           Sample size and number of included studies         < 10	Survey	1 (2%)		
2017–2021       9 (18%)       2017–2021       3 (6%)         2012–2016       11 (22%)       2012–2016       3 (6%)         2007–2011       14 (29%)       2007–2011       5 (10%)         2002–2006       3 (6%)       2002–2006       1 (2%)         Country of origin         United States       12 (24%)       United States       7 (14%)         Canada       7 (14%)       United Kingdom       4 (8%)         Denmark       5 (10%)       Israel       1 (2%)         Australia       3 (6%)       Sweden       2 (4%)         Republic of Korea       2 (4%)       Sweden       2 (4%)         Portugal       1 (2%)       Sweden       2 (4%)         Japan       1 (2%)       Sweden       3 (36%)         Sample size and number of included Kingdom       1 (2%)       Sweden       3 (36%)         Sample size and number of included studies       < 10		1 (2%)		
2012-2016       11 (22%)       2012-2016       3 (6%)         2007-2011       14 (29%)       2007-2011       5 (10%)         2002-2006       3 (6%)       2002-2006       1 (2%)         Country of origin         United States       12 (24%)       United States       7 (14%)         Canada       7 (14%)       United Kingdom       4 (8%)         Denmark       5 (10%)       Israel       1 (2%)         Australia       3 (6%)       Sweden       2 (4%)         Republic of Korea       2 (4%)       Israel       1 (2%)         Portugal       1 (2%)       Israel       1 (2%)         Germany       1 (2%)       Israel       1 (2%)         Germany       1 (2%)       Israel       1 (2%)         Sample size and number of included studies       < 10	Year of publication		Year of publication	
2007-2011       14 (29%)       2007-2011       5 (10%)         2002-2006       3 (6%)       2002-2006       1 (2%)         Country of origin         United States       12 (24%)       United States       7 (14%)         Canada       7 (14%)       United Kingdom       4 (8%)         Denmark       5 (10%)       Israel       1 (2%)         Australia       3 (6%)       Sweden       2 (4%)         Republic of Korea       2 (4%)       Republic of Korea       2 (4%)         Italy       2 (4%)       Portugal       1 (2%)         Germany       1 (2%)       Sample size and number of included studies       1 (2%)         < 10	2017-2021	9 (18%)	2017-2021	3 (6%)
2002–2006     3 (6%)     2002–2006     1 (2%)       Country of origin     Country of origin       United States     12 (24%)     United States     7 (14%)       Canada     7 (14%)     United Kingdom     4 (8%)       Denmark     5 (10%)     Israel     1 (2%)       Australia     3 (6%)     Sweden     2 (4%)       Republic of Korea     2 (4%)     Fortugal     1 (2%)       Italy     2 (4%)     Fortugal     1 (2%)       Japan     1 (2%)     Fortugal     1 (2%)       Germany     1 (2%)     Fortugal     1 (2%)       Sample size and number of included studies     < 10	2012-2016	11 (22%)	2012-2016	3 (6%)
Country of origin         Country of origin           United States         12 (24%)         United States         7 (14%)           Canada         7 (14%)         United Kingdom         4 (8%)           Denmark         5 (10%)         Israel         1 (2%)           Australia         3 (6%)         Sweden         2 (4%)           Republic of Korea         2 (4%)         Fortugal         1 (2%)           Italy         2 (4%)         Fortugal         1 (2%)           Japan         1 (2%)         Fortugal         1 (2%)           Germany         1 (2%)         Fortugal         1 (2%)           Sample size and number of included studies         < 10	2007-2011	14 (29%)	2007-2011	5 (10%)
United States 12 (24%) United States 7 (14%) Canada 7 (14%) United Kingdom 4 (8%) Denmark 5 (10%) Israel 1 (2%) Australia 3 (6%) Sweden 2 (4%) Republic of Korea 2 (4%) Italy 2 (4%) Portugal 1 (2%) Japan 1 (2%) Germany 1 (2%) United Kingdom 1 (2%) Sample size and number of included studies <10 10 (20%) 10–19 9 (18%) 20–29 2 (4%) 30–39 4 (8%) >50 9 (18%) Range 1–1998 Median 23 Interquartile range 41.5	2002-2006	3 (6%)	2002-2006	1 (2%)
Canada       7 (14%)       United Kingdom       4 (8%)         Denmark       5 (10%)       Israel       1 (2%)         Australia       3 (6%)       Sweden       2 (4%)         Republic of Korea       2 (4%)       Fortugal       1 (2%)         Italy       2 (4%)       Fortugal       1 (2%)         Japan       1 (2%)       Fortugal       1 (2%)         Germany       1 (2%)       Fortugal       1 (2%)         Sample size and number of included studies       410       10 (20%)         10-19       9 (18%)       9 (18%)         20-29       2 (4%)       30-39       4 (8%)         40-49       4 (8%)       50       9 (18%)         Range       1-1998       Median       23         Interquartile range       41.5       41.5	Country of origin		Country of origin	
Denmark       5 (10%)       Israel       1 (2%)         Australia       3 (6%)	United States	12 (24%)	United States	7 (14%)
Australia 3 (6%) Sweden 2 (4%) Republic of Korea 2 (4%) Italy 2 (4%) Portugal 1 (2%) Japan 1 (2%) Germany 1 (2%) United Kingdom 1 (2%) Sample size and number of included studies < 10 10 (20%) 10–19 9 (18%) 20–29 2 (4%) 30–39 4 (8%) 40–49 4 (8%) > 50 9 (18%) Range 1–1998 Median 23 Interquartile range 41.5	Canada	7 (14%)	United Kingdom	4 (8%)
Sweden       2 (4%)         Republic of Korea       2 (4%)         Italy       2 (4%)         Portugal       1 (2%)         Japan       1 (2%)         Germany       1 (2%)         United Kingdom       1 (2%)         Sample size and number of included studies         < 10	Denmark	5 (10%)	Israel	1 (2%)
Republic of Korea       2 (4%)         Italy       2 (4%)         Portugal       1 (2%)         Japan       1 (2%)         Germany       1 (2%)         United Kingdom       1 (2%)         Sample size and number of included studies         < 10	Australia	3 (6%)		
Italy       2 (4%)         Portugal       1 (2%)         Japan       1 (2%)         Germany       1 (2%)         United Kingdom       1 (2%)         Sample size and number of included studies         < 10	Sweden	2 (4%)		
Portugal 1 (2%) Japan 1 (2%) Germany 1 (2%) United Kingdom 1 (2%)  Sample size and number of included studies  < 10 10 (20%) 10–19 9 (18%) 20–29 2 (4%) 30–39 4 (8%) 40–49 4 (8%) > 50 9 (18%) Range 1–1998 Median 23 Interquartile range 41.5	Republic of Korea	2 (4%)		
Japan 1 (2%) Germany 1 (2%) United Kingdom 1 (2%)  Sample size and number of included studies  < 10 10 (20%) 10–19 9 (18%) 20–29 2 (4%) 30–39 4 (8%) 40–49 4 (8%) > 50 9 (18%) Range 1–1998 Median 23 Interquartile range 41.5	Italy	2 (4%)		
Germany 1 (2%) United Kingdom 1 (2%)  Sample size and number of included studies  < 10 10 (20%) 10–19 9 (18%) 20–29 2 (4%) 30–39 4 (8%) 40–49 4 (8%) > 50 9 (18%) Range 1–1998 Median 23 Interquartile range 41.5	Portugal	1 (2%)		
United Kingdom 1 (2%)  Sample size and number of included studies  < 10 10 (20%) 10–19 9 (18%) 20–29 2 (4%) 30–39 4 (8%) 40–49 4 (8%) > 50 9 (18%) Range 1–1998 Median 23 Interquartile range 41.5	Japan	1 (2%)		
Sample size and number of included studies         < 10	•	. ,		
included studies       < 10				
< 10		ber of		
10–19 9 (18%) 20–29 2 (4%) 30–39 4 (8%) 40–49 4 (8%) > 50 9 (18%) Range 1–1998 Median 23 Interquartile range 41.5	included studies			
20–29 2 (4%) 30–39 4 (8%) 40–49 4 (8%) > 50 9 (18%) Range 1–1998 Median 23 Interquartile range 41.5	< 10	10 (20%)		
30–39 4 (8%) 40–49 4 (8%) > 50 9 (18%) Range 1–1998 Median 23 Interquartile range 41.5	10–19	9 (18%)		
40-49       4 (8%)         > 50       9 (18%)         Range       1-1998         Median       23         Interquartile range       41.5		2 (4%)		
> 50 9 (18%) Range 1–1998 Median 23 Interquartile range 41.5	30–39	4 (8%)		
Range 1–1998 Median 23 Interquartile range 41.5				
Median 23 Interquartile range 41.5		9 (18%)		
Interquartile range 41.5	Range	1–1998		
· ·	Median	23		
Systematic review 19				
	Systematic review	19		

#### Narrative literature

The 'other' literature included narrative text (n=8) [56–61,66,67] and educational pieces (n=4) [62–65]. Narrative texts described legislation, directives and previous research and were largely published in the US (n=7). Additionally, narrative texts and opinion pieces provided education materials and guidance for improving manual patient handling in settings ranging from the community to operating theatres.

The following sections of the scoping review are focused on the primary research identified (n = 36) in order to address the review sub-questions.

#### Questions addressed by primary research

Aims and objectives of the included primary research were reviewed and multiple research questions were identified, with some comprising sub-questions. The research questions included: 1) physical demands during manual patient handling (n = 13) [2,22,24,28,31,34,36,41,43,44,47,51,52]; 2) patient handling practices and tasks performed (n = 13)[24,25,27,32–36,45,48,49,52,54]; 3) improving safety of patient handling (n=7) [21,26,34,42,43,46,47]; 4) risk assessment of patient handling tasks (n = 8 [21-23,32,36-38,46]); 5) investigation of kinetics (loading) experienced by the HCP (n = 7)[2,27,30,38-40,53]; 6) investigation of kinematics (joint motion) (n=7) [25,26,29,31,32,36,44]; and 7) personal factors affecting patient handling (n = 9) [31,33,35,37,39,41,50,51,55].

#### **Populations**

The summary of populations included in the review is displayed in Table 4. The largest staff population included in the review was nursing, with many studies investigating both qualified and unqualified staff (n = 9)

Table 4 Healthcare staff groups, age, gender, years of experience and patient populations included in research (n = 36).

Nursing	
Registered nurses	14 (39%)
Healthcare workers	13 (36%)
Nursing assistants	4 (11%)
Allied Health	
Physiotherapists	5 (14%)
Occupational therapists	4 (11%)
Paramedics	2 (5%)
Allied health assistants	1 (3%)
Therapy staff	1 (3%)
Radiological technologists	1 (3%)
Other	
Manual handling coordinators	1 (3%)
Health and safety professional	1 (3%)
Age range	, í
< 20–30	1 (3%)
31–40	9 (25%)
41–50	14 (39%)
51–60	4 (11%)
>61	2 (5%)
Gender	
All female	2 (5%)
Majority female	15 (42%)
All male	2 (5%)
Majority male	6 (17%)
Equal split	2 (5%)
Years of experience	
<1	1 (3%)
2–5	6 (17%)
6–10	7 (19%)
11–15	12 (33%)
16–20	9 (25%)
>20	7 (19%)
Patient population	
Simulated by volunteers	12 (33%)
Inpatient	11 (31%)
Care home	4 (11%)
Community	4 (11%)
Mannequins	2 (5%)
Unspecified	2 (5%)
Outpatient	1 (3%)

[21,25,33,34,41,42,44,49,52]. There was substantial heterogeneity in terminology for staff populations across studies limiting the accuracy of extraction. Physiotherapists were investigated in 5 of the included primary research [21,28,31,41,55]. Most participants were aged between 31 and 50 years and typically comprised all-female or majority female cohorts. The most common patient population included in primary research were volunteer simulated patients (n = 12) [26,28–31,33,38–41,43,53].

#### Settings

Laboratory (n = 13) [2,26,28-31,38-41,43,46,53] and hospital (n = 13) [21,22,24,32,33,42,44,46,47,49-52] settings were the most common settings used in primary research. Investigations into kinetics and kinematics were most frequently performed in laboratory settings with observation of patient handling practices investigated more frequently in hospital settings.

#### Aspects of manual patient handling

Relevant information was extracted to address the seven specific research questions of the scoping review and are presented in Table 5. A broad range of aspects were investigated within each research question. Investigating the tasks performed and estimation of spinal forces were some of the most frequently identified aspects in the primary research. Personal factors largely investigated practitioners experiences, perceptions and how staff and patient factors impact patient handling practices.

#### Outcome measures

Outcome measures were separated into measurement domains and tools, with the measurement tools grouped according to the domain they were used to measure, and are displayed proportionately in Fig. 2. Physical demands and tasks performed were the most frequently investigated domains. Kinematics showed the most variation in measurement tool. Two primary research studies investigated staff training and its effect on incidence of reported WRMSD [21,42]. One study quantified the risk of WRMSD with associated with nursing tasks by using ergonomic assessment tools [22]. A table of all included outcomes and references are included in the supplementary materials. Within primary research published in the last 5 years there has been an increase in use of qualitative measurement methods, especially to investigate practitioner perceptions and experiences.

#### Discussion

In this scoping review, the literature on manual patient handling in healthcare was identified and examined, providing a comprehensive map. This allowed for a focus on manual patient handling tasks that require manual assistance from HCPs or involve therapeutic handling. The literature comprised a range

Table 5
Summary of aspects of included literature.

Question	Aspect
Physical demands	Trunk position/loading [40,41]
	Fatiguing/straining tasks for HCPs [2,31,48]
	Potential solution for demanding tasks [2,48]
	Muscular effort [25]
	Perceived exertion[19,33,44,49]
	Level of assistance provided by HCPs [38,44]
	Heart rate [21]
Patient handling	Tasks performed by HCPs
practices	[21,24,32,42,45,46,49,51]
	Time taken for tasks [29,42,46,49]
	Staff positioning [21,29]
	Number of staff used [23,49]
	Patient factors (e.g. weight) [23,49]
	Ergonomic assessment [29–31]
Improving safety	Staff training [18,22,39,44]
	Identification of high risk tasks [40]
Risk assessment	Creation of staff training package [18]
	Risk assessing tasks [19,20,29,33,34]
	Risk reduction measurement [43]
	Risk prediction [35]
Kinetics	Estimation of spinal forces [2,24,27,35–37,50]
	Ground reaction forces [27,36]
	Action forces at hands/thighs [27]
Kinematics	Trunk movement during patient handling
	[23,26,33,41]
	Effect of training on trunk movement [22]
	Effect of taping on trunk movement [28]
	Analysis of full body postures [29]
Personal factors	Impact of patient factors (e.g. weight/level of
	disability) [36,48]
	Staff knowledge and impact on patient handling
	[48]
	Staff age, gender, experience level and impact
	on patient handling [28,30,32,38]
	Physical constraints experienced [34]
	Experiences of patient handling [47,48]
	Perceptions of moving and handling [44,46,52]

of records with a variety of outcome measurement tools. Primary research studies incorporated a variety of outcome tools to address their specific research question.

#### Research

This scoping review included 49 records, of which only one was a systematic review published in 2010. This suggests a need for further robust evidence synthesis, which can inform practice within healthcare. As will be highlighted further in the discussion, potential systematic reviews for which there may be adequate evidence include HCPs movement during manual patient handling tasks. The most common study design was observational cross-sectional. These studies are generally time efficient and inexpensive to perform and allow for investigation of multiple outcomes [68]. Observational studies, however, are generally unable to establish cause and effect and may identify a range of spurious factors that would lead to ineffective interventions if developed [68]. Further well-designed research




Fig. 2. Outcome measures from the primary research (n = 36) separated into measurement tools (outer ring) and the domain that they were used to collect data for (inner ring)(size of sections based on relative count for each measurement tool). Key: LMM – Lumbar motion monitor; Posture D – Posture dosimeter; TG – Tri-axial goniometer; TA – Tri-axial accelerometer; PWI – Physical workload index; TLC – Tri-axial load cells; Erg Assessment – Ergonomic assessment; 24-h – 24-h Survey.

including therapeutic handling, handler movement and handler perceptions is required to better establish the factors that increase risk of injury and to provide appropriate targets for interventions.

The evidence base investigating AHPs is small in comparison to nurses. Nursing forms the largest occupational group within healthcare globally [69] and a high rate of WRMSDs has been found within the profession [70]. However, AHPs perform many of the same tasks, in addition to therapeutic handling to aid patients' rehabilitation. Therapeutic handling can be manually intensive with little research exploring how these tasks are performed, how manually intensive they are, and the potential risks associated with performing them regularly. Within the five included records that investigated physiotherapists, three investigated patient transfers with only one investigating therapeutic handling.

This scoping review found that it is more common to explore which tasks are performed during manual patient handling than to investigate how they are performed and their burden through exploration of biomechanics including the joint angles and internal forces experienced. Research conducted more recently investigated personal factors, experiences and perceptions of patient handling with a variety of observational and qualitative methods. Many studies used healthy volunteers as simulated patients, especially in laboratory-based research. Volunteers enable laboratory investigations to occur where more detailed measurements can be made, but may lack ecological validity where they do not accurately represent patients with physical or cognitive deficits. There is a lack of research conducted within outpatient and community settings, where healthcare practitioners need to adapt their practice to each patient's environment and the equipment they have available [61], and this needs to be addressed. Research conducted outside a laboratory environment can provide a more realistic and comprehensive account of the movements involved. However, there is less control over variables with a risk of losing some of the precision laboratory settings can provide. A limitation of research in clinical settings, especially for

kinematic analysis, is the lack of access to gold-standard measurement tools such as Vicon or OPTOTRAK [71]. Instead, research frequently used 2-D video, photographs and inclinometers. On balance, the findings from this scoping review identify there is a need for both research with the tools and control of the laboratory as well as the naturalistic setting of hospitals and the community.

#### Narrative literature

The remaining literature sources included within this scoping review mostly comprised narrative summaries of legislation and government statistics. The legislation surrounding moving and handling in the UK was last updated in 2002. NHS staff involved in moving and handling are required to complete training following principles of safe manual handling, however, the evidence behind the training is unclear [72]. In addition, there is no specific AHP training for correct moving and handling of patients in a therapeutic manner.

Literature originating from the US has a focus on SPHM programs with an emphasis placed on using assistive devices to aid manual patient handling. Programs based on assistance devices are not found universally across the US, however, with HCPs still required to manually move and handle patients for certain tasks or movements.

#### Gaps in the literature

Few studies have focused on AHPs performing manual patient handling despite these staff groups being likely to be involved with therapeutic handling practices which are manually intensive. There are no current guidelines or formal training provided to AHPs within the UK for correct therapeutic handling, and this review found no evidence that such training occurs in other geographical locations. This is therefore a priority area for future research. Despite manual handling being identified as a significant factor involved in development of WRMSD, a lack of research was identified within this scoping review.

The records identified in this scoping review demonstrate that there is a lack of detailed measurement of manual patient handling in healthcare settings with real patients, with many articles measuring trunk position or using video or photos for full body analysis. Measuring patient handling in laboratory settings is more accurate via access to 3D motion analysis systems. However, these often involve simulated patients and thus may not accurately reflect how healthcare practitioners move and handle real patients in the healthcare environment. Technology used to collect data in healthcare settings does not allow for the recording accuracy of laboratory-based systems, however, due to recent advances in technology this is an area that could now be explored further.

Few studies qualitatively assessed moving and handling of patients in healthcare. Ethnographic research and further investigation into perceptions and experiences of patient handling could allow for improvement of staff training and guidance to reduce risk of injury.

Of the records included in this scoping review, one primary research study was conducted within the UK. More research is required internationally, and especially in the UK, to inform this local context and to allow relevant evidence to help guide and improve staff training content or methods of teaching to reduce the risk of injury on HCPs.

#### Strengths and limitations

This scoping review followed a comprehensive search strategy and protocol that was reviewed by an experienced research team with previous experience in scoping reviews. There remains the possibility that some relevant articles were not included which is also likely to be the case as the search was restricted to the English language. The methodological quality of literature included in this scoping review was not assessed in keeping with methodological guidance for scoping reviews where the aim was to map the available literature rather than to assess its quality or the implication of study findings. Most of the included research was conducted in the US and Canada, therefore there may be difficulty with generalizing findings to other healthcare contexts.

#### Acknowledgements

Thanks to Susan Simpson, moving and handling facilitator for NHS Grampian for her expertise in legislation related to moving and handling in the UK. This scoping review was completed as part of a Doctorate of Physiotherapy degree and formed the first step in a program of research on manual patient handling in the UK.

*Ethical approval*: Ethical approval was not required for this study.

*Funding*: This research did not receive any specific grant from funding agencies in the public, commercial or not-for-profit sectors.

#### **Conflicts of interest**

The authors declare no conflict of interest.

#### Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at doi:10.1016/j.physio.2023. 06.003.

#### References

- [1] Schoenfisch AL, Kucera KL, Lipscomb HJ, McIlvaine J, Becherer L, James T, et al. Use of assistive devices to lift, transfer, and reposition hospital patients. Nurs Res 2019;68(1):3-12.
- Baptise A. An evaluation of nursing tasks. Work. 2011;40(2):115–24.
   Smith J, Alexander P, Johnson C, Nelson V. Guidance on manual handling in physiotherapy: 4th Edition. Chartered Society of Physiotherapy [Internet]. London: CSP; 2014. Available from: https://www.csp.org.uk/system/files/guidance\_manual\_ andling\_physiotherapy\_4thed\_2014.pdf [Accessed 20 July 2020].
- [4]. Health and Safety Executive. Work related musculoskeletal disorders in Great Britain (WRMSD). Health and Safety Executive [Internet]. Merseyside: HSE; 2019. Available from: https://www.hse.gov.uk/statistics/causdis/msd.pdf [Accessed 10 June 2020]
- [5] O'Sullivan P. It's time for change with the management of non-specific chronic low back pain. Br J Sports Med 2012:46:224-7.
- [6] Bernal D, Campos-Serna J, Tobias A, Vargas-Prada S, Benavides FG, Serra C. Workrelated psychosocial risk factors and musculoskeletal disorders in hospital nurses and nursing aides: a systematic review and meta-analysis. Int J Nurs Stud 2015:52(2):635-48.
- [7] Chartered Society of Physiotherapy. Fitness for Work: Physiotherapy Works. Chartered Society of Physiotherapy [Internet] London: CSP; 2016. Available from: https:// sp.org.uk/system/files/fitness\_for\_work\_physiotherapy\_works.pdf [Accessed March 20211
- [8] Occupational Safety and Health Administration, Worker Safety in Your Hospital; Know the Facts. Occupational Safety and Health Administration [Internet] Washington: OSHA; 2013. Available from: https://doi.org/10.1016/j.january.com/ highlights\_508.pdf [Accessed 12 May 2021].
- Anderson S, Stuckey R, Fortington LV, Oakman J. Workplace injuries in the Australian allied health workforce. Aust Health Rev 2019;43(1):49–54.
- [10] Ribeiro T, Serranheira F, Loureiro H. Work related musculoskeletal disorders in primary health care nurses. Appl Nurs Res 2017;33:72-7.
- [11] McCrory B, Burnfield JM, Darragh AR, Meza JL, Irons SL, Chernyavskiy P, et al. Work injuries among therapists in physical rehabilitation. Proc Hum Factors Ergon Soc Annu Meet 2014;58(1):1072-6.
- [12] Glover W, McGregor A, Sullivan C, Hague J. Work-related musculoskeletal disorders affecting members of the Chartered Society of Physiotherapy. Physiotherapy. 2005:91(3):138-47.
- [13] Saraceni N, Kent P, Ng L, Campbell A, Straker L, O'Sullivan P. To flex or not to flex? Is there a relationship between lumbar spine flexion during lifting and low back pain? A systematic review with meta-analysis, J Orthopaed Sports Phys Ther 2020;50(3):121–30.
- [14] National Institute for Occupational Safety and Health. Safe patient handling and mobility (SPHM). Centers for Disease Control and Prevention [Internet]. Atlanta: CDC; 2013. Available from: https://www.cdc.gov/niosh/topics/safepatient/default.html [Accessed 8 March 2021].
- [15] Peters MDJ, Godfrey C, McInerney P, Munn Z, Tricco AC, Khalil, H. Chapter 11: Scoping Reviews (2020 version). In: Aromataris E, Munn Z (Editors). JBI Manual for Evidence Synthesis. Adelaide: JBI; 2020. Available from: https://synthesismanual.jbi. obal [Accessed 29 August 2020].
- [16] Johnson KS, Cooper K, Swinton S, Pavlova A. Moving and handling of people in the healthcare setting: a scoping review protocol. Centre for Open Science [Internet]. Charlottesville: OSF; 2021. Available from: https://osf.io/rh9tj/ [Accessed 8 April 20221.
- [17] United Nations Development Programme, Latest Human Development Index Rating, United Nations Development Programme. United Nations Development Programme [Internet]. New York: UNDP; 2020. Available from: http://hdr.undp.org/en/o atest-human-development-index-ranking [Accessed 18 August 2020].
- [18] Health and Safety Executive. The Manual Handling Operations Regulations 1992 (as amended) (MHOR). Health and Safety Executive [Internet]. Merseyside: HSE; 2020. Available from: https://www.hse.gov.uk/foi/internalops/ocs/300-399/313\_5.htm [Accessed 2021 03 08].
- [19] Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. BMJ
- [20] Tullar JM, Brewer S, Amick III BC, Irvin E, Mahood Q, Pompeii LA, et al. Occupational safety and health interventions to reduce musculoskeletal symptoms in the health care sector. J Occup Rehabil 2010;20(2):199-219.
- [21] Brusco NK, Taylor N, Stevens JA, Butler M, Searle A. Development of a manual handling programme for allied health. Int J Ther Rehabil 2007;14(5):204-9.
- [22] Cantarella C, Stucchi G, Menoni O, Consonni D, Cairoli S, Manno R, et al. MAPO method to assess the risk of patient manual handling in hospital wards; a validation study. Hum Factors 2020;62(7):1141-9.
- [23] Carneiro P, Martins J, Torres M. Musculoskeletal disorder risk assessment in home care urses. Work. 2015;51(4):657-65.
- [24] Chen J, Daraiseh NM, Davis KG, Pan W. Sources of work-related acute fatigue in United States hospital nurses. Nurs Health Sci 2014;16(1):19-25.
- [25] Hodder JN, MacKinnon SN, Ralhan A, Keir PJ. Effects of training and experience on patient transfer biomechanics. Int J Ind Ergonomics 2010;40(3):282-8.
- [26] Hodder JN, Holmes MW, Keir PJ. Continuous assessment of work activities and posture in long-term care nurses. Ergonomics 2010;53(9):1097-107.
- [27] Holmes MWR, Hodder JN, Keir PJ. Continuous assessment of low back loads in longterm care nurses. Ergonomics 2010;53(9):1108-16.
- [28] Howard NL, Bao S, Kim H, Silverstein B. Comparison of Muscle Activity of Four Types of Bed-To-Wheelchair Transfers. Am J Safe Patient Handl Mov 2013;3(1):16-29.
- [29] Hye-Knudsen CT, Schibye B, Hjortskov N, Fallentin N. Trunk motion characteristics during different patient handling tasks. Int J Ind Ergon 2004;33(4):327-37.

- [30] Jordan C, Luttmann A, Theilmeir A, Kuhn S, Wortmann N, Jager M. Characteristic values of the lumbar load of manual patient handling for the application in workers' compensation procedures. J Occip Med Toxicol 2011;6(1):1-13.
- [31] Kang M, Jung D, An D, Yoo W, Oh J. Acute effects of hamstring-stretching exercises on the kinematics of the lumbar spine and hip during stoop lifting. J Back Musculoskelet Rehabil 2013;26(3):329–36.
- [32] Kim T, Roh H. Analysis of risk factors for work-related musculoskeletal disorders in radiological technologists. J Phys Ther Sci 2014;26(9):1423-8.
- [33] Kjellberg K, Lagerstrom M, Hagberg M. Work technique of nurses in patient transfer tasks and associations with personal factors. Scand J Work Environ Health 2003:29(6):468-77
- [34] Kurowski A, Buchholz B, Punnet L, ProCare, Research T. A physical workload index to evaluate a safe resident handling program for nursing home personnel. Hum Factors 2014;56(4):669-83.
- [35] Kyriakidis S, Stevens ML, Karstad K, Sogaard K, Holtermann A. The influence of nursing home, ward, and eldercare workers on the number of resident handlings performed per shift in eldercare. Int J Env Res Public Health 2021;18(21). Arte Number: 11040. ate of Pubaton: Noember-1 2021.
- [36] Larouche D, Bellemare M, Prairie J, Hegg-Deloye S, Corbeil P. Overall risk index for patient transfers in total assistance mode executed by medical technician-paramedics in real work situations. Appl Ergon 2019;74:177-85.
- [37] Larouche D, Corbeil P, Bellemare M, Authier M, Prairie J, Hegg-Deloye S. To what extent do paramedics apply safe patient handling principles when transferring patients from stair chairs to stretchers? Ergonomics. 2019;62(10):1313-26.
- [38] Maekawa Y, Shiozaki A, Majima Y. Investigation of the load on the lumbar region in nursing technique's movements - relation between twist and surface electromyogram. Stud Health Technol Inform 2009;146:460-4.
- [39] Skotte J, Fallentin N. Low back injury risk during repositioning of patients in bed: the influence of handling technique, patient weight and disability. Ergonomics 2008;51(7):1042-52.
- [40] Skotte JH, Essendrop M, Hansen AF, Schibye B. A dynamic 3D biomechanical evaluation of the load on the low back during different patient-handling tasks. J Biomech 2002;35(10):1357-66.
- [41] Stringer EJ, Rice MS. Perception, Experience, and Ground Reaction Forces when Performing and Assessing Pivot Transfers. Am J Safe Patient Handl Mov 2014:4(3):76-85.
- [42] Theis JL, Finkelstein MJ, Long-term effects of safe patient handling program on staff injuries. Rehab Nurs 2014;39(1):26-35.
- [43] Veira E, Kumar S. Safety analysis of patient transfers and handling tasks. Qual Saf Health Care 2009;18(5):380-4.
- [44] Arias OE, Umukoro PE, Stoffel SD, Hopcia K, Sorensen G, Dennerlein JT. Associations between trunk flexion and physical activity of patient care workers for a single shift: a pilot study. Work. 2017;56(2):247–55.
- [45] Fiedler KM, Weir PL, van Wyk PM, Andrews DM. Analyzing what nurses do during work in a hospital setting: a feasibility study using video. Work. 2012;43(4):515-23.
- [46] Fragala G. Facilitating repositioning in bed. AAOHN J 2011;59(2):63–8.
- [47] Garzillo EM, Monaco MGL, Corvino AR, D'ancicco F, Feola D, Ventura DD, et al. Healthcare workers and manual patient handling: A pilot study for interdisciplinary training. Int J Environ Res Public Health 2020;17:1-14.
- [48] Newton K, Quinton A, Childs J. The incidence of Australian private practice sonographers moving patients unassisted and their level of training: A pilot study. Sonography 2020:7(2):48–54.
- [49] de Ruiter HP, Liaschenko J. To lift or not to lift: patient-handling practices. AAOHN J 2011;59(8):337-43.
- [50] Osborne ARH, Connell C, Morphet J. Investigating emergency nurses' beliefs and experiences with patient handling in the emergency department. Australas Emerg Care 2021;24(1):49-54.
- [51] Wångblad C, Ekblad M, Wijk H, Ivanoff SD. Experiences of physical strain during person transfer situations in dementia care units. Scand J Caring Sci 2009:23(4):644-50
- [52] McCoskey KL, Ergonomics and patient handling, AAOHN J 2007;55(11):454-62.
- [53] Lavender SA, Nagavarapu S, Marras WS. Biomechanical loads on the spine as patients are prepared for mechanical transfers: have patient lifts completely solved the problem? J Orthop Res Conf 2016;34(1):1.
- [54] Nikolajsen H, Nielsen CA. Manual patient handling and rehabilitation in health care settings: an observational study. Physiotherapy 2015;101:eS1092.
- [55] Wade H, Kneafsey R, Carpenter C. An ethnographic study of physiotherapists and occupational therapists experiences of implementing guidance on safe manual handling practice in a community setting. Clin Rehabil 2017;31(3):417–8.
- [56] Apple B, Letvak S. Ergonomic challenges in the perioperative setting. AORN J 2021;113(4):339-48.
- [57] Haney LL. Physical stresses related to the safe handling of residents. Director 2003;11(4):151-3.
- [58] Hignett S, Fray M, Rossi MA, Tamminen PL, Hermann S, Lomi C, et al. Implementation of the Manual Handling Directive in the healthcare industry in the European union for patient handling tasks. Int J Ind. Ergonomics. 2007;37(5):415–23.
- [59] Johnstone J. Manual handling: the challenges of different care environments. Br J Nurs 2020;29(6):358-63.
- [60] Rinds G. Manual handling in healthcare support work. BJHCA 2008;2(1):11-6.
- [61] Tofts D, Arnold M. Moving and handling in the community: update on legislation and best practice. Br J Community Nurs 2012;17(2):50-7.
- [62] Vatwani A. Caregiver guide and instructions for safe bed mobility. Arch Phys Med Rehabil 2017:98(9):1907-10
- [63] Waters T, Baptiste A, Short M, Plante-Mallon L, Nelson A. AORN ergonomic tool 1: Lateral transfer of a patient from a stretcher to an OR bed. AORN J 2011;93(3):334-9.
- [64] Waters T, Short M, Lloyd J, Baptiste A, Butler L, Petersen C, et al. AORN ergonomic tool 2: positioning and repositioning the supine patient on the OR bed. AORN J 2011;93(4):445-9.

- [65] Waters TR, Nelson A, Proctor C. Patient Handling Tasks with High Risk for Musculoskeletal Disorders in Critical Care. Crit Care Nurs Clin North Am 2007:19(2):131-43.
- [66] Weiner C, Alperovitch-Najenson D, Ribak J, Kalichman L. Prevention of nurses' workrelated musculoskeletal disorders resulting from repositioning patients in bed: comprehensive narrative review, Workplace Health Saf 2015;63(5):226–32.
- [67] Weinmeyer R. Safe patient handling laws and programs for health care workers. AMA J Ethics 2016;18(4):416–21.
- [68] Mann CJ. Observational research methods. Research design II: cohort, cross sectional, and case-control studies. Emerg Med J 2003;20(1):54-60.
- [69] Corazza S, Mündermann L, Gambaretto E. Markerless motion capture through visual hull, articulated icp and subject specific model generation. Int J Comput Vis 2010;87:156-69.
- [70] Serranheira F, Sousa-Uva M, Sousa-Uva A. Hospital nurses tasks and work-related
- musculoskeletal disorders symptoms: a detailed analysis. Work. 2010;15:401–9.

  [71] Cuesta-Vargas A, Galán-Mercant A, Williams JM. The use of inertial sensors system for human motion analysis. Phys Ther Rev 2010;15(6):462–73.
- [72] Haslam C, Clemes S, McDermott. Manual handling training: Investigation of current practices and development of guidelines. Health and Safety Executive [Internet]. Merseyside: HSE; 2007. Available from: https://www.hse.gov.uk/research/rrpdf/rr583. pdf [Accessed 14 May 2021].

Available online at www.sciencedirect.com

### **ScienceDirect**

## Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) Checklist

SECTION	ITEM	PRISMA-ScR CHECKLIST ITEM	REPORTED ON PAGE #
TITLE			
Title	1	Identify the report as a scoping review.	Title page
Structured summary	2	Provide a structured summary that includes (as applicable): background, objectives, eligibility criteria, sources of evidence, charting methods, results, and conclusions that relate to the review questions and objectives.	Abstract
INTRODUCTION			ı
Rationale	3	Describe the rationale for the review in the context of what is already known. Explain why the review questions/objectives lend themselves to a scoping review approach.	1-2
Objectives	4	Provide an explicit statement of the questions and objectives being addressed with reference to their key elements (e.g., population or participants, concepts, and context) or other relevant key elements used to conceptualize the review questions and/or objectives.	2
METHODS		<u> </u>	
Protocol and registration	5	Indicate whether a review protocol exists; state if and where it can be accessed (e.g., a Web address); and if available, provide registration information, including the registration number.	3-4
Eligibility criteria	6	Specify characteristics of the sources of evidence used as eligibility criteria (e.g., years considered, language, and publication status), and provide a rationale.	3-4
Information sources*	7	Describe all information sources in the search (e.g., databases with dates of coverage and contact with authors to identify additional sources), as well as the date the most recent search was executed.	4
Search	8	Present the full electronic search strategy for at least 1 database, including any limits used, such that it could be repeated.	Supplementary materials
Selection of sources of evidence†	9	State the process for selecting sources of evidence (i.e., screening and eligibility) included in the scoping review.	4
Data charting process‡	10	Describe the methods of charting data from the included sources of evidence (e.g., calibrated forms or forms that have been tested by the team before their use, and whether data charting was done independently or in duplicate) and any processes for obtaining and confirming data from investigators.	4-5
Data items	11	List and define all variables for which data were sought and any assumptions and simplifications made.	5
Critical appraisal of individual sources of evidence§	12	If done, provide a rationale for conducting a critical appraisal of included sources of evidence; describe the methods used and how this information was used in any data synthesis (if appropriate).	N/A

SECTION	ITEM	PRISMA-ScR CHECKLIST ITEM	REPORTED ON PAGE #
Synthesis of results	13	Describe the methods of handling and summarizing the data that were charted.	6-8
RESULTS			
Selection of sources of evidence	14	Give numbers of sources of evidence screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally using a flow diagram.	6
Characteristics of sources of evidence	15	For each source of evidence, present characteristics for which data were charted and provide the citations.	6
Critical appraisal within sources of evidence	16	If done, present data on critical appraisal of included sources of evidence (see item 12).	N/A
Results of individual sources of evidence	17	For each included source of evidence, present the relevant data that were charted that relate to the review questions and objectives.	6-8
Synthesis of results	18	Summarize and/or present the charting results as they relate to the review questions and objectives.	6-8
DISCUSSION			
Summary of evidence	19	Summarize the main results (including an overview of concepts, themes, and types of evidence available), link to the review questions and objectives, and consider the relevance to key groups.	8-11
Limitations	20	Discuss the limitations of the scoping review process.	12
Conclusions	21	Provide a general interpretation of the results with respect to the review questions and objectives, as well as potential implications and/or next steps.	10-12
FUNDING			
Funding	22	Describe sources of funding for the included sources of evidence, as well as sources of funding for the scoping review. Describe the role of the funders of the scoping review.	12

JBI = Joanna Briggs Institute; PRISMA-ScR = Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews.

19 20 21

<sup>\*</sup> Where sources of evidence (see second footnote) are compiled from, such as bibliographic databases, social media platforms, and Web sites.

<sup>†</sup> A more inclusive/heterogeneous term used to account for the different types of evidence or data sources (e.g., quantitative and/or qualitative research, expert opinion, and policy documents) that may be eligible in a scoping review as opposed to only studies. This is not to be confused with *information sources* (see first footnote).

<sup>‡</sup> The frameworks by Arksey and O'Malley (6) and Levac and colleagues (7) and the JBI guidance (4, 5) refer to the process of data extraction in a scoping review as data charting.

<sup>§</sup> The process of systematically examining research evidence to assess its validity, results, and relevance before using it to inform a decision. This term is used for items 12 and 19 instead of "risk of bias" (which is more applicable to systematic reviews of interventions) to include and acknowledge the various sources of evidence that may be used in a scoping review (e.g., quantitative and/or qualitative research, expert opinion, and policy document).

### 23 Supplementary material – Full search strategy

- 24 Key: TX All Text, KW Keyword, MH Exact Subject Heading
- 25 AMED (via EBSCOhost)
- 26 Search conducted 12 August 2020

Search	Query	Records retrieved
#1	TX moving and handling OR KW manual handling OR TX manual handling OR KW (moving and lifting patients) OR TX (moving and lifting patients) OR TX patient handling OR TX patient moving and handling OR TX patient positioning	399
#2	KW nurse OR TX nurs* OR KW physiotherapy OR KW physiotherapist TX physiotherap* OR KW physical therapy OR TX physical therap* OR KW allied health personnel OR TX allied health* OR TX support staff OR KW rehabilitation OR TX rehabilitation OR KW occupational therapist OR KW occupational therapy OR TX occupational therap*	127601
#3	#1 AND #2	319
Limited t	o English language	

### 28 AMED (via EBSCOhost)

### Updated search conducted 10 November 2021

Search	Query	Records retrieved
#1	TX moving and handling OR KW manual handling OR TX manual handling OR KW (moving and lifting patients) OR TX (moving and lifting patients) OR TX patient handling OR TX patient moving and handling OR TX patient positioning	408
#2	KW nurse OR TX nurs* OR KW physiotherapy OR KW physiotherapist TX physiotherap* OR KW physical therapy OR TX physical therap* OR KW allied health personnel OR TX allied health* OR TX support staff OR KW rehabilitation OR TX rehabilitation OR KW occupational therapist OR KW occupational therapy OR TX occupational therap*	131747
#3	#1 AND #2	347
#4	#1 AND #2 (Limited to literature since August 2020)	0
Limited t	to English language	

### 33 CINAHL (via EBSCOhost)

Search	Query	Records retrieved
#1	TX moving and handling OR TX patient moving and handling OR MH patient positioning OR TX patient positioning OR TX transfer* technique* OR TX manual handling OR TX moving and lifting patients	24460
#2	MH nurses OR TX nurs* OR TX physiotherap* OR MH physical therapy OR TX physical therap* OR MH rehabilitation OR TX rehabilitation OR MH allied health personnel OR TX allied health* OR TX support staff OR MH occupational therapy OR TX occupational therap*	3052157
#3	#1 AND #2	17814
#4	TX guidance OR TX guidelines OR TX practice guidelines OR TX clinical practice guidelines OR TX protocol OR TX regulation	800153
#5	#3 AND #4	8317
#6	MH ergonomics OR TX ergonomic* OR TX human factors	19336
#7	#5 AND #6	1304
Limited t	o English language	

### 37 EMBASE (via Ovid)

### 38 Search conducted 12 August 2020

Search	Query	Records retrieved
#1	KW moving and handling OR TX moving and handling OR KW moving and lifting patients OR TX moving and lifting patients OR KW manual handling OR TX manual handling OR KW patient handling OR TX patient handling OR KW patient moving and handling OR TX patient moving and handling OR KW patient positioning OR TX patient positioning	9542
#2	KW nurse OR TX nurs* OR KW physiotherapist OR KW physiotherapy OR TX physiotherap* OR KW physical therapy OR KW physical therapist OR TX physical therap* OR KW allied health personnel OR KW allied health professional OR TW allied health* OR KW support staff OR TX support staff OR KW rehabilitation OR TX rehabilitation OR KW occupational therapist OR KW occupational therapy OR TX occupational therap*	805052
#3	#1 AND #2	892

### 40 EMBASE (via Ovid)

### 41 Updated search conducted 10 November 2021

Search	Query	Records retrieved
#1	KW moving and handling OR TX moving and handling OR KW moving and lifting patients OR TX moving and lifting patients OR KW manual handling OR TX manual handling OR KW patient handling OR TX patient handling OR KW patient moving and handling OR TX patient moving and handling OR KW patient positioning OR TX patient positioning	7765
#2	KW nurse OR TX nurs* OR KW physiotherapist OR KW physiotherapy OR TX physiotherap* OR KW physical therapy OR KW physical therapist OR TX physical therap* OR KW allied health personnel OR KW allied health professional OR TW allied health* OR KW support staff OR TX support staff OR KW rehabilitation OR TX rehabilitation OR KW occupational therapist OR KW occupational therapy OR TX occupational therap*	879357
#3	#1 AND #2	881
#3	#1 AND #2 (Limited to literature since August 2020)	112

### 44 MEDLINE (via EBSCOhost)

### 45 Search conducted 12 August 2020

Search	Query	Records retrieved
#1	TX moving and handling OR TX moving and lifting patients OR TX manual handling OR TX patient handling OR TX patient moving and handling OR MH patient positioning OR TX patient positioning	16766
#2	MH nurses OR TX nurs* OR TX physiotherap* OR TX physical therap* OR MH allied health personnel OR TX allied health* OR TX support staff OR MH rehabilitation OR TX rehabilitation OR MH occupational therapy OR TX occupational therap*	1773156
#3	#1 AND #2	2735
Limited t	o English language	

### 47 MEDLINE (via EBSCOhost)

### Updated search conducted 10 November 2021

Search	Query	Records retrieved
#1	TX moving and handling OR TX moving and lifting patients OR TX manual handling OR TX patient handling OR TX patient moving and handling OR MH patient positioning OR TX patient positioning	19413
#2	MH nurses OR TX nurs* OR TX physiotherap* OR TX physical therap* OR MH allied health personnel OR TX allied health* OR TX support staff OR MH rehabilitation OR TX rehabilitation OR MH occupational therapy OR TX occupational therap*	1979659
#3	#1 AND #2	3725
#3	#1 AND #2 (Limited to literature since August 2020)	551
Limited t	o English language	

### 49

48

### 50 SPORTDiscus (via EBSCOhost)

Search	Query	Records retrieved
#1	KW moving and handling OR TX moving and handling OR KW patient	8977
	moving and handling OR TX patient moving and handling OR KW moving	
	and lifting patients OR TX moving and lifting patients OR KW manual	
	handling OR TX manual handling OR KW patient handling OR TX patient	
	handling OR KW patient positioning OR TX patient positioning OR TX	
	transfer* technique*	

#2	KW nurse OR TX nurs* OR KW physiotherapist OR KW physiotherapy OR TX physiotherap* OR KW physical therapist OR KW physical therapy OR TX physical therap* OR KW allied health personnel OR KW allied health professional OR TX allied health* OR KW support staff OR TX support staff OR KW rehabilitation OR TX rehabilitation OR KW occupational therapy OR	242724
#3	KW occupational therapist OR TX occupational therap*  #1 AND #2	3144
#3	#1 AND #2 (Limited to literature published after 2002)	2725
Limited to English language		

### 53 Google Scholar

### 54 Search conducted 10 November 2021

Search	Query	Records retrieved
#1	"Patient handling" AND "healthcare"	150
#2	"Moving and handling" AND "healthcare"	150
#3	"Moving and handling" AND "patients"	150
#4	"Manual handling" AND "healthcare"	150
#5	"Manual handling" AND "patients"	150
Records h	nand selected for screening	

55

57

### 56 ETHOS

Search	Query	Records retrieved
#1	Moving and handling in healthcare	10
#2	Moving and handling of patients	14
#3	Manual handling in healthcare	12
#4	Manual handling of patients	7
Records h	and selected for screening	

### 59 Open Grey

### 60 Search conducted 10 November 2021

Search	Query	Records retrieved
#1	Moving and handling in healthcare	1
#2	Moving and handling of patients	2
#3	Manual handling in healthcare	1
#4	Manual handling of patients	1

61

### 62 HSE

Search	Query	Records retrieved
#1	Moving and handling in healthcare	260

Records hand selected for screening	

### 65 NIOSH

### 66 Search conducted 10 November 2021

Search	Query	Records retrieved
#1	Moving and handling in healthcare	390
Records h	and selected for screening	

67

### 68 Safe Work Australia

### 69 Search conducted 10 November 2021

Search	Query	Records retrieved
#1	Moving and handling in healthcare	236
#2	Safe patient handling	715
#3	Moving and handling of patients	234
Records h	and selected for screening	

70

### 71 Canadian Centre of Occupational Health and Safety

Search	Query	Records retrieved
#1	Moving and handling in healthcare	3986

#2	Safe patient handling	3014
#3	Moving and handling of patients	3983
Records h	and selected for screening	

### 74 Worksafe New Zealand

Search	Query	Records retrieved
#1	Moving and handling in healthcare	1000
#2	Safe patient handling	1000
#3	Moving and handling of patients	1000
Records h	nand selected for screening	

#### 78 Excluded studies

- 79 1. Parkes KR, Carnell S, Farmer E. Musculo-skeletal disorders, mental
- health and the work environment. Health and Safety Executive
- [Internet]. Merseyside: HSE; 2005 [cited 2020 08 24]. [Available from:
- https://www.hse.gov.uk/research/rrpdf/rr316.pdf].
- Reason for exclusion: Ineligible setting.
- 2. Gillianders T. Upper Limb Disorders in the Workplace. Health & Safety at
- 85 Work. 2004;26(2):33.
- Reason for exclusion: Ineligible setting.
- 3. Health and Safety Executive. Manual handling at work: A brief guide.
- Health and Safety Executive [Internet]. Merseyside: HSE; 2020 [cited
- 89 2020 08 24]. [Available from:
- 90 https://www.hse.gov.uk/pubns/indg143.pdf].
- 91 Reason for exclusion: Ineligible concept.
- 92 4. Munro KD. Clinical reasoning for manual handling risk assessments in
- community settings: moving from rule based to intuitive reasoning
- 94 [dissertation]. University of Strathclyde; 2017.
- 95 Reason for exclusion: Ineligible concept.
- 96 5. Holtermann A, Clausen T, Jørgensen MB, Burdorf A, Andersen LL. Patient
- 97 Handling and Risk for Developing Persistent Low-Back Pain among
- 98 Female Healthcare Workers. Scandinavian journal of work, environment
- 99 & health. 2013;39(2):164-169.
- 100 Reason for exclusion: Ineligible concept.
- 101 6. Dorothée M, Benchekroun TH, Falzon P, Vidal-Gomel C. Patient handling
- as an healthcare activity: a simulation-based training methodology.
- 103 HEPS 2016. 2016;1-9.
- 104 Reason for exclusion: Ineligible concept.
- 105 7. Hignett S, Fray M. Manual handling in healthcare. Proceedings of the 1st
- 106 Conference of the Federation of the European Ergonomics Societies
- 107 [FEES]. 2010;1-10.
- 108 Reason for exclusion: Ineligible concept.
- 109 8. Waters TR. Introduction to Ergonomics for Healthcare Workers.
- 110 Rehabilitation Nursing. 2010;(5):185-191.
- 111 Reason for exclusion: Ineligible concept.

- 9. Owen BD, Keene K, Olson S. An ergonomic approach to reducing
- back/shoulder stress in hospital nursing personnel: a five year follow up.
- 114 Int J Nurs Stud. 2002;39(3):295-302.
- 115 Reason for exclusion: Ineligible concept.
- 116 10. Smedley J, Inskip H, Trevelyan F, Buckle P, Cooper C, Coggon D. Risk
- factors for incident neck and shoulder pain in hospital nurses. Occup
- 118 Environ Med. 2003;60(11):864-869.
- 119 Reason for exclusion: Ineligible concept.
- 120 11. Smedley J, Trevelyan F, Inskip H, Buckle P, Cooper C, Coggon D. Impact
- of ergonomic intervention on back pain among nurses. Scandinavian
- Journal of Work, Environment and Health. 2003;29(2):117-123.
- Reason for exclusion: Ineligible concept.
- 124 12. De Castro AB. Handle with care: The American nurses association's
- campaign to address work-related musculoskeletal disorders. Orthop
- 126 Nurs. 2006;25(6):356-365.
- 127 Reason for exclusion: Ineligible concept.
- 13. Guthrie PF, Westphal L, Dahlman B, Berg M, Behnam K, Ferrell D. A
- patient lifting intervention for preventing the work-related injuries of
- nurses. Work. 2004;22(2):79-88.
- 131 Reason for exclusion: Ineligible concept.
- 132 14. Radovanovic CAT, Alexandre NMC. Validation of an instrument for patient
- handling assessment. Appl Ergon. 2004;35(4):321-328.
- 134 Reason for exclusion: Ineligible concept.
- 135 15. Hignett S. Systematic review of patient handling activities starting in
- lying, sitting and standing positions. J Adv Nurs. 2003;41(6):545-552.
- 137 Reason for exclusion: Ineligible concept.
- 138 16. Mutch K. Changing manual-handling practice in a stroke rehabilitation
- unit. Prof Nurse. 2004;19(7):374-378.
- 140 Reason for exclusion: Ineligible concept.
- 141 17. Menzel NN, Brooks SM, Bernard TE, Nelson A. The physical workload of
- nursing personnel: Association with musculoskeletal discomfort. Int J
- 143 Nurs Stud. 2004;41(8):859-867.
- 144 Reason for exclusion: Ineligible concept.
- 145 18. Trossman S. Protecting the next generation: ANA, nursing partners work
- to educate students on safe patient handling techniques. Am Nurse.

- 147 2004;36(5):1, 7-9.
- 148 Reason for exclusion: Ineligible population.
- 149 19. Yip VYB. New low back pain in nurses: Work activities, work stress and
- sedentary lifestyle. J Adv Nurs. 2004;46(4):430-440.
- 151 Reason for exclusion: Ineligible concept.
- 20. Kjellberg K, Lagerstrom M, Hagberg M. Patient safety and comfort during
- transfers in relation to nurses' work technique. J Adv Nurs.
- 154 2004;47(3):251-259.
- 155 Reason for exclusion: Ineligible concept.
- 156 21. Menzel NN. Back pain prevalence in nursing personnel: measurement
- issues. AAOHN J. 2004;52(2):54-65.
- 158 Reason for exclusion: Ineligible concept.
- 159 22. Hartvigsen J, Lauritzen S, Lings S, Lauritzen T. Intensive education
- combined with low tech ergonomic intervention does not prevent low
- back pain in nurses. Occup Environ Med. 2005;62(1):13-17.
- 162 Reason for exclusion: Ineligible concept.
- 163 23. Pellatt GC. The safety and dignity of patients and nurses during patient
- handling. Br J Nurs. 2005;14(21):1150-1156.
- 165 Reason for exclusion: Ineligible concept.
- 166 24. Mitchell EA, Conlon AM, Armstrong M, Ryan AA. Towards rehabilitative
- handling in caring for patients following stroke: A participatory action
- research project. J Clin Nurs. 2005;14(3):3-12.
- 169 Reason for exclusion: Ineligible concept.
- 170 25. Tabone S. Safe patient handling. Tex Nurs. 2005;79(3):10-11.
- 171 Reason for exclusion: Ineligible concept.
- 172 26. Nelson A, Matz M, Chen F, Siddharthan K, Lloyd J, Fragala G.
- Development and evaluation of a multifaceted ergonomics program to
- prevent injuries associated with patient handling tasks. Int J Nurs Stud.
- 175 2006;43(6):717-733.
- 176 Reason for exclusion: Ineligible concept.
- 177 27. Waters T, Collins J, Galinsky T, Caruso C. NIOSH research efforts to
- prevent musculoskeletal disorders in the healthcare industry. Orthop
- 179 Nurs. 2006;25(6):380-389.
- 180 Reason for exclusion: Ineligible concept.

- 28. Castledine G. Nurse's poor moving and handling technique put patients
- at risk. Br J Nurs. 2006;15(6):307.
- 183 Reason for exclusion: Ineligible concept.
- 184 29. McKeon E. The ANA seeks investment in nursing workforce development:
- Safe patient handling and staffing at the forefront. Am J Nurs.
- 186 2006;106(5):79.
- 187 Reason for exclusion: Ineligible concept.
- 30. Smith DR, Mihashi M, Adachi Y, Koga H, Ishitake T. A detailed analysis of
- musculoskeletal disorder risk factors among Japanese nurses. J Saf Res.
- 190 2006;37(2):195-200.
- 191 Reason for exclusion: Ineligible concept,
- 192 31. Dawson AP, McLennan SN, Schiller SD, Jull GA, Hodges PW, Stewart S.
- 193 Interventions to prevent back pain and back injury in nurses: A
- systematic review. Occup Environ Med. 2007;64(10):642-650.
- 195 Reason for exclusion: Ineligible concept.
- 196 32. Pompeii LA, Lipscomb HJ, Dement JM. Surveillance of musculoskeletal
- injuries and disorders in a diverse cohort of workers at a tertiary care
- medical center. Am J Ind Med. 2008;51(5):344-356.
- 199 Reason for exclusion: Ineligible concept.
- 33. Warming S, Ebbehoj NE, Wiese N, Larsen LH, Duckert J, Tonnesen H.
- 201 Little effect of transfer technique instruction and physical fitness training
- in reducing low back pain among nurses: A cluster randomised
- 203 intervention study. Ergonomics. 2008;51(10):1530-1548.
- 204 Reason for exclusion: Ineligible concept.
- 205 34. Alexander P. The changing face of manual handling in the community. Br
- 206 J Community Nurs. 2008;13(7):316-322.
- 207 Reason for exclusion: Ineligible concept.
- 35. Hignett S, Crumpton E. Competency-based training for patient handling.
- 209 Appl Ergon. 2007;38(1):7-17.
- 210 Reason for exclusion: Ineligible concept.
- 211 36. Haney LL, Wright L. Sustaining Staff Nurse Support for a Patient Care
- 212 Ergonomics Program in Critical Care. Crit Care Nurs Clin North Am.
- 213 2007;19:197-204.
- 214 Reason for exclusion: Ineligible concept.

- 215 37. Knibbe HJJ, Knibbe NE, Klaassen, A. J. W. M. Safe Patient Handling
- 216 Program in Critical Care Using Peer Leaders: Lessons Learned in The
- Netherlands. Crit Care Nurs Clin North Am. 2007;19:205-211.
- 218 Reason for exclusion: Ineligible concept.
- 38. Pellatt GC. Clinical skills: bed making and patient positioning. Br J Nurs.
- 220 2007;16(5):302-305.
- Reason for exclusion: Ineligible concept.
- 39. Morse T, Fekieta R, Rubenstein H, Warren N, Alexander D, Wawzyniecki
- P. "Doing the heavy lifting: health care workers take back their backs".
- New solutions: a journal of environmental and occupational health policy.
- 225 2008;18(2):207-219.
- Reason for exclusion: Ineligible concept.
- 40. Pompeii LA, Lipscomb HJ, Schoenfisch AL, Dement JM. Musculoskeletal
- injuries resulting from patient handling tasks among hospital workers.
- 229 Am J Ind Med. 2009;52(7):571-578.
- 230 Reason for exclusion: Ineligible concept.
- 41. Akebi T, Inoue M, Harada N. Effects of educational intervention on joint
- angles of the trunk and lower extremity and on muscle activities during
- patient-handling tasks. Environmental Health and Preventive Medicine.
- 234 2009;14(2):118-127.
- 235 Reason for exclusion: Ineligible population.
- 42. Svensson AL, Stroyer J, Ebbehoj NE, SchultzLarsen K, Marott JL,
- Mortensen OS, et al. Multidimensional intervention and sickness absence
- in assistant nursing students. Occup Med. 2009;59(8):563-569.
- 239 Reason for exclusion: Ineligible population.
- 43. Sedlak CA, Doheny MO, Nelson A, Waters TR. Development of the
- National Association of Orthopaedic Nurses guidance statement on safe
- patient handling and movement in the orthopaedic setting. Orthop Nurs.
- 243 2009;28(2):2-8.
- 244 Reason for exclusion: Ineligible concept.
- 245 44. Kutash M, Short M, Shea J, Martinez M. The lift team's importance to a
- successful safe patient handling program. J Nurs Adm. 2009;39(4):170-
- 247 175.
- 248 Reason for exclusion: Ineligible concept.

- 45. Barnes AF. Reducing the incidence of back pain: student nurses'
- 250 recommendations. Br J Nurs. 2009;18(21):1334-1338.
- 251 Reason for exclusion: Ineligible population.
- 46. Knoblauch MD, Bethel SA. Safe patient-handling program "UPLIFTs"
- 253 nurse retention. Nursing. 2010;40(2):67-68.
- 254 Reason for exclusion: Ineligible concept.
- 255 47. Zadvinskis IM, Salsbury SL. Effects of a multifaceted minimal-lift
- environment for nursing staff: Pilot results. West J Nurs Res.
- 257 2010;32(1):47-63.
- 258 Reason for exclusion: Ineligible concept.
- 48. Sachs A. Nurses, patients alike benefit from safe patient handling. Am
- 260 Nurse. 2010;42(5):9.
- Reason for exclusion: Ineligible concept.
- 49. Cornish J, Jones A. Factors affecting compliance with moving and
- 263 handling policy: Student nurses' views and experiences. Nurse Education
- in Practice. 2010;10(2):96-100.
- 265 Reason for exclusion: Ineligible population.
- 50. Lee SJ, Faucett J, Gillen M, Krause N, Landry L. Factors associated with
- safe patient handling behaviors among critical care nurses. Am J Ind
- 268 Med. 2010;53(9):886-897.
- Reason for exclusion: Ineligible concept.
- 51. Waters TR, Rockefeller K. Safe patient handling for rehabilitation
- professionals. Rehabilitation nursing: the official journal of the
- Association of Rehabilitation Nurses. 2010;35(5):216-222.
- 273 Reason for exclusion: Ineligible concept.
- 52. Szeto PYG, Law RKY, Lee EWC, Chan SY, Lau TFO, Law SW. Effectiveness
- of an ergonomic intervention program for reducing work-related
- 276 musculoskeletal disorders in community nurses. Physiotherapy (United
- 277 Kingdom). Conference: World Physical Therapy 2011. Amsterdam
- Netherlands. Conference Publication. 2011;97 (SUPPL. 1):eS1201.
- 279 Reason for exclusion: Ineligible concept.
- 53. Genevay S, Cedraschi C, Courvoisier DS, Perneger TV, Grandjean R,
- Griesser AC, et al. Work related characteristics of back and neck pain
- among employees of a Swiss University Hospital. Joint Bone Spine.

- 283 2011;78(4):392-397.
- 284 Reason for exclusion: Ineligible concept.
- 285 54. Ogg MJ. Introduction to the Safe Patient Handling and Movement Series.
- 286 AORN J. 2011;93(3):331-333.
- 287 Reason for exclusion: Ineligible concept.
- 288 55. Schoenfisch AL, Lipscomb HJ, Myers DJ, Fricklas E, James T. A lift assist
- team in an acute care hospital-prevention of injury or transfer of risk
- during patient-handling tasks?. AAOHN J. 2011;59(8):329-334.
- 291 Reason for exclusion: Ineligible concept.
- 292 56. Mason MC. Back support. Nursing standard. 2011;25(30):23.
- 293 Reason for exclusion: Ineligible concept.
- 57. Black TR, Shah SM, Busch AJ, Metcalfe J, Lim HJ. Effect of transfer,
- 295 lifting, and repositioning (TLR) injury prevention program on
- 296 musculoskeletal injury among direct care workers. Journal of
- Occupational and Environmental Hygiene. 2011;8(4):226-235.
- 298 Reason for exclusion: Ineligible concept.
- 58. Kay K, Glass N. Debunking the manual handling myth: An investigation
- of manual handling knowledge and practices in the Australian private
- 301 health sector. Int J Nurs Pract. 2011;17(3):231-237.
- Reason for exclusion: Ineligible concept.
- 303 59. Griffiths H. Adverse risk: A 'dynamic interaction model of patient moving
- and handling'. J Nurs Manag. 2012;20(6):713-736.
- Reason for exclusion: Ineligible concept.
- 306 60. Callison MC, Nussbaum MA. Identification of physically demanding
- patient-handling tasks in an acute care hospital. Int J Ind Ergonomics.
- 308 2012;42(3):261-267.
- Reason for exclusion: Ineligible concept.
- 310 61. Krill C, Staffileno BA, Raven C. Empowering staff nurses to use research
- to change practice for safe patient handling. Nurs Outlook.
- 312 2012;60(3):157-162.
- Reason for exclusion: Ineligible concept.
- 314 62. Garg AG, Kapellusch JMK. Ergonomic interventions in nursing facilities:
- Long-term effectiveness of a comprehensive program. Occupational and
- 316 Environmental Medicine. Conference: 23rd Conference on Epidemiology
- in Occupational Health, EPICOH 2013: Improving the Impact. Utrecht

- Netherlands. Conference Publication. 2013;70 (SUPPL. 1):114.
- Reason for exclusion: Ineligible concept.
- 320 63. Burnfield JM, McCrory B, Shu Y, Buster TW, Taylor AP, Goldman AJ.
- Comparative kinematic and electromyographic assessment of clinician-
- and device-assisted sit-to-stand transfers in patients with stroke. Phys
- 323 Ther. 2013;93(10):1331-1341.
- Reason for exclusion: Ineligible population.
- 325 64. Lee SJ, Faucett J, Gillen M, Krause N. Musculoskeletal pain among
- critical-care nurses by availability and use of patient lifting equipment:
- 327 An analysis of cross-sectional survey data. Int J Nurs Stud.
- 328 2013;50(12):1648-1657.
- Reason for exclusion: Ineligible concept.
- 330 65. AlEisa E, AlAbbad H. Occupational back pain among rehabilitation nurses
- in Saudi Arabia: The influence of knowledge and awareness. Workplace
- 332 Health and Safety. 2013;61(9):401-407.
- Reason for exclusion: Ineligible concept.
- 334 66. Geiger JS. Establishing a physical therapist-driven model of safe patient
- handling and movement programs in a general hospital. Work.
- 336 2013;45(2):147-160.
- Reason for exclusion: Ineligible concept.
- 338 67. Campo M, Shiyko MP, Margulis H, Darragh AR. Effect of a safe patient
- handling program on rehabilitation outcomes. Arch Phys Med Rehabil.
- 340 2013;94(1):17-22.
- Reason for exclusion: Ineligible concept.
- 342 68. Belbeck A, Cudlip AC, Dickerson CR. Assessing the interplay between the
- 343 shoulders and low back during manual patient handling techniques in a
- nursing setting. JOSE. 2014;20(1):127-137.
- Reason for exclusion: Ineligible population.
- 346 69. Mayeda-Letourneau J. Safe patient handling and movement: A literature
- 347 review. Rehabil Nurs. 2014;39(3):123-129.
- Reason for exclusion: Ineligible concept.
- 70. Anderson MP, Carlisle S, Thomson C, Ross C, Reid HJ, Hart ND, et al.
- 350 Safe moving and handling of patients: an interprofessional approach.
- 351 Nursing standard. 2014;28(46):37-41.
- Reason for exclusion: Ineligible population.

- 353 71. Beauvais A, Frost L. Saving our backs: safe patient handling and mobility 354 for home care. Home Healthc Nurse. 2014;32(7):430-6.
- 355 Reason for exclusion: Ineligible concept.
- 356 72. Vroonland C, Baank S, Stokkel M. To manual handle, or to NOT manual handle? European Journal of Nuclear Medicine and Molecular Imaging.
- 358 Conference: 28th Annual Congress of the European Association of
- Nuclear Medicine, EANM 2015. Hamburg Germany. Conference
- 360 Publication. 42(1 SUPPL. 1):S847-S848.
- Reason for exclusion: Ineligible concept.
- 362
   73. Sundelin G, HarmsRingdahl K, Oberg B, Hansson SO, Hagg O, Lundberg
   363
   U, et al. The significance of occupational exposure for low back pain-a
   364
   systematic review. Physiotherapy (United Kingdom). Conference: World
- Confederation for Physical Therapy Congress 2015. Singapore Singapore.
- Conference Publication. 101(SUPPL. 1):eS1456-eS1457.
- Reason for exclusion: Ineligible population.
- 368 74. Bartley C, Webb JA, Bayly J. Multidisciplinary approaches to moving and handling for formal and informal carers in community palliative care. Int 370 J Palliat Nurs. 2015;21(1):17-23.
- Reason for exclusion: Ineligible population.
- 75. Kay K, Evans A, Glass N. Moments of speaking and silencing: Nursesshare their experiences of manual handling in healthcare. Collegian.
- 374 2015;22(1):61-70.
- 375 Reason for exclusion: Ineligible concept.
- 376 76. Abedini R, Choobineh AR, Hasanzadeh J. Patient manual handling risk 377 assessment among hospital nurses. Work. 2015;50(4):669-675.
- 378 Reason for exclusion: Ineligible context.
- 379 77. Aslam I, Davis SA, Feldman SR, Martin WE. A Review of Patient Lifting
   380 Interventions to Reduce Health Care Worker Injuries. Workplace health &
   381 safety. 2015;63(6):267-276.
- Reason for exclusion: Ineligible concept.
- 383 78. Bhimani R. Understanding Work-related Musculoskeletal Injuries in
- Rehabilitation from a Nursing Perspective. Rehabilitation nursing: the
- official journal of the Association of Rehabil Nurs. 2016;41(2):91-100.
- Reason for exclusion: Ineligible concept.

- 79. Boniface G, Ghosh S, Robinson L. District nurses' experiences of musculoskeletal wellbeing: a qualitative study. Br J Community Nurs.
- 389 2016;21(7):350-355.
- 390 Reason for exclusion: Ineligible concept.
- 391 80. Choi J, Cramer E. Reports From RNs on Safe Patient Handling and
- 392 Mobility Programs in Acute Care Hospital Units. J Nurs Adm.
- 393 2016;46(11):566-573.
- Reason for exclusion: Ineligible concept.
- 395 81. Freiberg A, Girbig M, Euler U, Scharfe J, Nienhaus A, Freitag S, et al.
- Influence of the Kinaesthetics care conception during patient handling on
- 397 the development of musculoskeletal complaints and diseases A scoping
- review. Journal of Occupational Medicine and Toxicology. 2016;11(1):1-
- 399 14.
- 400 Reason for exclusion: Ineligible concept.
- 401 82. Kuipers DA, Wartena BO, Dijkstra BH, Terlouw G, van t Veer, J. T. B.,
- van Dijk HW, et al. iLift: A health behavior change support system for
- 403 lifting and transfer techniques to prevent lower-back injuries in
- 404 healthcare. Int J Med Inf. 2016;96:11-23.
- 405 Reason for exclusion: Ineligible concept.
- 406 83. Coales P. Physiotherapists' perceptions of contributory and risk reduction
- factors for work-related spinal disorders in the profession. Physiotherapy
- 408 (United Kingdom). Conference: Physiotherapy UK Conference 2017.
- 409 2017;103(Supplement 1):e90.
- 410 Reason for exclusion: Ineligible concept.
- 411 84. Nagavarapu S, Lavender SA, Marras WS. Spine loading during the
- application and removal of lifting slings: the effects of patient weight,
- bed height and work method. Ergonomics. 2017;60(5):636-648.
- 414 Reason for exclusion: Ineligible population.
- 415 85. WhiteHeisel R, Canfield JP, YoungHughes S. Examining the Factor
- Structure and Reliability of the Safe Patient Handling Perception Scale:
- 417 An Initial Validation Study. Rehabil nurs. 2017;42(3):164-171.
- 418 Reason for exclusion: Ineligible concept.
- 419 86. Lee SJ, Lee JH. Safe patient handling behaviors and lift use among
- hospital nurses: A cross-sectional study. Int J Nurs Stud. 2017;74:53-

- 421 60.
- 422 Reason for exclusion: Ineligible concept.
- 423 87. Gold JE, Punnett L, Gore RJ. Predictors of low back pain in nursing home
- workers after implementation of a safe resident handling programme.
- 425 Occup Environ Med. 2017;74(6):389-395.
- 426 Reason for exclusion: Ineligible concept.
- 427 88. Contrada E. 1.5 CE Test Hours: Original Research: Patient Handling and
- 428 Mobility Course Content: A National Survey of Nursing Programs. Am J
- 429 Nurs. 2018;118(11):32-45.
- 430 Reason for exclusion: Ineligible concept.
- 431 89. Gold JE, Kurowski A, Gore RJ, Punnett L. Knee pain in nursing home
- workers after implementation of a safe resident handling program. Am J
- 433 Ind Med. 2018;61(10):849-860.
- 434 Reason for exclusion: Ineligible concept.
- 435 90. PowellCope G, Rugs D, Ialynytchev A, Devine D, McCoskey K, Zhang Y,
- et al. CE: Original Research: Patient Handling and Mobility Course
- 437 Content: A National Survey of Nursing Programs. Am J Nurs.
- 438 2018;118(11):22-31.
- 439 Reason for exclusion: Ineligible population.
- 91. Richardson A, McNoe B, Derrett S, Harcombe H. Interventions to prevent
- and reduce the impact of musculoskeletal injuries among nurses: A
- systematic review. Int J Nurs Stud. 2018;82:58-67.
- 443 Reason for exclusion: Ineligible concept.
- 444 92. Hurtado DA, Dumet LM, Greenspan SA, Rodriguez YI. Social Network
- Analysis of peer-specific safety support and ergonomic behaviors: An
- application to safe patient handling. Appl Ergon. 2018;68:132-137.
- 447 Reason for exclusion: Ineligible concept.
- 448 93. Havey R, Aebersold C, Terrasi J. "Add it to the Parking Lot": Safe Patient
- Handling and Mobility in the Emergency Department. Crit Care Nurs Q.
- 450 2018;41(3):302-311.
- 451 Reason for exclusion: Ineligible concept.
- 94. Karstad K, Jorgensen AFB, Greiner BA, Burdorf A, Sogaard K, Rugulies R,
- et al. Danish Observational Study of Eldercare work and musculoskeletal
- disorderS (DOSES): A prospective study at 20 nursing homes in

- 455 Denmark. BMJ Open. 2018;8(2):e019670. 456 Reason for exclusion: Ineligible concept. 457 95. Totzkay DL. Multifactorial Strategies for Sustaining Safe Patient Handling 458 and Mobility. Crit Care Nurs Q. 2018;41(3):340-344. 459 Reason for exclusion: Ineligible concept. 96. Worobey LA, Zigler CK, Huzinec R, Rigot SK, Sung J, Rice LA. Reliability 460 461 and validity of the revised transfer assessment instrument. Topics in Spinal Cord Injury Rehabilitation. 2018;24(3):217-226. 462 463 Reason for exclusion: Ineligible concept. 464 97. Nolan D, O'Sullivan K, Stephenson J, O'Sullivan P, Lucock M. What do 465 physiotherapists and manual handling advisors consider the safest lifting 466 posture, and do back beliefs influence their choice?. Musculoskeletal 467 Science and Practice. 2018;33:35-40. 468 Reason for exclusion: Ineligible setting. 469 98. Hegewald J, Berge W, Heinrich P, Staudte R, Freiberg A, Scharfe J, et al. 470 Do technical aids for patient handling prevent musculoskeletal complaints 471 in health care workers?-A systematic review of intervention studies. 472 International Journal of Environmental Research and Public Health 473 2018;15(3):1-19. Reason for exclusion: Ineligible concept. 474 475 99. Dias A, Bernardes JM. Low back pain and manual patient handling among 476 healthcare workers: A cross-sectional study. Occupational and 477 Environmental Medicine. Conference: 27th Epidemiology in Occupational 478 Health Conference, EPICOH 2019. New Zealand. 2019;76(Supplement 479 1):97-A98. 480 Reason for exclusion: Ineligible concept. 481 100. Johnstone J. Manual handling: the challenges of different care 482 environments. Br J Nurs. 2020;29(6):358-363. 483 Reason for exclusion: Ineligible concept.
- 101.Naidoo D, Govender P, Naidoo SN, Ngubane N, Nkosi Z, Mulla A.
   Occupational Risks in Occupational Therapy Service Learning: A Single-Site "Fear Factor" Study in South Africa. Occup Ther Int. 2020.
   2020:e4746813.
   Reason for exclusion: Ineligible rank HDI.

489 102.Rugs D, PowellCope G, Campo M, Darragh A, Harwood K, Kuhn J, et al. 490 The use of safe patient handling and mobility equipment in rehabilitation. 491 Work. 2020;66(1):31-40. 492 Reason for exclusion: Ineligible concept. 493 103. Gilthorpe J, Fairbairn S, Taylor S, Hutchinson A, Ford C, Tweedlie J. 494 Interprofessional education for practice: moving and handling for people 495 with complex needs in contemporary healthcare. Physiotherapy (United 496 Kingdom). Conference: Physiotherapy UK Conference 2019. 497 2019;107(Supplement 1):e170-e171. 498 Reason for exclusion: Ineligible population. 499 104. Garzillo EM, Monaco MGL, Corvino AR, D'ancicco F, Feola D, Ventura DD, 500 et al. Healthcare workers and manual patient handling: A pilot study for 501 interdisciplinary training. Int J Env Res Pub He. 2020;17(14):1-14. 502 Reason for exclusion: Ineligible concept. 503 105.Lecca LI, Fabbri D, Portoghese I, Pilia I, Meloni F, Marcias G, et al. 504 Manual handling of patients: role of kinesiophobia and catastrophizing in 505 health workers with chronic low back pain. Eur J Phys Rehabil Med. 506 2020;56(3):307-312. 507 Reason for exclusion: Ineligible concept. 508 106. Tang K, Diaz J, Lui O, Proulx L, Galle E, Packham T. Do active assist 509 transfer devices improve transfer safety for patients and caregivers in 510 hospital and community settings? A scoping review. Disability and 511 rehabilitation. Assist technol. 2020;15(6):614-624. 512 Reason for exclusion: Ineligible concept. 107. Campo M, Weiser S, Koenig KL, Nordin M. Work-Related Musculoskeletal 513 514 Disorders in Physical Therapists: A Prospective Cohort Study With 1-Year 515 Follow-up. Phys Ther. 2008;88(5):608-619. 516 Reason for exclusion: Ineligible concept. 517 108.Cromie JE, Robertson VJ, Best MO. Work-Related Musculoskeletal Disorders and the Culture of Physical Therapy. Phys Ther. 518 519 2002;82(5):459-472. 520 Reason for exclusion: Ineligible concept. 521 109. Vieira ER, Kumar S. Working postures: a literature review. J Occup 522 Rehabil. 2004;14(2):143-159. 523 Reason for exclusion: Ineligible population.

524 110. Mani S, Sharma S, Omar B, Ahmad K, Muniandy Y, Singh DKA. 525 Quantitative measurements of forward head posture in a clinical settings: 526 a technical feasibility study. Eur J Physiother. 2017;19(3):119-123. 527 Reason for exclusion: Ineligible population. 528 111. Thomas K. Pondering Posture. Co-Kinetic Journal. 2020(84):24-29. 529 Reason for exclusion: Ineligible concept. 530 112. Tam GYT, Yeung SS. Perceived effort and low back pain in non-531 emergency ambulance workers: Implications for rehabilitation. J Occup 532 Rehabil. 2006;16(2):231-240. 533 Reason for exclusion: Ineligible concept. 534 113. Dewey AJ, Olkowski BF, Stolfi AM. On "Safe patient handling perceptions 535 and practices . . . ". Phys Ther. 2014;94(7):1054-1055. 536 Reason for exclusion: Ineligible concept. 537 114. Cooklin A, Joss N, Husser E, Oldenburg B. Integrated Approaches to 538 Occupational Health and Safety: A Systematic Review. Am J Health 539 Promot. 2017;31(5):401-412. 540 Reason for exclusion: Ineligible concept. 541 115. Campo M, Darragh AR. Impact of Work-Related Pain on Physical 542 Therapists and Occupational Therapists. Phys Ther. 2010;90(6):905-920. 543 Reason for exclusion: Ineligible concept. 544 116. Kurowski A, Pransky G, Punnett L. Impact of a Safe Resident Handling 545 Program in Nursing Homes on Return-to-Work and Re-injury Outcomes 546 Following Work Injury. J Occup Rehabil. 2019;29(2):286-294. 547 Reason for exclusion: Ineligible concept. 117. Roussel NA, Kos D, Demeure I, Heyrman A, De Clerck M, Zinzen E, et al. 548 549 Effect of a multidisciplinary program for the prevention of low back pain 550 in hospital employees: A randomized controlled trial. J Back 551 Musculoskelet Rehabil. 2015;28(3):539-549. 552 Reason for exclusion: Ineligible concept. 118. Mustard C, Kalcevich C, Steenstra I, Smith P, Amick B. Disability 553 554 Management Outcomes in the Ontario Long-Term Care Sector. J Occup 555 Rehabil. 2010;20(4):481-488. 556 Reason for exclusion: Ineligible concept. 557 119. Ferguson SA, Marras WS, Burr DL. Differences Among Outcome

Measures in Occupational Low Back Pain. J Occup Rehabil.

559	2005;15(3):329-341.
560	Reason for exclusion: Ineligible population.
561	120. Frings-Dresen M, Sluiter JK. Development of a Job-Specific FCE Protocol:
562	The Work Demands of Hospital Nurses as an Example. J Occup Rehabil.
563	2003;13(4):233-248.
564	Reason for exclusion: Protocol only.
565	121.Reesink D, Jorritsma W, Reneman M. Basis for a Functional Capacity
566	Evaluation Methodology for Patients with Work-related Neck Disorders. J
567	Occup Rehabil. 2007;17(3):436-449.
568	Reason for exclusion: Ineligible population.
569	122. Darragh AR, Campo M, King P. Work-related activities associated with
570	injury in occupational and physical therapists. Work. 2012;42(3):373-
571	384.
572	Reason for exclusion: Ineligible concept.
573	123.Lapane KL, Dubé C,E., Jesdale BM. Worker Injuries in Nursing Homes: is
574	Safe Patient Handling Legislation the Solution? J Nurs Home Res Sci.
575	2016;2:110-117.
576	Reason for exclusion: Ineligible concept.
577	124.Clemes SA, Haslam CO, Haslam RA. What constitutes effective manual
578	handling training? A systematic review. Occup Med. 2010;60(2):101-
579	107.
580	Reason for exclusion: Ineligible concept.
581	125.Alberto R, Draicchio F, Varrecchia T, Silvetti A, Iavicoli S. Wearable
582	Monitoring Devices for Biomechanical Risk Assessment at Work: Current
583	Status and Future Challenges-A Systematic Review. Int J Environ Res
584	Public Health. 2018;15(9).
585	Reason for exclusion: Ineligible setting.
586	126.Gusenius TM, Decker MM, Weidemann AG. Using shared governance to
587	achieve a culture change in safe patient handling. Int J Orthop Trauma
588	Nurs. 2018;31:35-39.
589	Reason for exclusion: Ineligible concept.
590	127.Carta A, Parmigiani F, Roversi A, Rossato R, Milini C, Parrinello G, et al.
591	Training in safer and healthier patient handling techniques. Br J Nurs.
592	2010;19(9):576-582.
593	Reason for exclusion: Ineligible concept.

594	128.Lin F, Wang A, Cavuoto L, Xu W. Toward Unobtrusive Patient Handling
595	Activity Recognition for Injury Reduction Among At-Risk Caregivers. IEEE
596	J Biomed Health Inform. 2017;21(3):682-695.
597	Reason for exclusion: Ineligible concept.
598	129. Jones S. The nurse who could lift. Nursing. 2017;47(5):42-44.
599	Reason for exclusion: Ineligible concept.
600	130.Koppelaar E, Knibbe HJJ, Miedema HS, Burdorf A. The influence of
601	ergonomic devices on mechanical load during patient handling activities
602	in nursing homes. Ann Occup Hyg. 2012;56(6):708-718.
603	Reason for exclusion: Ineligible concept.
604	131. Eitzen D, Byard RW. The handling of bariatric bodies. J Forensic Leg Med.
605	2013;20(1):57-59.
606	Reason for exclusion: Ineligible concept.
607	132.Peterson EL, McGlothlin JD, Blue CL. The development of an ergonomics
608	training program to identify, evaluate, and control musculoskeletal
609	disorders among nursing assistants at a state-run veterans' home. J
610	Occup Environ Hyg. 2004;1(1):10-16.
611	Reason for exclusion: Ineligible concept.
612	133. Trossman S. Taking safe patient handling to the home front. Am Nurse.
613	2009;41(6):14.
614	Reason for exclusion: Ineligible concept.
615	134.Gilbert JH, Vermillion B, Chase LK. Stop the pain: Reinforcing a
616	successful ergonomics program. Nurs Manage. 2012;43(7):18-20.
617	Reason for exclusion: Ineligible concept.
618	135.Cimolin V, Cau N, Tacchini E, Galli M, Rigoldi C, Rinolfi M, et al. Spinal
619	load in nurses during emergency lifting of obese patients: preliminary
620	results. Med Lav. 2016;107(5):356-363.
621	Reason for exclusion: Ineligible concept.
622	136.Douglas B, Fitzpatrick D, Golub-Victor A, Lowe SM. Should my patient
623	use a mechanical lift? Part 2: algorithm and case application. Home
624	Healthc Nurse. 2014;32(3):172-180.
625	Reason for exclusion: Ineligible concept.
626	137.Bacharach DW, Miller K, von Duvillard S,P. Saving your back: How do
627	horizontal patient transfer devices stack up? Nursing. 2016;46(1):59-64.
628	Reason for exclusion: Ineligible concept.

629 138. Dunning E. Safer patient handling in your grasp. Nurs Manage. 630 2009;40(2):33-37. 631 Reason for exclusion: Ineligible concept. 632 139. Wilson CB. Safer handling practice: influence of staff education on older 633 people. Br J Nurs. 2002;11(20):1332-1339. 634 Reason for exclusion: Ineligible concept. 635 140. Bunke JF. Safe strategies for lifting large patients. Nursing. 2008;38:4-6. 636 Reason for exclusion: Ineligible concept. 637 141. Watters C. Safe patient handling: are we doing it yet? Orthop Nurs. 638 2008;27(1):38-41. 639 Reason for exclusion: Ineligible concept. 640 142. Kennedy B, Kopp T. Safe patient handling protects employees too. Nursing. 2015;45(8):65-67. 641 642 Reason for exclusion: Ineligible concept. 643 143. Dieker E, Steffes A, Tidwell S. Safe patient handling curriculum content. 644 Kans Nurse. 2007;82(3):8-9. 645 Reason for exclusion: Ineligible concept. 646 144. Durham CF. Safe patient handling and movement: time for a culture 647 change. Tar Heel Nurse. 2007;69(4):16-18. 648 Reason for exclusion: Ineligible concept. 649 145. Nelson A, Owen B, Lloyd JD, Fragala G, Matz MW, Amato M, et al. Safe 650 patient handling and movement. Am J Nurs. 2003;103(3):32-43. 651 Reason for exclusion: Ineligible concept. 652 146. Spruce L. Safe Patient Handling and Movement. AORN J. 653 2020;112(1):63-71. 654 Reason for exclusion: Ineligible concept. 147.Restrepo TE, Schmid FA, Gucer PW, Shuford HL, Shyong CJ, McDiarmid 655 656 MA. Safe lifting programs at long-term care facilities and their impact on 657 workers' compensation costs. J Occup Environ Med. 2013;55(1):27-35. 658 Reason for exclusion: Ineligible concept. 659 148. Relias Media. Safe lifting becomes standard practice. Hosp Case Manag. 660 2013;21(2):26-28.

Reason for exclusion: Ineligible concept.

662 149. Patrick G. Risks of moving and handling. Nurs Stand. 2016;30(52):64-663 65. 664 Reason for exclusion: Ineligible concept. 665 150.Ngan K, Drebit S, Siow S, Yu S, Keen D, Alamgir H. Risks and causes of 666 musculoskeletal injuries among health care workers. Occup Med. 667 2010;60(5):389-394. 668 Reason for exclusion: Ineligible concept. 669 151.Lee S, Faucett J, Gillen M, Krause N, Landry L. Risk perception of 670 musculoskeletal injury among critical care nurses. Nurs Res. 671 2013;62(1):36-44. 672 Reason for exclusion: Ineligible concept. 673 152. Holtermann A, Clausen T, Aust B, Mortensen OS, Andersen LL. Risk for 674 low back pain from different frequencies, load mass and trunk postures 675 of lifting and carrying among female healthcare workers. Int Arch Occup 676 Environ Health. 2013;86(4):463-470. 677 Reason for exclusion: Ineligible concept. 153. Caspi CE, Dennerlein JT, Kenwood C, Stoddard AM, Hopcia K, Hashimoto 678 679 D, et al. Results of a pilot intervention to improve health and safety for 680 health care workers. J Occup Environ Med. 2013;55(12):1449-1455. 681 Reason for exclusion: Ineligible concept. 682 154. Makhoul PJ, Sinden KE, MacPhee RS, Fischer SL. Relative Contribution of 683 Lower Body Work as a Biomechanical Determinant of Spine Sparing 684 Technique During Common Paramedic Lifting Tasks. J Appl Biomech. 685 2017;33(2):137-143. 686 Reason for exclusion: Ineligible concept. 687 155. Nam S, Song M, Lee S. Relationships of Musculoskeletal Symptoms, 688 Sociodemographics, and Body Mass Index With Leisure-Time Physical 689 Activity Among Nurses. Workplace Health Saf. 2018;66(12):577-587. 690 Reason for exclusion: Ineligible concept. 691 156.Kitagawa K, Uezono T, Nagasaki T, Nakano S, Wada C. Relationship 692 between surface electromyography of the spinae erector muscles and 693 subjectively adjusted step length in the supporting standing-up motion. J 694 Phys Ther Sci. 2019;31(11):869-872. 695 Reason for exclusion: Ineligible population.

696 157. Alperovitch-Najenson D, Sheffer D, Treger I, Finkels T, Kalichman L. 697 Rehabilitation versus Nursing Home Nurses' Low Back and Neck-Shoulder 698 Complaints. Rehabil Nurs. 2015;40(5):286-293. 699 Reason for exclusion: Ineligible concept. 700 158. Davies N. Reducing your risk of back injury at work. Nurs Stand. 701 2015;29(25):73. 702 Reason for exclusion: Ineligible concept. 703 159. Resnick ML, Sanchez R. Reducing patient handling injuries through 704 contextual training. J Emerg Nurs. 2009;35(6):504-508. 705 Reason for exclusion: Ineligible concept. 706 160. Fragala G. Reducing Occupational Risk to Ambulatory Caregivers. 707 Workplace Health Saf. 2016;64(9):414-419. 708 Reason for exclusion: Ineligible concept. 709 161. Owlia M, Kamachi M, Dutta T. Reducing lumbar spine flexion using real-710 time biofeedback during patient handling tasks. Work. 2020;66(1):41-711 51. Reason for exclusion: Ineligible population. 712 713 162. Ann Adamczyk M. Reducing Intensive Care Unit Staff Musculoskeletal 714 Injuries With Implementation of a Safe Patient Handling and Mobility 715 Program. Crit Care Nurs Q. 2018;41(3):264-271. 716 Reason for exclusion: Unable to access paper. 717 163. Lipley N. Progress slow on safe handling policies. Emerg Nurse. 718 2002;10(1):2. 719 Reason for exclusion: Ineligible concept. 720 164.Rugs D, Toyinbo P, Patel N, Powell-Cope G, Hahm B, Elnitsky C, et al. 721 Processes and outcomes of the veterans health administration safe 722 patient handling program: study protocol. JMIR Res Protoc. 723 2013;2(2):e49. 724 Reason for exclusion: Ineligible concept. 725 165.de Castro A,B., Hagan P, Nelson A. Prioritizing safe patient handling: The 726 American Nurses Association's Handle With Care Campaign. J Nurs Adm.

727

728

2006;36(7-8):363-369.

Reason for exclusion: Ineligible concept.

729	166.Bhimani R. Prevention of Work-related Musculoskeletal Injuries in
730	Rehabilitation Nursing. Rehabil Nurs. 2016;41(6):326-335.
731	Reason for exclusion: Ineligible concept.
732	167.Spruce L. Preventing Staff Member Injury Through Safe Patient Handling
733	and Movement. AORN J. 2020;111(5):567-568.
734	Reason for exclusion: Ineligible concept.
735	168.Brown MR, Personeus K, Langan J. Preventing lower back pain among
736	caregivers. Arch Phys Med Rehabil. 2015;96(7):1365-1366.
737	Reason for exclusion: Ineligible concept.
738	169. Homsted L. Preventing back injuries: safe patient handling and
739	movement. Fla Nurse. 2004;52(3):3.
740	Reason for exclusion: Ineligible concept.
741	170.Bell J, Collins J, Galinsky TL, Waters TR. Preventing back injuries. Ala
742	Nurse. 2009;36(2):16.
743	Reason for exclusion: Ineligible concept.
744	171.Blocks M. Practical solutions for safe patient handling. Nursing.
745	2005;35(10):44-45.
746	Reason for exclusion: Ineligible concept.
747	172.Chan F, Hsu C, Lin H, Wang J, Su S, Huang C, et al. Physicians as well as
748	nonphysician health care professionals in Taiwan have higher risk for
749	lumbar herniated intervertebral disc than general population. Medicine.
750	2018;97(1):e9561.
751	Reason for exclusion: Ineligible concept.
752	173.Alperovitch-Najenson D, Treger I, Kalichman L. Physical therapists
753	versus nurses in a rehabilitation hospital: comparing prevalence of work-
754	related musculoskeletal complaints and working conditions. Arch Environ
755	Occup Health. 2014;69(1):33-39.
756	Reason for exclusion: Ineligible concept.
757	174.Andersen LL, Vinstrup J, Villadsen E, Jay K, Jakobsen MD. Physical and
758	Psychosocial Work Environmental Risk Factors for Back Injury among
759	Healthcare Workers: Prospective Cohort Study. Int J Environ Res Public
760	Health. 2019;16(22).
761	Reason for exclusion: Ineligible concept.
762	175. Vinstrup J, Madeleine P, Jakobsen MD, Jay K, Andersen LL. Patient
763	Transfers and Risk of Back Injury: Protocol for a Prospective Cohort

764 Study With Technical Measurements of Exposure. JMIR Res Protoc. 765 2017;6(11):e212. 766 Reason for exclusion: Protocol only. 767 176.Blair S, Bratton M. Patient handling and time outs: The way to sustain 768 safety. Nurs Manage. 2015;46(6):46-50. 769 Reason for exclusion: Ineligible concept. 770 177.Lee J, Kim S, Jung H, Koo J, Woo K, Kim MT. Participatory action 771 oriented training for hospital nurses (PAOTHN) program to prevent 772 musculoskeletal disorders. J Occup Health. 2009;51(4):370-376. 773 Reason for exclusion: Ineligible concept. 774 178. Tideiksaar R. Part I: Achieving safe resident handling. Director. 775 2008;16(2):27-29. 776 Reason for exclusion: Ineligible concept. 777 179.Rodríguez-Acosta RL, Richardson DB, Lipscomb HJ, Chen JC, Dement JM, 778 Myers DJ, et al. Occupational injuries among aides and nurses in acute 779 care. Am J Ind Med. 2009;52(12):953-964. 780 Reason for exclusion: Ineligible concept. 781 180. Zubair A. Nurses at risk of injuries. Aust Nurs Midwifery J. 782 2015;23(2):26-28. 783 Reason for exclusion: Ineligible concept. 784 181.NJSNA. NJSNA members participate in multi-state study to assess safe 785 patient handling. N J Nurse. 2002;32(5):20. 786 Reason for exclusion: Ineligible concept. 787 182.Cameron SJ, Armstrong-Stassen M, Kane D, Moro FBP. Musculoskeletal 788 problems experienced by older nurses in hospital settings. Nurs Forum. 789 2008;43(2):103-114. 790 Reason for exclusion: Ineligible concept. 791 183. Caponecchia C, Coman RL, Gopaldasani V, Mayland EC, Campbell L. 792 Musculoskeletal disorders in aged care workers: a systematic review of 793 contributing factors and interventions. Int J Nurs Stud. 794 2020;110:103715. 795 Reason for exclusion: Ineligible concept. 796 184. Gillespie T, Lane S. Moving the Bariatric Patient. Crit Care Nurs Q. 797 2018;41(3):297-301. 798 Reason for exclusion: Ineligible concept.

799	185. Warren G. Moving and handling: reducing risk through assessment. Nurs
800	Stand. 2016;30(40):49-58.
801	Reason for exclusion: Ineligible concept.
802	186. Wanless S, Page A. Moving and handling education in the community:
803	technological innovations to improve practice. Br J Community Nurs.
804	2009;14(12):530-532.
805	Reason for exclusion: Ineligible concept.
806	187. Theilmeier A, Jordan C, Luttmann A, Jäger M. Measurement of action
807	forces and posture to determine the lumbar load of healthcare workers
808	during care activities with patient transfers. Ann Occup Hyg.
809	2010;54(8):923-933.
810	Reason for exclusion: Ineligible concept.
811	188. Winkelman C, Chiang L. Manual turns in patients receiving mechanical
812	ventilation. Crit Care Nurse. 2010;30(4):36-44.
813	Reason for exclusion: Ineligible concept.
814	189. Cowley SP, Leggett S. Manual handling risks associated with the care,
815	treatment and transportation of bariatric (severely obese) clients in
816	Australia. Work. 2011;39(4):477-483.
817	Reason for exclusion: Ineligible concept.
818	190. Thompson S, Jevon P. Manual handling 1assisted stand. The manual
819	handling procedure to help a seated patient into a standing position.
820	Nurs Times. 2008;104(50-51):36-37.
821	Reason for exclusion: Ineligible concept.
822	191.Dennerlein JT, O'Day ET, Mulloy DF, Somerville J, Stoddard AM, Kenwood
823	C, et al. Lifting and exertion injuries decrease after implementation of an
824	integrated hospital-wide safe patient handling and mobilisation
825	programme. Occup Environ Med. 2017;74(5):336-343.
826	Reason for exclusion: Ineligible concept.
827	192. Stevenson A. Learning to handle patients safely. Nurs N Z.
828	2009;15(4):24-25.
829	Reason for exclusion: Ineligible concept.
830	193. Thomas DR, Thomas YLN. Interventions to reduce injuries when
831	transferring patients: a critical appraisal of reviews and a realist
832	synthesis. Int J Nurs Stud. 2014;51(10):1381-1394.
833	Reason for exclusion: Ineligible concept.

834	194. Hignett S. Intervention strategies to reduce musculoskeletal injuries
835	associated with handling patients: a systematic review. Occup Environ
836	Med. 2003;60(9):E6.
837	Reason for exclusion: Ineligible concept.
838	195.Wyatt S, Meacci K, Arnold M. Integrating Safe Patient Handling and Early
839	Mobility: Combining Quality Initiatives. J Nurs Care Qual.
840	2020;35(2):130-134.
841	Reason for exclusion: Ineligible concept.
842	196.Dickinson S, Taylor S, Anton P. Integrating a Standardized Mobility
843	Program and Safe Patient Handling. Crit Care Nurs Q. 2018;41(3):240-
844	252.
845	Reason for exclusion: Ineligible concept.
846	197.Przybysz L, Levin PF. Initial Results of an Evidence-Based Safe Patient
847	Handling and Mobility Program to Decrease Hospital Worker Injuries.
848	Workplace Health Saf. 2017;65(2):83-88.
849	Reason for exclusion: Ineligible concept.
850	198. Wanless S, Wanless SG. Improving training and education in patient
851	handling. Nurs Times. 2011;107(23):17-19.
852	Reason for exclusion: Ineligible population.
853	199. Fragala G, Fragala M. Improving the safety of patient turning and
854	repositioning tasks for caregivers. Workplace Health Saf.
855	2014;62(7):268-273.
856	Reason for exclusion: Ineligible concept.
857	200.Park RM, Bushnell PT, Bailer AJ, Collins JW, Stayner LT. Impact of
858	publicly sponsored interventions on musculoskeletal injury claims in
859	nursing homes. Am J Ind Med. 2009;52(9):683-697.
860	Reason for exclusion: Ineligible concept.
861	201. Varnam W. How to mobilise patients with dementia to a standing
862	position. Nurs Older People. 2011;23(8):31-36.
863	Reason for exclusion: Ineligible concept.
864	202. Taylor J. How to assist patients with sit-stand transfers. Nurs Stand.
865	2017;31(37):41-45.
866	Reason for exclusion: Ineligible concept.
867	203.Serranheira F, Sousa-Uva M, Sousa-Uva A. Hospital nurses tasks and

work-related musculoskeletal disorders symptoms: A detailed analysis.

869 Work. 2015;51(3):401-409. 870 Reason for exclusion: Ineligible concept. 204. Enos L. Hidden costs: the case for ergonomics and safe patient handling. 871 872 Oreg Nurse. 2010:5. 873 Reason for exclusion: Ineligible concept. 874 205. Worthington KA. Handling and moving patients--safely: protect patients 875 and lighten your load. Am J Nurs. 2002;102(6):112. 876 Reason for exclusion: Ineligible concept. 877 206. Allen LK, Durham CE. Handle with care: a new curriculum for safe patient 878 handling and movement at the University of North Carolina at Chapel 879 Hill. Tar Heel Nurse. 2006;68(2):21-23. 880 Reason for exclusion: Ineligible concept. 881 207. Carlson AH. Handle with care. Rehab Manag. 2008;21(9):33-35. 882 Reason for exclusion: Ineligible concept. 883 208.AORN. Guideline Quick View: Safe Patient Handling and Movement. 884 AORN J. 2018;108(2):223-227. 885 Reason for exclusion: Ineligible concept. 886 209.Link T. Guideline Implementation: Safe Patient Handling and Movement: 887 1.8 www.aornjournal.org/content/cme. AORN J. 2018;108(6):663-674. 888 Reason for exclusion: Ineligible concept. 889 210. Hauk L. Guideline for safe patient handling and movement. AORN J. 890 2018;107(6):10-12. 891 Reason for exclusion: Ineligible concept. 892 211.AORN. Guideline at a Glance: Positioning. AORN J. 2017;106(5):460-893 465. 894 Reason for exclusion: Ineligible concept. 895 212. Trossman S. Getting a lift. ANA, CMA and RN efforts continue to build 896 momentum for safe patient handling movement. Am Nurse. 897 2007;39(4):1. 898 Reason for exclusion: Ineligible concept. 899 213. Thiede M, Liebers F, Seidler A, Gravemeyer S, Latza U. Gender specific 900 analysis of occupational diseases of the low back caused by carrying, 901 lifting or extreme trunk flexion--use of a prevention index to identify 902 occupations with high prevention needs. Am J Ind Med. 2014;57(2):233-

903	244.
904	Reason for exclusion: Ineligible population.
905	214.Kayser SA, Wiggermann NE, Kumpar D. Factors associated with safe
906	patient handling practice in acute care and its relationship with patient
907	mobilization: A cross-sectional study. Int J Nurs Stud. 2020;104:103508.
908	Reason for exclusion: Ineligible concept.
909	215.Jootun D, MacInnes A. Examining how well students use correct handling
910	procedures. Nurs Times. 2005;101(4):38-40.
911	Reason for exclusion: Ineligible population.
912	216. Hignett S, Crumpton E, Ruszala S, Alexander P, Fray M, Fletcher B.
913	Evidence-based patient handling: systematic review. Nurs Stand.
914	2003;17(33):33-36.
915	Reason for exclusion: Ineligible concept.
916	217.Johnsson C, Carlsson R, Lagerström M. Evaluation of training in patient
917	handling and moving skills among hospital and home care personnel.
918	Ergonomics. 2002;45(12):850-865.
919	Reason for exclusion: Ineligible concept.
920	218. Hallmark B, Mechan P, Shores L. Ergonomics: safe patient handling and
921	mobility. Nurs Clin North Am. 2015;50(1):153-166.
922	Reason for exclusion: Ineligible concept.
923	219.Brusin JH. Ergonomics in radiology. Radiol Technol. 2011;83(2):141-157.
924	Reason for exclusion: Ineligible concept.
925	220. Ergonomic principles of safe patient handling. Healthc Hazard Manage
926	Monit. 2003;16(5):1-7.
927	Reason for exclusion: Unable to access paper.
928	221.Dawson JM, Harrington S. Embracing safe patient handling. Nurs
929	Manage. 2012;43(10):15-17.
930	Reason for exclusion: Ineligible concept.
931	222.Martin PJ, Harvey JT, Culvenor JF, Payne WR. Effect of a nurse back
932	injury prevention intervention on the rate of injury compensation claims.
933	J Safety Res. 2009;40(1):13-19.
934	Reason for exclusion: Ineligible concept.
935	223. Price C, Sanderson LV, Talarek DP. Don't pay the price: utilize safe
936	patient handling. Nursing. 2013;43(12):13-15.
937	Reason for exclusion: Ineligible concept.

938	224. Holtermann A, Clausen T, Jørgensen MB, Aust B, Mortensen OS, Burdorf
939	A, et al. Does rare use of assistive devices during patient handling
940	increase the risk of low back pain? A prospective cohort study among
941	female healthcare workers. Int Arch Occup Environ Health.
942	2015;88(3):335-342.
943	Reason for exclusion: Ineligible concept.
944	225.D'Arcy L,P., Sasai Y, Stearns SC. Do assistive devices, training, and
945	workload affect injury incidence? Prevention efforts by nursing homes
946	and back injuries among nursing assistants. J Adv Nurs.
947	2012;68(4):836-845.
948	Reason for exclusion: Ineligible concept.
949	226.Edlich RF, Hudson MA, Buschbacher RM, Winters KL, Britt LD, Cox MJ, et
950	al. Devastating injuries in healthcare workers: description of the crisis
951	and legislative solution to the epidemic of back injury from patient lifting.
952	J Long Term Eff Med Implants. 2005;15(2):225-241.
953	Reason for exclusion: Ineligible concept.
954	227.McKinney M. Curbing injuries from moving patients. Mod Healthc.
955	2015;45(13):28.
956	Reason for exclusion: Ineligible concept.
957	228. Stevens L, Rees S, Lamb KV, Dalsing D. Creating a culture of safety for
958	safe patient handling. Orthop Nurs. 2013;32(3):155-164.
959	Reason for exclusion: Ineligible concept.
960	229. Stenger K, Montgomery LA, Briesemeister E. Creating a culture of
961	change through implementation of a safe patient handling program. Crit
962	Care Nurs Clin North Am. 2007;19(2):213-222.
963	Reason for exclusion: Ineligible concept.
964	230.Kotejoshyer R, Punnett L, Dybel G, Buchholz B. Claim Costs,
965	Musculoskeletal Health, and Work Exposure in Physical Therapists,
966	Occupational Therapists, Physical Therapist Assistants, and Occupational
967	Therapist Assistants: A Comparison Among Long-Term Care Jobs. Phys
968	Ther. 2019;99(2):183-193.
969	Reason for exclusion: Ineligible concept.
970	231.Black JM, Salsbury S, Vollman KM. Changing the Perceptions of a Culture
971	of Safety for the Patient and the Caregiver: Integrating Improvement
972	Initiatives to Create Sustainable Change. Crit Care Nurs Q.

9/3	2018;41(3):226-239.
974	Reason for exclusion: Ineligible concept.
975	232.Rose MA, Drake DJ, Baker G, Watkins FR, Waters W, Pokorny M. Caring
976	for morbidly obese patients: safety considerations for nurse
977	administrators. Nurs Manage. 2008;39(11):47-50.
978	Reason for exclusion: Ineligible concept.
979	233.Alberta RN. Care for your back and it will love you back. Alta RN.
980	2013;68(4):26-27.
981	Reason for exclusion: Ineligible concept.
982	234.Knibbe JJ, Knibbe NE, Waaijer E. Assessments of patients with a 5-
983	category or a 3-category practical classification system: validity and
984	practicality. Work. 2012;41 Suppl 1:5655-5656.
985	Reason for exclusion: Ineligible concept.
986	235.Burdorf A, Koppelaar E, Evanoff B. Assessment of the impact of lifting
987	device use on low back pain and musculoskeletal injury claims among
988	nurses. Occup Environ Med. 2013;70(7):491-497.
989	Reason for exclusion: Ineligible concept.
990	236.Hwang J, Ari H, Matoo M, Chen J, Kim JH. Air-assisted devices reduce
991	biomechanical loading in the low back and upper extremities during
992	patient turning tasks. Appl Ergon. 2020;87:103121.
993	Reason for exclusion: Ineligible concept.
994	237. Nathenson P. Adapting OSHA ergonomic guidelines to the rehabilitation
995	setting. Rehabil Nurs. 2004;29(4):127-130.
996	Reason for exclusion: Ineligible concept.
997	238. Stutzman S. A rehabilitation hospital approach to safe patient
998	movement. Nebr Nurse. 2010;43(3):12.
999	Reason for exclusion: Ineligible concept.
1000	239. Johnson CL, Demass Martin S,L., Markle-Elder S. A national solution to
1001	patient handling injuries. Am J Nurs. 2007;107(9):73-75.
1002	Reason for exclusion: Ineligible concept.
1003	240.Lees L. A guide for HCAs on safe patient transfers. Nurs Times.
1004	2013;109(26):20-22.
1005	Reason for exclusion: Ineligible concept.
1006	Reason for exclusion: Ineligible concept.

1007	241.Dutta T, Holliday PJ, Gorski SM, Baharvandy MS, Fernie GR. A
1008	biomechanical assessment of floor and overhead lifts using one or two
1009	caregivers for patient transfers. Appl Ergon. 2012;43(3):521-531.
1010	Reason for exclusion: Ineligible concept.
1011	242.Choi SD, Brings K. Work-related musculoskeletal risks associated with
1012	nurses and nursing assistants handling overweight and obese patients: A
1013	literature review. Work. 2016;53(2):439-448.
1014	Reason for exclusion: Ineligible concept.
1015	243.Bolin M. Workplace safety in the surgical suite: OSHA's effect. AORN J.
1016	2012;96(1):1.
1017	Reason for exclusion: Ineligible concept.
1018	244.Spratt D, Cowles Jr C,E., Berguer R, Dennis V, Waters TR, Rodriguez M,
1019	et al. Workplace safety equals patient safety. AORN J. 2012;96(3):235-
1020	244.
1021	Reason for exclusion: Ineligible concept.
1022	245. Peled K. Workplace safety assessment and injury prevention in hospital
1023	settings. Work. 2005;25(3):273-277.
1024	Reason for exclusion: Ineligible concept.
1025	246. Converso A, Murphy C. Winning the battle against back injuries. RN.
1026	2004;67(2):52-82.
1027	Reason for exclusion: Ineligible concept.
1028	247. Stetler CB, Burns M, Sander-Buscemi K, Morsi D, Grunwald E. Use of
1029	evidence for prevention of work-related musculoskeletal injuries. Orthop
1030	Nurs. 2003 Jan;22(1):32-41.
1031	Reason for exclusion: Ineligible concept.
1032	248.Jabłońska R, Gralik M, Królikowska A, Haor B, Antczak A. The Problem of
1033	Back Pain Among Nurses of Neurology and Neurosurgery Wards. J Neurol
1034	Neurosurg Nurs. 2016;5(3):84-91.
1035	Reason for exclusion: Ineligible concept.
1036	249.Bakola H, Zyga S, Stergioulas A, Kipreos G, Sheikhzadeh A,
1037	Panoutsopoulos G. The "Perception of Musculoskeletal Risk Factors
1038	(PMRF)" for Operating Room Nurses (ORN): Translation and Cultural
1039	Adaption of the Questionnaire in the Greek Language. Int J Caring Sci.
1040	2017 May;10(2):834-841.
1041	Reason for exclusion: Ineligible concept.

1042	250. Jarus T, Ratzon NZ. The implementation of motor learning principles in
1043	designing prevention programs at work. Work. 2005;24(2):171-182.
1044	Reason for exclusion: Ineligible concept.
1045	251.Gallagher AM, Gilligan R, McGrath M. The effect of DVD training on the
1046	competence of occupational therapy students in manual handling: A pilot
1047	study. Int J Ther Rehabil. 2014;21(12):575-583.
1048	Reason for exclusion: Ineligible population.
1049	252.Ratzon NZ, Bar-Niv N, Froom P. The effect of a structured personalized
1050	ergonomic intervention program for hospital nurses with reported
1051	musculoskeletal pain: An assigned randomized control trial. Work.
1052	2016;54(2):367-377.
1053	Reason for exclusion: Ineligible concept.
1054	253. Chadwick M. Test your manual handling knowledge: answers. Nurs
1055	Residential Care. 2008;10(11):561-563.
1056	Reason for exclusion: Ineligible concept.
1057	254. Chadwick M. Test your manual handling knowledge. Nurs Residential
1058	Care. 2008;10(10):503-507.
1059	Reason for exclusion: Ineligible concept.
1060	255.Cabeças J,Martin Miquel. Taxonomy to characterize occupational hazards
1061	(risk factors) at the workplace level. Work. 2015;51(4):703-713.
1062	Reason for exclusion: Ineligible concept.
1063	256.Kennedy CA, III AB, Dennerlein JT, Brewer S, Catli S, Williams R, et al.
1064	Systematic review of the role of occupational health and safety
1065	interventions in the prevention of upper extremity musculoskeletal
1066	symptoms, signs, disorders, injuries, claims and lost time. J Occup
1067	Rehabil. 2010;20(2):127-162.
1068	Reason for exclusion: Ineligible concept.
1069	257.American Association for Safe Patient Handling and Movement. Spotlight
1070	On: American Association for Safe Patient Handling & Movement United
1071	States of America. Am J Safe Patient Handling Movement. 2014;4(2):62-
1072	63.
1073	Reason for exclusion: Ineligible concept.
1074	258. Walker L, Docherty T, Hougendobler D, Guanowsky C, Rosenthal M.
1075	Sharing the Lessons: the 10-Year Journey of a Safe Patient Movement

1076	Program. Int J Safe Patient Handl Mobil. 2017;7(1):20-28.
1077	Reason for exclusion: Ineligible concept.
1078	259. Monaghan HM. Selecting Effective Training and Education Strategies as
1079	Part of a Safe Patient Handling and Movement Program. Am J Safe
1080	Patient Handling Movement. 2011;1(4):17-21.
1081	Reason for exclusion: Ineligible concept.
1082	260.Scalzitti DA, Harwood KJ, Campo M. Searching the Literature for Safe
1083	Patient Handling and Mobility Evidence. Am J Safe Patient Handling
1084	Movement. 2016;6(1):10-15.
1085	Reason for exclusion: Ineligible concept.
1086	261.Nelson AL, Collins J, Knibbe H, Cookson K, Castro AB, Whipple KL. Safer
1087	patient handling. Nurs Manage. 2007;38(3):26-33.
1088	Reason for exclusion: Ineligible concept.
1089	262. Way H. Safe Patient Handling Initiative in Level I Trauma Center Results
1090	in Reduction of Hospital-Acquired Pressure Injury and Fewer Patient
1091	Handling Injuries. Am J Safe Patient Handling Movement.
1092	2016;6(4):160-165.
1093	Reason for exclusion: Ineligible concept.
1094	263. Murray E. Safe Patient Handling in the Morgue. Am J Safe Patient
1095	Handling Movement. 2012;2(2):48-53.
1096	Reason for exclusion: Ineligible concept.
1097	264. Capewell RA. Safe Patient Handling in Home Health Care: a Review of
1098	the Literature. Am J Safe Patient Handling Movement. 2011;1(2):25-34.
1099	Reason for exclusion: Ineligible concept.
1100	265.Matz MW. Safe Patient Handling Building Design Guidance: an Overview
1101	of the Patient Handling and Movement Assessment. Am J Safe Patient
1102	Handling Movement. 2011;1(2):13-18.
1103	Reason for exclusion: Ineligible concept.
1104	266.Enos L. Safe Patient Handling and Patient Safety: Identifying the Current
1105	Evidence Base and Gaps in Research. Am J Safe Patient Handling
1106	Movement. 2013;3(3):94-102.
1107	Reason for exclusion: Ineligible concept.
1108	267. Nelson K. Safe Patient Handling and Health Care Reform: an Opportunity
1109	to Link Patient and Worker Safety. Am J Safe Patient Handling

1110	Movement. 2015;5(1):9-12.
1111	Reason for exclusion: Ineligible concept.
1112	268.Monaghan HM, Murray E, Severson L, Kissing J. Safe Patient Handling &
1113	Movement: Long Term Care. Am J Safe Patient Handling Movement.
1114	2013;3:S1-S19.
1115	Reason for exclusion: Ineligible concept.
1116	269. Monaghan HM. Safe Patient Handling & Movement: Legislation,
1117	Accreditation, Quality, and Recognition. Am J Safe Patient Handling
1118	Movement. 2013;3(3):103-109.
1119	Reason for exclusion: Ineligible concept.
1120	270.Gallagher S, Hilton T, Monaghan HM, Muir M, Dye A. Safe Patient
1121	Handling & Movement: Bariatric Considerations. Am J Safe Patient
1122	Handling Movement. 2014:1-16.
1123	Reason for exclusion: Ineligible concept.
1124	271. Monaghan HM. Safe Patient Handling & Movement Around the World:
1125	United States of America-Past, Present, and Future. Am J Safe Patient
1126	Handling Movement. 2012;2(2):58-60.
1127	Reason for exclusion: Ineligible concept.
1128	272.Knibbe HJJ. Safe Patient Handling & Movement Around the World: the
1129	NetherlandsPast, Present, and Future. Am J Safe Patient Handling
1130	Movement. 2011;1(4):37-39.
1131	Reason for exclusion: Ineligible concept.
1132	273.Kuo W, Chang J. Safe Patient Handling & Movement Around the World:
1133	TaiwanPast, Present, and Future. Am J Safe Patient Handling
1134	Movement. 2013;3(1):38-39.
1135	Reason for exclusion: Ineligible concept.
1136	274.Lindahl O. Safe Patient Handling & Movement Around the World:
1137	ScandinaviaPast, Present, and Future. Am J Safe Patient Handling
1138	Movement. 2011;1(3):36-37.
1139	Reason for exclusion: Ineligible concept.
1140	275.Carton M. Safe Patient Handling & Movement Around the World: Saudi
1141	ArabiaPast, Present, and Future. Am J Safe Patient Handling
1142	Movement. 2013;3(2):74-75.
1143	Reason for exclusion: Ineligible concept

1144	276. Menoni O, Battevi N. Safe Patient Handling & Movement Around the
1145	World: ItalyPast, Present, and Future. Am J Safe Patient Handling
1146	Movement. 2012;2(3):100-101.
1147	Reason for exclusion: Ineligible concept.
1148	277.Geiger JS. Safe Patient Handling & Movement Around the World: Israel
1149	Past, Present, and Future. Am J Safe Patient Handling Movement.
1150	2012;2(1):33-35.
1151	Reason for exclusion: Ineligible concept.
1152	278. Monaghan HM. Safe Patient Handling & Movement Around the World:
1153	Editorial. Am J Safe Patient Handling Movement. 2012;2(4):138-139.
1154	Reason for exclusion: Ineligible concept.
1155	279.0'shea L. Safe Patient Handling & Movement Around the World:
1156	AustraliaPast, Present, and Future. Am J Safe Patient Handling
1157	Movement. 2011;1(2):35-37.
1158	Reason for exclusion: Ineligible concept.
1159	280.Blood R. Safe moving and handling of individuals. Nurs Residential Care.
1160	2005;7(10):439-441.
1161	Reason for exclusion: Ineligible concept.
1162	281.Wardell H. Reduction of injuries associated with patient handling. AAOHN
1163	J. 2007;55(10):407-412.
1164	Reason for exclusion: Ineligible concept.
1165	282.Punchihewa HKG, Gyi DE. Reducing work-related Musculoskeletal
1166	Disorders (MSDs) through design: Views of ergonomics and design
1167	practitioners. Work. 2016;53(1):127-142.
1168	Reason for exclusion: Ineligible population.
1169	283.Sampath SL, Wilson K, Davis K, Kotowski S. Reality of Safe Patient
1170	Handling Policies and Programs in Hospitals Across the United States. Int
1171	J Safe Patient Handl Mobil. 2019;9(2):69-76.
1172	Reason for exclusion: Ineligible concept.
1173	284. Jaklic A, Jagodic V. Quantitative assessment of nurses' work in intensive
1174	care units: comparison of the TISS-28 scoring system with the
1175	classification system used in Slovenia. Connect World Crit Care Nurs.
1176	2005;4(3):122.
1177	Reason for exclusion: Ineligible concept.

1178	285.Zontek TL, Isernhagen JC, Ogle BR. Psychosocial factors contributing to
1179	occupational injuries among direct care workers. AAOHN J.
1180	2009;57(8):338-347.
1181	Reason for exclusion: Ineligible concept.
1182	286.Nelson A, Lloyd JD, Menzel N, Gross C. Preventing nursing back injuries:
1183	redesigning patient handling tasks. AAOHN J. 2003;51(3):126-134.
1184	Reason for exclusion: Ineligible concept.
1185	287. White KM. Policy spotlight: patient care ergonomics: are we moving in
1186	the right direction? Nurs Manage. 2007;38(4):26-30.
1187	Reason for exclusion: Ineligible concept.
1188	288.Franklin J, Yeung Y. Play it safe: use proper ergonomics to protect
1189	patients and staff. Crit Care Nurse. 2008;28(2):40.
1190	Reason for exclusion: Ineligible concept.
1191	289.Kim I, Geiger-Brown J, Trinkoff A, Muntaner C. Physically demanding
1192	workloads and the risks of musculoskeletal disorders in homecare
1193	workers in the USA. Health Soc Care Community. 2010;18(5):445-455.
1194	Reason for exclusion: Ineligible concept.
1195	290. Chadwick M, Titcomb P. Patient handling: getting the training right. Nurs
1196	Residential Care. 2007;9(10):490-492.
1197	Reason for exclusion: Ineligible concept.
1198	291. Sabbath EL, Yang J, Dennerlein JT, Boden LI, Hashimoto D, Sorensen G.
1199	Paradoxical Impact of a Patient-Handling Intervention on Injury Rate
1200	Disparity Among Hospital Workers. Am J Public Health.
1201	2019;109(4):618-625.
1202	Reason for exclusion: Ineligible concept.
1203	292. Sedlak CA. Overview and summary: nurse safety: have we addressed
1204	the risks? Online J Issues Nurs. 2004;9(3):4.
1205	Reason for exclusion: Ineligible concept.
1206	293.Teeple E, Collins JE, Shrestha S, Dennerlein JT, Losina E, Katz JN.
1207	Outcomes of safe patient handling and mobilization programs: A meta-
1208	analysis. Work. 2017;58(2):173-184.
1209	Reason for exclusion: Ineligible concept.
1210	294. Yoder A, Coley K, Harrison D, Wright K. Outcomes of a Safe Patient
1211	Handling Program Implementation. Am J Safe Patient Handling

1212	Movement. 2014;4(4):111-117.
1213	Reason for exclusion: Ineligible concept.
1214	295.Larsen AK, Holtermann A, Mortensen OS, Punnett L, Rod MH, Jørgensen
1215	MB. Organizing workplace health literacy to reduce musculoskeletal pain
1216	and consequences. BMC Nurs. 2015;14(1):1-13.
1217	Reason for exclusion: Ineligible concept.
1218	296.Alnaser MZ. Occupational musculoskeletal injuries in the health care
1219	environment and its impact on occupational therapy practitioners: a
1220	systematic review. Work. 2007;29(2):89-100.
1221	Reason for exclusion: Ineligible concept.
1222	297. Filek S, Brims M. Nurssafe Patient Handling & Movement Around the
1223	World: British Columbia (Canada)Past, Present, and Future. Am J Safe
1224	Patient Handling Movement. 2013;3(3):110-112.
1225	Reason for exclusion: Ineligible concept.
1226	298.van Wyk PM, Andrews DM, Weir PL. Nurse perceptions of manual patient
1227	transfer training: implications for injury. Work. 2010;37(4):361-373.
1228	Reason for exclusion: Ineligible concept.
1229	299.Krill C, Raven C, Staffileno BA. Moving from a Clinical Question To
1230	Research: The Implementation Of a Safe Patient Handling Program.
1231	Medsurg Nurs. 2012 Mar;21(2):104-116.
1232	Reason for exclusion: Ineligible concept.
1233	300. Rinds G. Moving and handling: part one. Nurs Residential Care.
1234	2007;9(6):260-262.
1235	Reason for exclusion: Ineligible concept.
1236	301. Sturman-Floyd M. Moving and handling: assessing the handler. Nurs
1237	Residential Care. 2013;15(2):98-101.
1238	Reason for exclusion: Ineligible concept.
1239	302. Chadwick M. More principles for a positive approach to manual handling.
1240	Nurs Residential Care. 2008;10(3):151-153.
1241	Reason for exclusion: Ineligible concept.
1242	303.van Wyk PM, Weir PL, Andrews DM. Manual patient transfers used most
1243	often by student and staff nurses are consistent with their perceptions of
1244	transfer training, and performance confidence. Work. 2015;50(2):249-
1245	260.
1246	Reason for exclusion: Ineligible population.

1247	304. Enos L. Making the Business Case to Initiate, Evaluate, and Sustain Safe
1248	Patient Handling Programs: Part 2. Am J Safe Patient Handling
1249	Movement. 2011;1(4):22-30.
1250	Reason for exclusion: Ineligible concept.
1251	305. Enos L. Making the Business Case to Initiate, Evaluate, and Sustain Safe
1252	Patient Handling Programs. Am J Safe Patient Handling Movement.
1253	2011;1(3):8-15.
1254	Reason for exclusion: Ineligible concept.
1255	306. Hanson H, Wagner M, Monopoli V, Keysor J. Low back pain in physical
1256	therapists: a cultural approach to analysis and intervention. Work.
1257	2007;28(2):145-151.
1258	Reason for exclusion: Ineligible concept.
1259	307. Stevenson JM. Looking forward by looking back: Helping to reduce work-
1260	related musculoskeletal disorders. Work. 2014;47(1):137-141.
1261	Reason for exclusion: Ineligible population.
1262	308. Kane J. Intervention needed for manual handling safety. Kai Tiaki Nurs
1263	NZ. 2015;21(10):18-20.
1264	Reason for exclusion: Ineligible concept.
1265	309. Enos L, Monaghan HM, Odum M. International Round Table Discussion:
1266	Safe Patient Handling & Mobility: what is Working, what Needs to
1267	Change, and Where does the Future Lie? Int J Safe Patient Handl Mobil.
1268	2019;9(3):122-127.
1269	Reason for exclusion: Ineligible concept.
1270	310.Gallagher A, Dube J, Wawzyniecki P. International Round Table
1271	Discussion: Engaging Patients and Family in Safe Patient Handling and
1272	Early Mobility Decision-Making. Int J Safe Patient Handl Mobil.
1273	2019;9(1):42-46.
1274	Reason for exclusion: Ineligible concept.
1275	311.Francis R, Fray M, Morley S. International Round Table Discussion: Does
1276	Federal, Government, or State Legislation Mandating SPHM Make a
1277	Difference to Compliance? Int J Safe Patient Handl Mobil. 2019;9(4):149-
1278	153.
1279	Reason for exclusion: Ineligible concept.
1280	312.Costa-Black KM, Durand M, Imbeau D, Baril R, Loisel P. Interdisciplinary
1281	team discussion on work environment issues related to low back

1282	disability: a multiple case study. Work. 2007;28(3):249-265.
1283	Reason for exclusion: Ineligible concept.
1284	313. Soares MM, Jacobs K, Selis K, Vanacker H, Hermans V, Mylle G, et al.
1285	Integration of ergonomics in the prevention policy of a large university
1286	hospital. Work. 2012;41:5917-5920.
1287	Reason for exclusion: Ineligible concept.
1288	314.Perlow E, Tunney N, Lucado A. Integrating Safe Patient Handling Into
1289	Physical Therapist Education: Reducing the Incidence of Physical
1290	Therapist Injury and Improving Patient Outcomes. J Phys Ther Educ.
1291	2016;30(2):32-37.
1292	Reason for exclusion: Ineligible concept.
1293	315. Soares MM, Jacobs K, Sharan D, Ajeesh P. Injury prevention in
1294	physiotherapists - a scientific review. Work. 2012;41:1855-1859.
1295	Reason for exclusion: Ineligible concept.
1296	316.Szeto G, Cheng A, Lee E, Schonstein E, Gross D. Implementing the Work
1297	Disability Prevention Paradigm Among Therapists in Hong Kong:
1298	Facilitators and Barriers. J Occup Rehabil. 2011;21(1):76-83.
1299	Reason for exclusion: Ineligible concept.
1300	317. Sofie J, Belza B, Young H. Health and safety risk at a skilled nursing
1301	facility: nursing assistants' perceptions. J Gerontol Nurs. 2003;29(2):13-
1302	21.
1303	Reason for exclusion: Ineligible concept.
1304	318. Powell-Cope G, Hughes NL, Sedlak C, Nelson A. Faculty perceptions of
1305	implementing an evidence-based safe patient handling nursing
1306	curriculum module. Online J Issues Nurs. 2008;13(3):13p.
1307	Reason for exclusion: Ineligible concept.
1308	319.Kotowski S, Davis K. Factors Impacting the Patient Migration in Hospital
1309	Beds: Pathway to Reduce Patient Handling Injuries. Int J Safe Patient
1310	Handl Mobil. 2019;9(3):102-110.
1311	Reason for exclusion: Ineligible concept.
1312	320.Adams J. Factors Associated with Manual Handling Injuries: a Case-
1313	Control Study Employing the Health Belief Model. Am J Safe Patient
1314	Handling Movement. 2013;3(3):84-93.
1315	Reason for exclusion: Ineligible concept.

1316	321.Gallagher SM. Exploring the Relationship between Obesity, Patient
1317	Safety, and Caregiver Injury. Am J Safe Patient Handling Movement.
1318	2011;1(2):8-12.
1319	Reason for exclusion: Ineligible concept.
1320	322.Rinds G. Exploring recent developments in manual handling. Nurs
1321	Residential Care. 2009;11(6):312-315.
1322	Reason for exclusion: Ineligible concept.
1323	323. Nelson A, Baptiste AS. Evidence-based practices for safe patient handling
1324	and movement. Orthop Nurs. 2006;25(6):366-379.
1325	Reason for exclusion: Ineligible concept.
1326	324. Egerod I. Evidence-based practice among Danish critical care nurses.
1327	Connect World Crit Care Nurs. 2005;4(3):103-104.
1328	Reason for exclusion: Ineligible concept.
1329	325.Knibbe Hjj, Knibbe Ne, Heitink, Diede E. B. L. Evaluating Different
1330	Methods of Showering and Washing Patients: Assessing Ergonomic,
1331	Time, and Quality Aspects. Am J Safe Patient Handling Movement.
1332	2016;6(2):49-64.
1333	Reason for exclusion: Ineligible concept.
1334	326.Potter M, Jones S. Entry-level physiotherapists' strategies to lower
1335	occupational injury risk in physiotherapy: a qualitative study. Physiother
1336	Theory Pract. 2006;22(6):329-336.
1337	Reason for exclusion: Ineligible concept.
1338	327.Enos L, Eldredge D, Rockefeller K. Educational Case Report: a Safe
1339	Patient Handling and Mobilization Training Program in an Academic
1340	Medical Center. Am J Safe Patient Handling Movement. 2016;6(3):120-
1341	129.
1342	Reason for exclusion: Ineligible concept.
1343	328. Hignett S, Crumpton E. Development of a patient handling assessment
1344	tool. Int J Ther Rehabil. 2005;12(4):178-181.
1345	Reason for exclusion: Ineligible concept.
1346	329.Powell-Cope G, Haun J, Rugs D. Description of a Social Marketing
1347	Framework for Implementing an Evidence-Based Safe Patient Handling
1348	Program. Am J Safe Patient Handling Movement. 2013;3(2):64-72.
1349	Reason for exclusion: Ineligible concept

1350	330.Mansour M. Current assessment of patient safety education. Br J Nurs.
1351	2012;21(9):536-543.
1352	Reason for exclusion: Ineligible population.
1353	331. Hooper J, Charney W. Creation of a safety culture: reducing workplace
1354	injuries in a rural hospital setting. AAOHN J. 2005;53(9):394-398.
1355	Reason for exclusion: Ineligible concept.
1356	332.Knibbe Hjj, Waaijer E. Covid-19 and Safe Patient Handling and Mobility in
1357	the Netherlands. Int J Safe Patient Handl Mobil. 2020;10(2):80-82.
1358	Reason for exclusion: Ineligible concept.
1359	333.Monaghan HM. Considering Hospital Transporters in a Safe Patient
1360	Handling Program. Am J Safe Patient Handling Movement. 2014;4(3):93-
1361	96.
1362	Reason for exclusion: Ineligible concept.
1363	334. Soares MM, Jacobs K, Macdonald W. Conceptual framework for
1364	development of a toolkit for prevention of work-related musculoskeletal
1365	disorders. Work. 2012;41:3933-3936.
1366	Reason for exclusion: Ineligible concept.
1367	335.Enos L, Monaghan HM, Thomas S. Comprehensive Review of SPHM
1368	Articles from 2005-2016. Am J Safe Patient Handling Movement.
1369	2016;6:1-28.
1370	Reason for exclusion: Ineligible concept.
1371	336.Surrey LR, Fineran B, Howerton B, Mock A, Naylor D, Wiegand-Swihart T.
1372	Comparison of a simulated versus work environment lifting task on L5/S1
1373	compression forces. Work. 2003;21(3):265-270.
1374	Reason for exclusion: Ineligible population.
1375	337. Haas AD, Hunter DA, Howard NL. Bringing a structural perspective to
1376	work: Framing occupational safety and health disparities for nursing
1377	assistants with work-related musculoskeletal disorders. Work.
1378	2018;59(2):211-229.
1379	Reason for exclusion: Ineligible concept.
1380	338.Harwood KJ. Blazing a New Trail: Advocacy for Safe Patient Handling and
1381	Mobility. Am J Safe Patient Handling Movement. 2015;5(1):21-26.
1382	Reason for exclusion: Ineligible concept.
1383	339.Nourollahi M, Afshari D, Dianat I. Awkward trunk postures and their
1384	relationship with low back pain in hospital nurses. Work.

1385	2018;59(3):317-323.
1386	Reason for exclusion: Ineligible setting.
1387	340.Rockefeller K. Appraising Evidence about Safe Patient Handling and
1388	Mobility. Am J Safe Patient Handling Movement. 2016;6(1):33-38.
1389	Reason for exclusion: Ineligible concept.
1390	341. Wanless S. Applying Theories of Health Behavior and Change to Moving
1391	and Handling Practice. Int J Safe Patient Handl Mobil. 2017;7(3):105-
1392	109.
1393	Reason for exclusion: Ineligible concept.
1394	342.AORN. ANA releases national standards for safe patient handling and
1395	mobility. AORN J. 2013;98(2):4.
1396	Reason for exclusion: Ineligible concept.
1397	343. Trossman S. ANA leads initiative to develop national safe patient
1398	handling standards. Am Nurse. 2012;44(5):4.
1399	Reason for exclusion: Ineligible concept.
1400	344.ANA. ANA leading the effort to develop national safe patient handling
1401	standards. Am Nurse. 2012;44(3):14.
1402	Reason for exclusion: Ineligible concept.
1403	345.Shaw L, Domanski S, Freeman A, Hoffele C. An investigation of a
1404	workplace-based return-to-work program for shoulder injuries. Work.
1405	2008;30(3):267-276.
1406	Reason for exclusion: Ineligible concept.
1407	346. Soares MM, Jacobs K, Knibbe JJ, Knibbe NE. An international validation
1408	study of the care thermometer: a tool supporting the quality of
1409	ergonomic policies in health care. Work. 2012;41:5639-5641.
1410	Reason for exclusion: Ineligible concept.
1411	347.Roll S, Roll SC, Evans KD, Hutmire CD, Baker JP. An analysis of
1412	occupational factors related to shoulder discomfort in diagnostic medical
1413	sonographers and vascular technologists. Work. 2012;42(3):355-365.
1414	Reason for exclusion: Ineligible concept.
1415	348. Tompa E, Dolinschi R, de Oliveira C, III AB, Irvin E. A systematic review
1416	of workplace ergonomic interventions with economic analyses. J Occup
1417	Rehabil. 2010;20(2):220-234.
1418	Reason for exclusion: Ineligible concept.

1419	349.Harwood KJ, Darragh AR, Campo M, Rockefeller K, Scalzitti DA. A
1420	Systematic Review of Safe Patient Handling and Mobility Programs to
1421	Prevent Musculoskeletal Injuries in Occupational and Physical Therapists
1422	and Assistants. Int J Safe Patient Handl Mobil. 2018;8(1):46-56.
1423	Reason for exclusion: Ineligible concept.
1424	350.Harwood KJ, Scalzitti DA, Campo M, Darragh AR. A Systematic Review of
1425	Safe Patient Handling and Mobility Programs to Improve Patient
1426	Outcomes in Rehabilitation. Am J Safe Patient Handling Movement.
1427	2016;6(4):141-150.
1428	Reason for exclusion: Ineligible concept.
1429	351.Deen M, Gibson L, Strong J. A survey of occupational therapy in
1430	Australian work practice. Work. 2002;19(3):219-230.
1431	Reason for exclusion: Ineligible concept.
1432	352.Dropkin J, Moline J, Power PM, Kim H. A qualitative study of health
1433	problems, risk factors, and prevention among Emergency Medical Service
1434	workers. Work. 2015;52(4):935-951.
1435	Reason for exclusion: Ineligible concept.
1436	353.Rue CA, Rayson MP, Walker EF, Doherty J, Thompson J, Myers SD, et al.
1437	A job task analysis to describe the physical demands of specialist
1438	paramedic roles in the National Ambulance Resilience Unit (NARU). Work.
1439	2019;63(4):547-557.
1440	Reason for exclusion: Ineligible concept.
1441	354. Noonan J, Wagner SL. A biopsychosocial perspective on the management
1442	of work-related musculoskeletal disorders. AAOHN J. 2010;58(3):105-
1443	114.
1444	Reason for exclusion: Ineligible concept.
1445	355.Kurowski A, Boyer J, Fulmer S, Gore R, Punnett L. Changes in ergonomic
1446	exposures of nursing assistants after the introduction of a safe resident
1447	handling program in nursing homes. Int J Ind Ergon. 2012;42(6):525-
1448	32.
1449	Reason for exclusion: Ineligible concept.
1450	356.Kneafsey R, Haigh C. Moving and handling rehabilitation patients: A
1451	survey of nurses' views. Int J Ther Rebabil. 2009;16(8):431-41.
1452	Reason for exclusion: Ineligible concept.

1453	357. Prairie J, Corbeil P. Paramedics on the job: Dynamic trunk motion
1454	assessment at the workplace. Appl Ergon. 2014;45(4):895-903.
1455	Reason for exclusion: Ineligible concept.
1456	358.Treadwell L. Safe manual handling practices in the UK and USA: More
1457	than just a training exercise. Int J Ther Rehabil. 2013;20(7):326-7.
1458	Reason for exclusion: Ineligible concept.
1459	359.Taylor J, Sims J, Haines TP. The emergent relevance of care staff
1460	decision-making and situation awareness to mobility care in nursing
1461	homes: an ethnographic study. J Adv Nurs. 2014;70(12):2767-78.
1462	Reason for exclusion: Ineligible concept.
1463	360.Darragh AR, Shiyko M, Margulis H, Campo M. Effects of a safe patient
1464	handling and mobility program on patient self-care outcomes. Am J of
1465	Occup Ther. 2014;68(5):589-96.
1466	Reason for exclusion: Ineligible concept.
1467	361. Piccenna L, Lannin NA, Scott K, Bragge P, Gruen R. Guidance for
1468	community-based caregivers in assisting people with moderate to severe
1469	traumatic brain injury with transfers and manual handling: Evidence and
1470	key stakeholder perspectives. Health Soc Care Community.
1471	2017;25(2):458-65.
1472	Reason for exclusion: Ineligible population.
1473	362.Risor BW, Casper SD, Andresen LL, Sorensen J. A multi-component
1474	patient-handling intervention improves attitudes and behaviors for safe
1475	patient handling and reduces aggression experienced by nursing staff: A
1476	controlled before-after study. Appl Ergon. 2017;60:74-82.
1477	Reason for exclusion: Ineligible concept.
1478	363. Fragala G. Bed care for patients in palliative settings: Considering risks
1479	to caregivers, and bed surfaces. Int J Palliat Nurs. 2015;21(2):66-70.
1480	Reason for exclusion: Ineligible concept.
1481	364.Rodrigues FF, de Oliveira Sato T. Effectiveness of ergonomic training to
1482	reduce physical demands and musculoskeletal symptoms - An overview
1483	of systematic reviews. Int J Ind Ergon. 2019;74.
1484	Reason for exclusion: Ineligible concept.
1485	365.Rinds G. Moving and handling, part two: Risk management. Nurs
1486	Residential Care. 2007;9(7):306-9.
1487	Reason for exclusion: Ineligible concept.

1488	366.Hall N, Bryer M. Moving and handling education: Educating the
1489	experienced practitioner. Int J Ther Rehabil. 2005;12(10):449-53.
1490	Reason for exclusion: Ineligible concept.
1491	367.Rice MS, Dusseau JM, Kopp Miller B. A questionnaire of musculoskeletal
1492	injuries associated with manual patient lifting in occupational therapy
1493	practitioners in the state of Ohio. Occup Ther Health Care. 2011;25:95-
1494	107.
1495	Reason for exclusion: Ineligible concept.
1496	368.Nelson A, Harwood KJ, Tracey CA, Dunn KL. Myths and Facts About Safe
1497	Patient Handling in Rehabilitation. Rehabil Nurs. 2008;33(1):10-7.
1498	Reason for exclusion: Ineligible concept.
1499	369.Donnelly C, Macmillan MS. Devising and evaluating a model for teaching
1500	therapeutic handling and moving skills to complementary therapists.
1501	Complement Ther Clin Pract. 2007;13(3):201-9.
1502	Reason for exclusion: Ineligible concept.
1503	370. Habibi E, Pourabdian S, Atabaki AK, Hoseini M. Evaluation of work-
1504	related psychosocial and ergonomics factors in relation to low back
1505	discomfort in emergency unit nurses. Int J Prev Med. 2012;3(8):564-8.
1506	Reason for exclusion: Ineligible concept.
1507	371.Olson DL, King PM. Estimated energy expenditure of nursing assistants in
1508	long term care. Work. 2012;42(3):331-40.
1509	Reason for exclusion: Ineligible concept.
1510	372.Darragh AR, Campo MA, Frost L, Miller M, Pentico M, Margulis H. Safe-
1511	patient-handling equipment in therapy practice: Implications for
1512	rehabilitation. Am J Occup Ther. 2013;67(1):45-53.
1513	Reason for exclusion: Ineligible concept.
1514	373.Dyrkacz AR, Mak LY, Heck CS. Work-related injuries in Canadian
1515	occupational therapy practice. Can J Occup Ther. 2012;79(4):237-47.
1516	Reason for exclusion: Ineligible concept.
1517	374. Mawston GA, Boocock MG. The effect of lumbar posture on spinal loading
1518	and the function of the erector spinae: Implications for exercise and
1519	vocational rehabilitation. N Z J Physiother. 2012;40(3):135-40.
1520	Reason for exclusion: Ineligible concept.

1521	375.Nelson A. Promoting Safe Patient Handling. Rehabil Nurs.
1522	2008;33(1):42-43.
1523	Reason for exclusion: Ineligible concept.
1524	376.Parry JS. Guidance on manual handling in physiotherapy. Aqualines News
1525	Hydro Assoc Chart Physiother. 2008;2:15-6.
1526	Reason for exclusion: Ineligible concept.
1527	377. Arnold M, Radawiec S, Campo M, Wright LR. Changes in functional
1528	independence measure ratings associated with a safe patient handling
1529	and movement program. Rehabil Nurs. 2011;36(4):138-44.
1530	Reason for exclusion: Ineligible population.
1531	378.Collins JW, Bell JL, Gronqvist R. Developing evidence-based interventions
1532	to address the leading causes of workers' compensation among
1533	healthcare workers. Rehabil Nurs. 2010;35(6):225-35.
1534	Reason for exclusion: Ineligible concept.
1535	379.Haladay DE, Blorstad AL, McBrief NM, Denegar CR, Lengerich EJ. Back
1536	pain among health care workers in Pennsylvania 2002-2006. Work.
1537	2012;41(1):93-8.
1538	Reason for exclusion: Ineligible concept.
1539	380. White J, Gray H. Factors influencing the effectiveness of manual handling
1540	education. Int J Ther Rehabil. 2004;11(5):226-31.
1541	Reason for exclusion: Ineligible concept.
1542	381.van Zoest GG, van den Berg, H. T., Holtkamp FC. Three-dimensionality
1543	of contact forces during clinical manual examination and treatment: a
1544	new measuring system. Clin Biomech. 2002;17:719-22.
1545	Reason for exclusion: Ineligible concept.
1546	382. Li J, Sommerich CM, Chipps E, Lavender SA, Stasny EA. A framework for
1547	studying risk factors for lower extremity musculoskeletal discomfort in
1548	nurses. Ergonomics 2020;63(12):1535-1550.
1549	Reason for exclusion: Ineligible concept.
1550	383. Brooker KJ, Vikan M, Thyli B. A Qualitative Exploratory Study of
1551	Norwegian OR Nurses' Patient Positioning Priorities. AORN J
1552	2020;111(2):211-220.
1553	Reason for exclusion: Ineligible concept.
1554	384. Storey BJ, Verkerk MM, Hashtroudi A, GoldingWood D. A systematic
1555	review of interventions to prevent work-related musculoskeletal

1556	disorders in ENT surgeons. J Laryngol Otol 2021.
1557	Reason for exclusion: Ineligible concept.
1558	385. Muller A, Corbeil P. Back loading estimation during team handling: Is
1559	the use of only motion data sufficient? PloS one 2020;15(12).
1560	Reason for exclusion: Ineligible concept.
1561	386. XU H, SHI J, NI W, Xin Z, WANG A, ZHENG X. Biomechanical Analysis on
1562	Optimization of Manual Handing Technique in Manual Lifting and
1563	Lowering Tasks. Journal of Beijing Sport University 2013;36(11):76-36.
1564	Reason for exclusion: Not English Language.
1565	387.Callihan ML, Eyer JC, McCoy CJ, Dailey AM, Diket KM, Robinson AT, et al.
1566	Development and Feasibility Testing of a Contextual Patient Movement
1567	Intervention. Journal of emergency nursing 2021;47(1):101.
1568	Reason for exclusion: Ineligible population.
1569	388. Claycomb M. Guidelines to Safe Patient Transfers. Rehab Management:
1570	The Interdisciplinary Journal of Rehabilitation 2015;28(1):14-17.
1571	Reason for exclusion: Ineligible concept.
1572	389. Bernardes JM, MonteiroPereira PE, GomezSalgado J, RuizFrutos C, Dias
1573	A. Healthcare workers' knowledge for safe handling and moving of the
1574	patient. International journal of occupational safety and ergonomics :
1575	JOSE 2021:1-20.
1576	Reason for exclusion: Ineligible concept.
1577	390. Kurowski A, Pransky G, Punnett L. Impact of a Safe Resident Handling
1578	Program in Nursing Homes on Return-to-Work and Re-injury Outcomes
1579	Following Work Injury. J Occup Rehabil 2019;29(2):286-294.
1580	Reason for exclusion: Ineligible concept.
1581	391. Lee SJ, Stock L, Michalchuk V, Adesoye K, Mullen K. Impact of California
1582	Safe Patient Handling Legislation: Health Care Workers' Perspectives.
1583	Workplace health & safety 2021;69(3):124-133.
1584	Reason for exclusion: Ineligible concept.
1585	392. Improving patient and health care provider safety: task force develops
1586	recommendations on patient handling. PT: Magazine of Physical Therapy
1587	2005;13(4):48-52.
1588	Reason for exclusion: Ineligible concept.
1589	393. Alghadir AH, AlAbbad H, Buragadda S, Iqbal A. Influence of work-related

safety and health guidelines on knowledge and prevalence of

1591 occupational back pain among rehabilitation nurses in saudi arabia: A 6-1592 month follow-up study. International Journal of Environmental Research 1593 and Public Health 2021;18(16). 1594 Reason for exclusion: Ineligible concept. 1595 394. Gilthorpe J, Fairbaurn S, Taylor S, Hutchison A, Ford C, Tweedlie J. 1596 Interprofessional education for practice: moving and handling for people 1597 with complex needs in contemporary healthcare. Physiotherapy (United 1598 Kingdom). Conference: Physiotherapy UK Conference 2019. Birmingham 1599 United Kingdom. 107(Supplement 1) (pp e170-e171); Elsevier Ltd; 1600 2020. 1601 Reason for exclusion: Ineligible population. 1602 395. Schroder C, Nienhaus A. Intervertebral disc disease of the lumbar spine 1603 in health personnel with occupational exposure to patient handling-a 1604 systematic literature review and meta-analysis. International Journal of 1605 Environmental Research and Public Health 2020;17(1-20). 1606 Reason for exclusion: Ineligible concept. 1607 396. Horgan D, Purtill H, O'Sullivan P, Dankaerts W, O'Sullivan K. 1608 Investigation of the impact of people manual handling training on back 1609 beliefs. Man Ther 2016;25. 1610 Reason for exclusion: Ineligible concept. 1611 397. Lecca LI, Fabbri D, Portoghese I, Pilia I, Meloni F, Marcias G, et al. 1612 Manual handling of patients: role of kinesiophobia and catastrophizing in 1613 health workers with chronic low back pain. European journal of physical 1614 and rehabilitation medicine 2020;56(3):307-312. 1615 Reason for exclusion: Ineligible concept. 1616 398. Tullar JM, Brewer S, Amick III BC, Irvin E, Mahood Q, Pompeii LA, et al. 1617 Occupational Safety and Health Interventions to Reduce Musculoskeletal 1618 Symptoms in the Health Care Sector. J Occup Rehabil 2010;20(2):199-1619 219. 1620 Reason for exclusion: Duplicate. 1621 399. Dewey AJ, Olkowski BF, Stolfi AM. On "Safe patient handling perceptions and practices . . . ". Phys Ther 2014;94(7):1054-1055. 1622

Reason for exclusion: Ineligible study design.

1624	400. Wyatt D. Our Shared Purpose: Workplace Safety. AORN J
1625	2020;112(4):325-326.
1626	Reason for exclusion: Ineligible concept.
1627	401. SAFE LIFTING TECHNIQUES. Rehab Management: The Interdisciplinary
1628	Journal of Rehabilitation 2010;23(6):32.
1629	Reason for exclusion: Ineligible concept.
1630	402. Silberstein N. Safe Patient Handling. Physical Therapy Products
1631	2009;20(6):12-15.
1632	Reason for exclusion: Ineligible concept.
1633	403. Satusky MJ. Safe Patient Handling and Mobility in the Orthopaedic
1634	Setting. Orthopedic nursing 2021;40(2):60-63.
1635	Reason for exclusion: Ineligible concept.
1636	404. Lee SJ, Kang KJ, Lee JH. Safe patient handling legislation and changes
1637	in programs, practices, perceptions, and experience of musculoskeletal
1638	disorders by hospital characteristics: A repeated cross-sectional survey
1639	study. Int J Nurs Stud 2021;113:103791.
1640	Reason for exclusion: Ineligible concept.
1641	405. Olkowski BF, Stolfi AM. Safe Patient Handling Perceptions and Practices:
1642	A Survey of Acute Care Physical Therapists. Phys Ther 2014;94(5):682-
1643	695.
1644	Reason for exclusion: Ineligible concept.
1645	406. Boninger ML. Safe Transfer Technique. Archives of Physical Medicine &
1646	Rehabilitation 2013;94(12):2579-2580.
1647	Reason for exclusion: Ineligible concept.
1648	407. Long F. Safer Patient Handling: Stat! Rehab Management: The
1649	Interdisciplinary Journal of Rehabilitation 2019;32(6):32-35.
1650	Reason for exclusion: Ineligible concept.
1651	408. Zhou J, Wiggermann N. The effects of hospital bed features on physical
1652	stresses on caregivers when repositioning patients in bed. Appl Ergon
1653	2021;90.
1654	Reason for exclusion: Ineligible concept.
1655	409. Vieira ER, Kumar S. Working Postures: A Literature Review. J Occup
1656	Rehabil 2004;14(2):143-159.
1657	Reason for exclusion: Ineligible population.

## 1658 <u>Data extraction instrument</u>

Author,	Aims/	Study type	Setting	Participant	Participant:	Patient	Outcome measure:	Date of
Year,	purpose			profession	Age,	population	Domain,	extraction
Country of origin					Gender,		Measurement tools	
					Years of experience,			
					Sample size			

## Outcome measure count and references

Outcome measure	Measurement tool	Relative Count		
Kinematics	Video	3 [29,33,40]		
	Goniometer	3 [26,35,40]		
	Inclinometer	2 [22,24]		
	Observation	2 [31,33]		
	Lumbar motion monitor	2 [23,26]		
	OPTOTRAK	1 [27]		
	Tri-axial accelerometer	1 [41]		
	Tri-axial goniometer	1 [26]		
	Vicon	1 [28]		
	Posture Dosimeter	1 [33]		
	Photo	1 [40]		
Physical demands	EMG	7 [23,25,26,28,35,37,50]		
	Borg scale	3 [19,37,44]		
	Video	1 [25]		
	Accelerometer	1 [41]		
	Interviews	1 [33]		
	Heart rate	1 [21]		
	VAS	1 [38]		
Tasks performed	Observation	7 [21,22,33,46,51,52]		
	Videos	2 [33,42]		
	Survey	2 [45,49]		
	Ergonomic Assessment	1 [29]		
Kinetics	Digitised video	5 [27,36,37,40,50]		
	Force platforms	3 [36-38]		
	Force transducers	3 [36,37,43]		
	Tri-axial load cell	1 [2]		
	Physical workload index	1 [31]		
Safety	Observation	4 [19,30,33,52]		
	Staff training	2 [18,44]		
	Video	2 [20,30]		
	Photo	1 [20]		
	Survey	1 [45]		
WRMSD	Incidence	3 [18,19,39]		
	Risk factors	1 [20]		
Staff perceptions/ opinions	Focus groups	3 [47,48,52]		
•	Interviews	2 [34,46]		
	Questionnaire	1 [30]		
	24-h survey	1 [49]		