

Dose-response modelling of resistance exercise across outcome domains in strength and conditioning: a meta-analysis.

SWINTON, P.A., SCHOENFELD, B.J. and MURPHY, A.

2024

© The Author(s) 2024. The version of record of this article, first published in *Sports Medicine*, is available online at Publisher's website: <https://doi.org/10.1007/s40279-024-02006-3>
Supplementary materials are appended after the main text of this document.



Dose–Response Modelling of Resistance Exercise Across Outcome Domains in Strength and Conditioning: A Meta-analysis

Paul Alan Swinton¹ · Brad J. Schoenfeld² · Andrew Murphy^{1,3}

Accepted: 14 February 2024 / Published online: 23 April 2024
© The Author(s) 2024

Abstract

Background Resistance exercise is the most common training modality included within strength and conditioning (S&C) practice. Understanding dose–response relationships between resistance training and a range of outcomes relevant to physical and sporting performance is of primary importance for quality S&C prescription.

Objectives The aim of this meta-analysis was to use contemporary modelling techniques to investigate resistance-only and resistance-dominant training interventions, and explore relationships between training variables (frequency, volume, intensity), participant characteristics (training status, sex), and improvements across a range of outcome domains including maximum strength, power, vertical jump, change of direction, and sprinting performance.

Methods Data were obtained from a database of training studies conducted between 1962 and 2018, which comprised healthy trained or untrained adults engaged in resistance-only or resistance-dominant interventions. Studies were not required to include a control group. Standardized mean difference effect sizes were calculated and interventions categorized according to a range of training variables describing frequency (number of sessions per week), volume (number of sets and repetitions performed), overall intensity (intensity of effort and load, categorised as low, medium or high), and intensity of load (represented as % of one-repetition maximum [1RM] prescribed). Contemporary modelling techniques including Bayesian mixed-effects meta-analytic models were fitted to investigate linear and non-linear dose-responses with models compared based on predictive accuracy.

Results Data from a total of 295 studies comprising 535 groups and 6,710 participants were included with analyses conducted on time points ≤ 26 weeks. The best performing model included: duration from baseline, average number of sets, and the main and interaction effects between outcome domain and intensity of load (% 1RM) expressed non-linearly. Model performance was not improved by the inclusion of participant training status or sex.

Conclusions The current meta-analysis represents the most comprehensive investigation of dose–response relationships across a range of outcome domains commonly targeted within strength and conditioning to date. Results demonstrate the magnitude of improvements is predominantly influenced by training intensity of load and the outcome measured. When considering the effects of intensity as a % 1RM, profiles differ across outcome domains with maximum strength likely to be maximised with the heaviest loads, vertical jump performance likely to be maximised with relatively light loads ($\sim 30\%$ 1RM), and power likely to be maximised with low to moderate loads (40–70% 1RM).

1 Introduction

Resistance exercise is established as one of the most effective training modalities within strength and conditioning (S&C) [1, 2]. Researchers have shown that improvements in strength and power can be transferred to a range of important activities associated with sports performance including sprinting and jumping [1, 3, 4]. Results from a recent large meta-analysis also highlighted the importance of training specificity, in relation to the imposed demands of training, with the greatest improvements made when

Key Points

Results from the current large-scale meta-analysis demonstrate that intervention duration combined with manipulation of training intensity are the most relevant factors in altering magnitude of improvement following resistance training.

Different % 1RM intensity profiles exist across outcome domains, highlighting the importance of selecting domain-specific loads.

Across the heterogeneous research, training status, sex and training frequency were shown to provide limited predictive capabilities. These factors should still be considered when developing holistic periodised strength and conditioning programs with further research required.

matching the training stimulus and the outcomes assessed (e.g., traditional resistance exercise with heavy loads matched with one-repetition maximum (1RM) assessment) [5]. Beyond specificity, appropriate prescription of resistance exercise requires consideration of a range of acute program variables including volume, intensity, frequency and potentially more subtle variables including exercise selection and exercise order [6]. There have been attempts to provide general recommendations of training dose to maximise a range of outcome domains including strength, hypertrophy, power and muscular endurance [6]. Previous researchers have also indicated that the training status of participants quantified as years of resistance training experience may play an important role and interact with training dose [7, 8]. Greater understanding of dose–response relationships across a range of factors including the training modality, outcome domain, participant characteristics (e.g., training status) and length of intervention is key for continued development of resistance exercise and the desire to avoid over- or under-loading.

Several systematic reviews have investigated training dose within S&C, with most focusing on development of strength and hypertrophy [7–17]. Seminal studies conducted by Rhea and colleagues [7–9] were among the first to use meta-analytic techniques to quantify dose–response relationships. Initial research from Rhea et al. [9] incorporated data from 16 studies to compare single versus multiple sets and concluded that performance of three sets was more effective than a single set. Follow-up research was conducted by the authors by (1) substantially increasing the number of studies included in the analysis [7], and (2) focusing on higher-level athletes [8]. The large-scale

meta-analysis conducted by Rhea et al. [7] included data from 140 studies and provided further support for the superiority of multiple sets, with four sets per muscle group concluded to produce the greatest improvements [7]. The results from the comprehensive analysis indicated different dose–response relationships for trained and untrained participants, with a higher intensity of load (80% 1RM) and a frequency of 2 days per week judged most effective for trained participants; and lower intensity of load (60% 1RM) and a frequency of 3 days per week judged most effective for untrained participants. In the follow-up meta-analysis restricted to 37 studies including competitive athletes, Peterson et al. [8] concluded higher volumes (eight sets per muscle group) and intensity of load intensities (85% 1RM) resulted in the largest effects with no differences found between frequencies of 2 or 3 days per week. Collectively, the work from Rhea and colleagues synthesised results from almost 200 studies and confirmed the existence of dose–response relationships for the development of strength and the likely moderation by participant characteristics including training status [18].

The most recent meta-analyses investigating dose–response relationships in S&C have tended to focus on the manipulation of a smaller number of program variables and restricting analyses to more homogeneous studies [10–17]. Meta-analyses from Grgic et al. [12] and Ralston et al. [13] investigated the effects of manipulating training frequency on strength improvements and incorporated data from 22 studies directly comparing frequencies of 1–4 days per week, and from 12 studies directly comparing frequencies of 1–3 days per week, respectively. Grgic et al. [12] concluded that higher training frequencies resulted in greater improvements in strength, but that these increases appeared to be primarily mediated through increased weekly volume. Similar conclusions were presented by Ralston et al. [13] showing that when resistance volume was equated across low (1 day week⁻¹), medium (2 days week⁻¹) or high (≥ 3 days week⁻¹) frequencies, similar increases in strength were obtained for both isolation and multi-joint exercises. Restricting analyses to trained athletes, Cuthbert et al. [17] identified no differences in improvements to either lower (effect size: 0.06; CI – 0.20, 0.32) or upper (effect size: 0.09; CI – 0.17, 0.35) body strength when comparing training frequencies during competitive periods.

Another smaller and more focused meta-analysis was conducted by Schoenfeld et al. [15] comparing low-load ($\leq 60\%$ 1RM) versus high-load ($> 60\%$ 1RM) resistance training. The meta-analysis included 21 relatively homogeneous studies tending to focus on exercises including the bench press, knee extension and leg press, all performed to momentary muscular failure. Results from the meta-analysis identified greater improvements in 1RM strength with

high-load resistance training; however, the transfer of training to isometric strength testing showed marginally greater effects favouring higher versus lower loads (effect size: 0.16; 95% CI – 0.06, 0.37) [15]. Moreover, similar improvements were observed in muscle hypertrophy across conditions. The results presented by Schoenfeld et al. [15] highlight the potential for different dose–response relationships across outcomes routinely targeted in S&C.

Whilst both large meta-analyses comprising heterogeneous studies and smaller, more focussed meta-analyses present different strengths and weaknesses, there is likely a benefit in simultaneously modelling dose–response relations across a broader range of outcome domains than has been investigated previously. The response to any training program is ultimately a complex interaction of a range of variables and participant characteristics; however, large modelling analyses have the potential to identify general trends that provide researchers and practitioners with important information on which to design more specific programs. With this perspective in mind, the purpose of this encompassing meta-analysis was to use contemporary modelling techniques to investigate resistance training interventions and explore relationships between training variables (e.g., frequency, volume, overall intensity, and intensity of load), participant characteristics (training status, sex), and a range of outcome domains including maximum strength, power, vertical jump, change of direction (COD), and sprinting performance, which are key targets in S&C program design.

2 Methods

2.1 Overview of Meta-analysis

The meta-analysis was conducted on a database of S&C training studies with analyses restricted to interventions that comprised resistance training only, or combined interventions where resistance training accounted for more than half of the training volume (e.g., resistance combined with plyometrics, speed, COD or power training). The database included information describing outcome variables, participant characteristics, training-dose parameters along with baseline and follow-up means and standard deviations, as has been described elsewhere [1]. The information was used to calculate intervention-only (e.g., non-controlled) effect sizes designed to draw inferences from indirect comparisons. To conduct the meta-analysis, sequential hierarchical models were fitted to account for dependencies (e.g., reporting of multiple outcomes at multiple time points from the same study) and structure within the data (e.g., time points nested in outcomes, nested in studies). Participant, training dose, and intervention characteristics were sequentially added to

models to identify the most influential factors whilst monitoring changes caused by underlying associations among the variables. The focus of this meta-analysis was to quantify and describe the general influence of training dose across a range of outcomes commonly investigated in S&C studies.

2.2 Search Strategy and Reporting

The present review was conducted as a follow-up to a larger review that featured a broader range of training interventions [1]. The search for the original review was performed using Embase, Medline, Web of Science, SPORTDiscus and Google Scholar. Hand searching of relevant journals including *Medicine and Science in Sports and Exercise*, the *Journal of Strength and Conditioning Research*, and *Research Quarterly* was also conducted. Database search terms were included to identify various training modes and a range of outcome measures. The following keywords and phrases were combined with Boolean operators; “strength” OR “resistance” OR “sprint” OR “plyometric” OR “exercise” AND “intervention” OR “training” OR “program” OR “programme” AND “1RM” OR “repetition maximum” OR “speed” OR “velocity” OR “power” OR “jump” OR “change of direction” OR “agility” OR “acceleration” OR “rate of force development”. No restriction was placed on the date of the study, with searching conducted in January 2018. Reporting of this review was guided by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2020 [19] statement, with the checklist included for transparency (Online Supplementary Material (OSM) 1). Risk of bias assessment was not conducted.

2.3 Inclusion Criteria and Data Extraction

Inclusion and exclusion criteria for the current meta-analysis were set to include as many relevant S&C training studies as possible. Inclusion criteria comprised: (1) any resistance training or majority resistant training-based study ≥ 4 weeks; (2) healthy trained or untrained participants with a mean age between 14 and 60 years; (3) training group with a minimum of four participants; (4) pre- and post-training means and standard deviations; and (5) sufficient information provided to quantify training intensity, volume, and frequency. Studies did not require a control or comparator group to be included. A standardised extraction codebook was developed using Microsoft Excel, with data extracted and coded independently by four researchers in duplicate, with one reviewer (AM) completing extraction for all studies to provide consistency. Study selection followed a two-stage selection strategy (title or abstract screen and full-text screen) undertaken primarily by AM and two researchers as part of the original larger review during 2018–2020. The independent screeners convened at

the end of each screening stage to resolve any discrepancies. Data regarding the study (authors, year, total number of active intervention groups); participant characteristics (final study n, sex, training status, and age); outcome domain (maximum strength, power, jump performance, and sprinting performance); training dose (overall intensity, intensity of load, volume, frequency, number of exercises, number of sets, number of repetitions), and pre- and post-training means and standard deviations were obtained. The definitions used to categorise outcome domains included: (1) maximum strength: a measure of maximum force production where time was not limited (e.g., 1–6 repetition maximum, isometric mid-thigh pull, peak torque); (2) power: a direct measurement (e.g., Wingate test) or indirect estimation (e.g., vertical jump) of mechanical power output measured in Watts (absolute and normalised relative to body mass); (3) jump performance: measure of jump height or distance; (4) sprint performance: a measurement of the time to complete a specified linear distance or the velocity achieved; and (5) COD performance: a measurement of the time to complete a non-reactive change of direction or

reactive task. Training status was categorised based on the mean S&C training experience as untrained (<1 year), recreationally trained (1–5 years), or highly trained (>5 years). If the mean S&C training experience was not stated, the minimum required experience was used for categorisation. Sex of the groups was categorised as male-only, female-only or mixed sex. Criteria used to quantify overall intensity and volume for each training mode can be found in Table 1. In brief, overall intensity was categorised specific to each domain and considered both intensity of effort and mechanical factors. In addition, intensity of load was also quantified based on the mean percentage of one repetition maximum (% 1RM) used in the resistance training. In cases where % 1RM was not explicitly stated, % 1RM values were estimated based on the number of repetitions performed per set using methods outlined by Haff and Triplett [20] (OSM 2). Training frequency was classified as the average number of sessions per week throughout the intervention.

Table 1 Criteria used to categorise overall intensity and volume of each training mode included

Training mode	Intensity categorisation	Volume categorisation
Resistance training	Coding based on % 1RM 1 = Low 0–59.9% 1RM 2 = Moderate 60–84.9% 1RM 3 = High \geq 85% 1RM In cases where % 1RM was not explicitly stated, % 1RM value and category were estimated based on the number of repetitions performed per set using methods outlined by Haff and Triplett [19] (Online Supplementary Material 2)	Average number of repetitions performed per set in key exercises 1: Low – 1–5 2: Mid – 6–10 3: High – 11 +
Plyometric	Based on the exercises included, for example: 1 = Low – low amplitude hopping, box jumps, squat jumps 2 = Moderate – bounding, lateral jumps, hurdle jumps, countermovement jump, drop jump with < 30 cm drop 3 = High – drop jump with > 30 cm drop, multidirectional bounding, single leg jumps, rebounding jumps	Average number of foot contacts per session 1: Low – < 80 2: Mid – 80–120 3: High – 120 +
Ballistic	Always categorised as high intensity due to high levels of relative effort required, unless it was explicitly stated sub-maximal effort was used	Average number of repetitions performed per set 1: Low – 1–3 2: Mid – 4–6 3: High – 7 +
Sprint	Always categorised as high intensity due to high levels of relative effort required, unless it was explicitly stated sub-maximal effort was used (e.g. skipping, marching sub-maximal runs)	Average number of runs per session 1: Low – 1–4 2: Mid – 5–9 3: High 10 +
Change of direction	Based on exercises included: 1 = Low – ladder drills, footwork drills, single turn run with < 90° change of direction (COD), 2 = Moderate – lateral movement drills, single turn with > 90° COD, 3 = High – multiple sharp CODs, 505 drills, reactive drills	Average number of runs per session 1: Low – 1–5 2: Mid – 6–9 3: High 10 +
Combined	Combinations of resistance, sprint, ballistic, plyometric and agility training. To be considered a combined training mode, the secondary mode must account for at least 30% of total lower body training volume Categorisation of 1 = Low, 2 = Mid, 3 = High, based on categorisation of included training types	Categorisation of 1 = Low 2 = Mid 3 = High based on categorisation of included training types

COD change of direction, cm centimetre, % 1RM percentage one-repetition maximum

2.4 Statistical Analysis

Effect sizes and their sampling variance were calculated using group mean and standard deviation values calculated pre-intervention and at any subsequent time-point. SMD_{pre} was calculated by dividing the relevant mean difference by the pre-intervention standard deviation. The sampling variance $\sigma_e^2(SMD_{pre})$ of the effect size [21] was calculated using the following formula:

$$\sigma_e^2(SMD_{pre}) = \frac{n-1}{n(n-3)} \left(2(1-r) + nSMD_{pre}^2 \right) - \frac{SMD_{pre}^2}{c(n-1)^2}$$

where n is the sample size, r is the correlation between repeated measures, and $c(n-1)$ is the bias function, which was approximated by $1 - \frac{3}{4n-5}$ [22]. To account for the small sample sizes generally used in S&C, a bias correction was applied to the effect size and sampling variance by multiplying by the approximated bias function and its square, respectively.

All meta-analyses were conducted using a nested four-level mixed-effects meta-analytic model. The series of nestings and a full overview of the model framework are presented in the Online Supplementary files (OSM 3). Predictors were added at level 2 (time of measurement from baseline), level 3 (outcome domain as a categorical predictor (strength, power, sprint, vertical jump, COD)) and level 4 (number of repetitions per set as a categorical predictor (low < 8, high \geq 8); number of repetitions per set as a smooth predictor; number of sets; number of sets as a smooth predictor; number of sets as a categorical predictor (low < 4, high \geq 4); number of sessions per week as a categorical predictor (low < 3, high \geq 3); number of sessions per week as a smooth predictor; number of exercises as a categorical predictor (low < 4, high \geq 4); number of exercises as a smooth predictor; volume as a categorical predictor (low < 2, mid = 2, high > 2); overall intensity as a categorical predictor (low < 2, mid = 2, high > 2); intensity of load (% 1RM) as a categorical predictor (low < 80; high \geq 80); intensity of load (% 1RM) as a smooth predictor; sex as a categorical variable (males, females, mixed); and training status as a categorical predictor (untrained, recreationally trained, highly trained)).

Candidate models were fitted and compared based on predictive accuracy using ELPD-LOO (see OSM for further details) [23]. Median values and 95% credible intervals (CrIs) were presented for regression coefficients where predictors were found to improve the previous model, with the marginal effect of smooth terms visualised using plots and illustrating uncertainties. All analyses were conducted in R [24], with models fit using the brms package interfaced with Stan [25] to perform sampling, and leave-one-out cross-validation performed using the loo package [26]. Analyses were completed across the entire data set including both

resistance-only and resistance-dominant training interventions, with a sensitivity analysis completed with resistance-only training interventions. Outlier SMD_{pre} values were identified by adjusting the distribution by a Tukey g -and- h distribution and obtaining the 0.0035- and 0.9965-quantiles, with values beyond these points removed prior to further analysis [27]. Convergence of parameter estimates was obtained for all models with Gelman-Rubin R -hat values below 1.1 [28]. No attempts were made to assess certainty in the body of evidence for an outcome.

3 Results

3.1 Descriptions of Data

Data from a total of 295 studies comprising 535 groups and 6,710 participants were obtained (Fig. 1 and reference list provided in OSM 4). Of the 535 groups, 372 comprised resistance training only ($n=4,664$), and 163 comprised resistance training combined with other training modalities ($n=2,046$). Sixty-five percent of groups were categorised as male-only, 23% were categorised as mixed sex, and 12% were categorised as female-only. Fifty-four percent of groups were categorised as untrained, 41% were categorised as recreationally trained, and 5% were categorised as highly trained. The duration of interventions ranged from 4 to 208 weeks, with 97% of the data obtained from interventions ≤ 26 weeks. Analyses were thus restricted to time points ≤ 26 weeks following baseline, which provided data from 3,065 outcomes (maximum strength: 1,546 (50%); power: 550 (18%); jump performance: 512 (17%); sprint performance: 370 (12%); COD performance 87 (3%)). Results presented in text are from the complete data set comprising both resistance-only and resistance-dominant training interventions. Sensitivity analyses conducted with resistance-only training interventions were consistent with the larger data set. Full details of the best predictor models at each level for both resistance-only and resistance-dominant training interventions are presented in Tables 2 and 3.

3.2 Null Model

A total of 61 outliers were removed from the analysis such that effect sizes ranged from -0.83 to 4.7 . For the null model the pooled mean effect size was $SMD_{0.5} = 0.55$ [95% CrI 0.51–0.60], with between-study standard deviation $\tau_{0.5} = 0.33$ [75% CrI 0.31–0.36] (Table 2).

3.3 Level 2 Predictors

The only level 2 predictor included was time of measurement following baseline. The median time of measurement




Fig. 1 Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram for included studies. S&C strength and conditioning

was 8 weeks (interquartile range (IQR): 6–12), with binary (short: ≤ 8 weeks, long: > 8 weeks) and linear predictors investigated. Binary categorisation showed an increase in mean pooled effect size with longer interventions ($SMD_{\text{short:long},0.5} = 0.17$ [95% CrI 0.11–0.23]), and for the continuous linear predictor the weekly increase was estimated as $SMD_{\beta_{\text{Time},0.5}} = 0.03$ [95% CrI 0.02–0.04]. The ELPD-LOO difference comparing the null model and inclusion of categorical time or continuous time showed improved model performance and was equal to -11.9 (se: 5.2) and -12.1 (se: 4.2), respectively. All subsequent models including those assessed as part of the sensitivity analysis featured time as a linear predictor (Tables 2 and 3).

3.4 Level 3 Predictors

The only level 3 predictor assessed was the outcome domain measured (Fig. 2). Using maximum strength as the reference level, the mean pooled value was higher in this domain compared to all others ($SMD_{\text{strength:COD},0.5} = -0.28$ [95% CrI -0.39 to -0.17]; $SMD_{\text{strength:jump},0.5} = -0.26$ [95% CrI -0.32 to -0.21]; $SMD_{\text{strength:power},0.5} = -0.28$ [95% CrI -0.33 to -0.23]; $SMD_{\text{strength:sprint},0.5} = -0.40$ [95% CrI -0.47 to -0.34]). The effect of time remained positive after including outcome domain ($SMD_{\beta_{\text{Time},0.5}} = 0.03$ [95% CrI 0.02–0.03]), with a large improvement in model performance (ELPD-LOO difference: -46.8 (se: 12.0)). All subsequent models including those assessed as part of the sensitivity analysis

Table 2 Best performing models at each stage of analysis for complete data set comprising both resistance-only and resistance-dominant training interventions

Level	Included data	Additional included variables [95% CrI]	ELPD-LOO [Standard error]	ELPD-LOO difference	Level 2 standard deviation [75% CrI]	Level 3 standard deviation [75% CrI]	Level 4 standard deviation [75% CrI]
1	291 studies 3004 outcomes	Null model; Mean: 0.55 [0.51–0.60]	– 1805 [60.9]		0.33 [0.31–0.36]	0.25 [0.24–0.26]	0.05 [0.02–0.07]
2	291 studies 3004 outcomes	Time (Weeks): 0.03 [0.02–0.04]	– 1792 [61.2]	– 12.1 [4.2]	0.33 [0.32–0.35]	0.24 [0.22–0.25]	0.05 [0.02–0.08]
2+3	291 studies 3004 outcomes	Outcome domain: Strength: 0.79 [0.74–0.84] Jump: 0.53 [0.47–0.59] Power: 0.52 [0.45–0.58] COD: 0.51 [0.39–0.63] Sprint: 0.39 [0.32–0.46]	– 1728 [62.0]	– 46.8 [12.0]	0.33 [0.31–0.36]	0.20 [0.19–0.22]	0.05 [0.02–0.08]
2+3+4	265 studies 2795 outcomes	Average number sets 0.05 [0.03–0.07] Smooth(Intensity value % IRM)	– 1600 [59.9]	– 12.8 [3.8]	0.33 [0.31–0.36]	0.21 [0.20–0.22]	0.04 [0.01–0.07]
2+3*4	265 studies 2795 outcomes	Interaction between Outcome domain and Smooth(Intensity value % IRM)	– 1588 [59.8]	– 11.7 [4.8]	0.33 [0.31–0.36]	0.20 [0.19–0.22]	0.05 [0.02–0.08]

COD change of direction, *CrI* credible interval

Table shows the additional variables included at each stage of the sequential process along with posterior estimates of model parameters, except for variables where smooth terms were added. At each stage, models also include variables identified as providing the best performance at the preceding stage

featured time as a continuous predictor and outcome domain (Table 2 and 3).

3.5 Level 4 Predictors

Initial analyses were conducted using the categorical volume and overall intensity predictors. No improvement in model performance relative to inclusion of time and outcome domain was obtained for volume (ELPD-LOO difference: + 5.2 (se: 4.6)). In contrast, improvement was observed for overall intensity (ELPD-LOO difference: – 6.6 (se: 2.2)) with evidence of greater mean pooled values for medium ($SMD_{low:medium,0.5} = 0.07$ [95% CrI 0.00–0.15]) and high categories ($SMD_{low:high,0.5} = 0.10$ [95% CrI 0.01–0.19]). The inclusion of smooth terms showed no improvement in model performance when adding the number of sessions per week (ELPD-LOO difference: + 3.4 (se: 2.0)) or the average number of exercises

per session (ELPD-LOO difference: + 1.0 (se: 2.6)). Limited evidence of model improvement was obtained with inclusion of smooth terms for the average number of repetitions per session (ELPD-LOO difference: – 6.4 (se: 3.7)). The marginal effect illustrated a consistent decrease in effect size with a greater number of repetitions that slowed after ten repetitions. Stronger evidence of model improvement using smooth terms was obtained for intensity of load with values expressed as percentage of maximum (ELPD-LOO difference: – 18.0 (se: 4.5)) and the average number of sets per session (ELPD-LOO difference: – 17.5 (se: 4.4)) Marginal smooths illustrated that the relationship was monotonic but non-linear for intensity of load with a reduced incline between 65 and 100% of maximum. The relationship was found to be linear for the average number of sets per session, with an increasing mean effect size association with a greater number of sets ($SMD_{\beta_{Sets,0.5}} = 0.05$ [95% CrI 0.03–0.07]).

Table 3 Best performing models at each stage of analysis for resistance-only training interventions

Level	Included data	Additional Included variables [95% CrI]	ELPD-LOO [Standard error]	ELPD-LOO difference	Level 2 standard deviation [75% CrI]	Level 3 standard deviation [75% CrI]	Level 4 standard deviation [75% CrI]
1	197 studies 1,876 outcomes	Null model; Mean: 0.58 [0.52–0.64]	– 1197 [45.8]		0.36 [0.33–0.39]	0.26 [0.24–0.27]	0.10 [0.08–0.13]
2	197 studies 1,876 outcomes	Time (Weeks): 0.03 [0.02–0.04]	– 1189 [45.7]	– 8.3 [3.3]	0.36 [0.34–0.39]	0.24 [0.22–0.26]	0.11 [0.08–0.13]
2+3	197 studies 1,876 outcomes	Outcome domain: Strength: 0.79 [0.74–0.85] Jump: 0.54 [0.46–0.62] Power: 0.50 [0.42–0.58] COD: 0.58 [0.42–0.70] Sprint: 0.45 [0.36–0.55]	– 1162 [62.0]	– 18.4 [6.5]	0.34 [0.32–0.38]	0.22 [0.20–0.23]	0.10 [0.08–0.13]
2+3+4	177 studies 1,714 outcomes	Average number sets: 0.07 [0.04–0.09] Smooth(Intensity value %1RM)	– 1077 [41.6]	– 8.5 [3.1]	0.36 [0.33–0.40]	0.22 [0.20–0.24]	0.11 [0.09–0.13]
2+3*4	177 studies 1,714 outcomes	Interaction between Outcome domain and Smooth(Intensity value %1RM)	– 1061 [41.8]	– 12.0 [4.2]	0.35 [0.31–0.39]	0.23 [0.20–0.25]	0.10 [0.08–0.13]

CrI credible interval, ELPD-LOO expected log pointwise predictive density for a new dataset that was estimated with leave-one-out cross validation, COD change of direction

Table shows the additional variables included at each stage of the sequential process along with posterior estimates of model parameters, except for variables where smooth terms were added. At each stage, models also include variables identified as providing the best performance at the preceding stage

No improvement in model performance was obtained for the addition of sex (ELPD-LOO difference: +2.9 (se: 2.3)) or training status (ELPD-LOO difference: +0.2 (se: 2.3)). The same lack of improvement in model performance was obtained with sensitivity analyses conducted with resistance-only interventions. A final best performing model including both the average number of sets per session ($SMD_{\beta_{Sets,0.5}} = 0.05$ [95% CrI 0.03–0.07]) and smoothed intensity of load was obtained for the complete data set and for resistance-only training interventions (Table 2 and 3).

3.6 Cross-Level Interactions Between Levels 3 and 4

Potential cross-level interactions were investigated separately between outcome domain and both average number sets per session and intensity of load. No improvement in model performance was obtained for the cross-level interaction between outcome domain and average number of sets per session (ELPD-LOO difference: – 0.6 (se: 3.5)). In

contrast, model performance was increased for the cross-level interaction between outcome domain and intensity of load (ELPD-LOO difference: – 0.6 (se: 3.5)), with spline modelled intensity showing markedly different relationships across the different outcome domains (Table 2 and Fig. 3).

4 Discussion

The primary aim of this meta-analysis was to produce a comprehensive modelling of the dose–response relationships between resistance exercise and commonly used measures of physical performance within S&C. The analyses identified that a range of factors are associated with the magnitude of change across resistance-only and resistance-dominant training interventions. These factors include the length of the intervention, outcome type, volume, overall intensity of training, and the intensity of load. The analyses also




Fig. 2 (Top): Posterior distributions of pooled mean effect sizes across outcome domains. (Bottom): Values represent shrunken values after fitting meta-analytic model also accounting for time of measurement. Black lines represent 75% and 95% credible intervals

identified important interactions between loading intensity and outcome domain, such that some outcomes are more likely to experience greater improvements with a range of sub-maximum loads (30–70% 1RM).

Length of intervention has generally not been explored in dose–response modelling due to most reviews focusing on a smaller number of homogeneous studies with a restricted range of durations. Across the studies included in this meta-analysis, durations were found to be relatively short, with the median duration equal to 8 weeks and 97% of interventions lasting less than 26 weeks. The prevalence of shorter duration studies may be due to multiple challenges associated with longer duration interventions including an increased need for resources as well as greater inclusion and agreement from key stakeholders (athlete, organisation and management), increased dropouts, and scheduling difficulties to fit within non-competitive periods [29]. Despite the relatively short intervention durations, the results show that substantive improvements can be made, and that longer durations within these time frames create greater mean improvements. Therefore, practitioners and athletes could use shorter time periods, potentially during off-season or preseason, to maximise physiological improvements when

competition or sporting demands are lowest. The effect of duration remained consistent throughout the model-building process with standardised mean differences estimated to increase by approximately 0.03 for each additional week of training. Given the relatively short and homogeneous durations included, there was limited ability to explore the functional form of changes over longer durations (e.g., playing seasons, years). In a recent large modelling study investigating the time-course of strength adaptations, Steele et al. [30] showed that linear-log growth models were appropriate to describe improvements of relatively untrained participants over the course of almost 7 years, with improvements tending to plateau after approximately 1 year. The training stimulus investigated by Steele et al. [30] focused on minimal dose resistance training (1x/week, single sets to momentary failure of six exercises), which is likely to have influenced the parameters obtained. Our analysis was limited to durations of no more than 26 weeks and thus we cannot draw inferences as to how results might change over longer time frames. Further research is required to better understand the influence of duration of an intervention and likely interactions between participant characteristics, the specific outcome, the training stimulus, and changes in the




Fig. 3 (Top): Marginal effects of smooth terms illustrating interactions between intensity of load expressed as percentage of maximum and outcome domain. (Bottom): Density plots at the top of each figure illustrate the distribution of the load intensity variable for the

given outcome domain. Solid lines represent the best estimate of the smooth relationship and shaded regions represent intervals of uncertainty (75% and 95%)

training stimulus following, for example, different periodized approaches.

The manipulation of acute program variables within resistance training interventions is often focused on the development of a single outcome domain. Previous meta-analyses investigating dose–response relationships have

predominantly focused on the development of muscular strength and/or hypertrophy [7–17]. The current meta-analysis demonstrated varying effects across multiple outcome domains commonly targeted with the greatest effect sizes obtained for strength, and the lowest obtained for sprint performance. Resistance training for the purpose of

improving maximum strength is arguably the most investigated and well-understood area within S&C [1, 2], and greater effect sizes may reflect this increased refinement and specificity between traditional resistance-training methods and maximum strength outcomes [1]. In addition, researchers frequently test maximum strength using the same exercises included in the training intervention, further increasing specificity and potentially the improvements measured [31]. Results from a previous meta-analysis indicate that the dose–response relationship between the % 1RM and strength gains diminishes when testing is carried out isometrically [15]. Practitioners should be advised to select an appropriate testing mode for the given training stimulus applied, whilst being aware of any potential upward or downward shift in expected results dependent on the measurement used. Further study is needed to provide greater context to the transfer of strength from varied magnitudes of load to neutral testing modalities.

Following maximum strength, jump performance and power generated the next largest effects. Similar magnitude improvements in jump performance and power may be expected given the well-established relationship between the two factors [32–35]. In addition, many of the studies measured power during loaded and unloaded jumps, further increasing associations and similar magnitude improvements. Outcomes relating to sprint performance demonstrated the lowest magnitude improvements. Sprinting comprises a substantial and complex technique element [36–40], and given the relatively low number of studies (~12%) that included sprint specific interventions, a lower effect size distribution may be expected. More broadly, the lower effect size distribution for sprint outcomes may also reflect a lack of specificity with regards to development of relevant physical outputs. Transference between improvements and long-term adaptations in S&C is dependent primarily on the training principles of specificity and progressive overload, respectively [41, 42]. Most training methods included in the meta-analysis focused on bilateral production of maximum vertical forces over long durations. In contrast, sprinting activities require high forces produced over short ground contact times that are predominantly unilateral with substantive horizontal components in relation to the body's position relative to the ground [37, 38, 43, 44]. In a recent meta-analysis, Murphy et al. [4] showed moderate to strong relationships between improvements in strength, power and sprint performance in team sport athletes, concluding that greater development of physical capacities may result in further improvements in sprint performance. Despite these correlations, however, researchers have also shown that large increases in maximum strength (~12–18%) translate into only small decreases in sprint times (~–2–8%) [45–47]. With evidence to suggest restrictions may exist in the transference of physical capacities to sprinting, further sprint-specific

training modes such as resisted sprint training may provide additional benefits allowing for the ability to overload kinetic output with increased kinematic specificity to the complex movement of sprinting [48]. Collectively, there appears to be scope for future research to investigate why improvements in sprint performance are generally much smaller than other outcome domains and whether this difference can be ameliorated with a focus on certain training practices.

Movements associated with COD and agility could be considered even more complex than sprinting due to the high acceleration and deceleration demands, the ability to rapidly alter body position, combined with the need in some activities to react to an external stimulus [49, 50]. The results of the current meta-analysis suggest improvements in COD are likely to be of similar magnitude to those measured during vertical jump and tasks focusing on the development of power, albeit with a greater level of uncertainty. Outcomes measuring COD and the related construct of change of direction speed and agility represent a developing area within S&C [51, 52] with only ~3% of outcomes assessing COD performance. Whilst reasons for the larger effect size distribution in comparison to linear sprinting require further study, potential explanations include the complex, multifaceted nature of the tasks and the scope for multiple limiting factors to be addressed. Additionally, it is recognised that many agility and COD tasks include substantive skill elements [52], such that failure to appropriately familiarize participants could lead to systematic biases in regard to learning effects and subsequent overestimations of effect sizes.

The results of the current meta-analysis demonstrate the importance of training intensity. Previous researchers investigating dose–response relationships have tended to contextualise and quantify intensity based on load and thereby % 1RM [7, 8, 14, 15]. This approach works best when considering traditional strength or hypertrophy focused interventions comprising large compound movements where 1RMs can be measured and appropriately summarise a relevant feature of intensity [53]. An aim of the current meta-analysis was to investigate dose–response relationships across a range of resistance-based training modalities and outcomes; therefore, in addition to intensity of load expressed as a % 1RM, a more general categorisation scheme was included. Interventions comprising predominantly ballistic, loaded jumping or sprinting exercises were always considered high intensity, due to the high mechanical loads and assumption that they are conducted with maximal intent, unless stated otherwise. Across all outcomes, evidence was obtained that greater overall intensity was associated with increased effect sizes, with interventions judged to be of medium and high overall intensity expected to increase effect sizes by approximately 0.10 relative to low overall intensity. When prescribing intensity, practitioners should consider training intensity beyond simply intensity of load (% 1RM). For

example, ensuring maximum intent during ballistic exercise exemplifies a simplistic method of prescribing higher intensity training. Incorporating methods such as velocity-based training provides the opportunity for instantaneous feedback during ballistic movements to encourage maximal intent [54]. Adjusting the intensity of plyometric training may be more complex and largely dependent on exercise selection to increase or decrease take-off and landing ground reaction forces [55].

Additional detailed information on the dose–response relationship of intensity was obtained when investigating potential interaction effects between outcome and intensity of load measured by % 1RM. The results identified a range of different profiles, with no clear pattern for COD, monotonic increases for strength and speed, a monotonic decrease for jump performance, and a parabolic profile for power. The best estimate profile for maximum strength appeared non-linear with an inflection point ~70% 1RM, where further increases in effect size estimates started to slow with additional load. The results of the present meta-analysis are consistent with previous reviews, e.g. Peterson et al. [8] identified increased effects with heavier % 1RM loads but diminishing effects, particularly with untrained participants [18]. The results also align with previous research indicating that heavy load training may increase muscle activation by up to 30% [56], conceivably providing a stronger stimulus for adaptation. Some authors have also suggested, however, that improvements in strength with greater loads may be inflated due to high specificity of task and outcome, given that strength testing is often conducted performing the 1RM of the movement being trained [15].

Analysis of sprint performance also identified a monotonic increase in effect, with the greatest increases obtained with the heaviest loads. The most common sprint outcomes investigated in S&C research include the time to sprint between 5 and 50 m, with the most frequent intervals comprising 10, 20 and 30 m [4]. Most studies have been conducted with either team sport or untrained participants who achieve maximum velocity between 15 and 40 m, in comparison to trained sprinters who require distances of 40–80 m to achieve maximum velocity [38, 44, 57, 58]. Consequently, sprint data collected over 10–30 m may provide researchers with divergent outcomes describing both acceleration and maximum velocity. Previous studies have reported strong associations with outcomes designed to assess acceleration (e.g., 10 m) and horizontal force, power and relative strength with longer duration ground contact times (approximately 200 ms) [37, 38]. In contrast, maximum velocity sprinting has been shown to be dependent on the ability to maintain large horizontal and vertical forces whilst minimizing braking forces with reduced ground contact time (approximately 100 ms) [38, 39, 58]. Previous meta-analyses have concluded that high-intensity non-specific resistance exercise

is among the most effective training methods to improve sprint performance in team sport athletes [4, 36, 59]. Neural and morphological adaptations associated with high-load resistance exercise and improved force output may provide a mechanism for positive transfer to the high levels of horizontal force required during early-phase acceleration to improve sprint performance. Highly trained individuals may require more specific training methods, however, that target improvements in physical qualities while matching the kinematic demands of sprinting [1, 4].

In contrast to the increasing dose–response relationships with intensity for strength and sprint performance, results identified monotonic decreases for jump performance with the largest effects obtained at ~30% 1RM. Jump performance is dependent on net impulse and take-off velocity such that % 1RM loads lifted with maximum intent provide sufficient stimulus but do not limit velocity to a large extent may provide the greatest transfer to improvements in jump performance [33, 60]. Previous researchers have also demonstrated that jump squat training with low (<30% 1RM) or no additional load can produce velocity-specific adaptations associated with improvements in jump performance [61]. The intensity profile for outcomes measuring power production was parabolic, with the greatest improvements obtained between ~40 and 70% 1RM. These results support the hypothesis that performing resistance exercise with loads that elicit the largest power outputs is the most effective method to improve power and that for most exercises power is maximised between 30 and 70% 1RM [62]. During weightlifting exercises (clean, snatch, hang and pull variations), power is maximised with heavier loads ($\geq 70\%$ 1RM), whereas loads of 0–30% 1RM maximise power during jump squat exercises [62]. As the optimum load for power production is exercise-dependent, practitioners should be aware of the appropriate load required to stimulate peak power output within the prescribed exercises and endeavour to create athlete-specific profiles where access to relevant measurement devices is available.

The influence of training frequency and volume on strength and hypertrophy has been assessed in several previous meta-analyses [10–12, 16]. The results obtained herein were mixed but showed limited evidence that these factors were influential. No improvement in model performance was obtained when including the number of sessions per week as a categorical or continuous predictor, or the average number of exercises per session. Only limited evidence of model improvement was obtained for the average number of repetitions per set, with the marginal effect showing declines as the number of repetitions increased. In contrast, evidence was obtained for greater effect sizes with increasing number of sets per session. Seminal research by Rhea et al. [9] was among the first, within S&C, to use meta-analytical techniques to assess the use of single versus multiple

sets in resistance exercise for strength development. The authors concluded multiple sets were more beneficial than single-set training. A follow-up meta-regression by Krieger et al. [63] found a 46% increase in muscular strength when completing two to three sets, in comparison to single sets, although no further difference was found for resistance exercise with more than four sets. More recently, researchers have concluded that increasing weekly training volume through increased number of sets can produce similar results to increasing training frequency [11, 12]. The current meta-analysis demonstrates that focussing on a smaller number of key exercises while completing multiple sets at an appropriate intensity for a targeted outcome may be more beneficial than attempting to perform many exercises with an increased frequency.

The training status of participants is a key consideration when designing and implementing resistance exercise. Previous meta-analyses have demonstrated rank-order effects, with the largest improvements obtained by untrained participants [1, 7], followed by recreationally trained and then highly trained participants. In contrast, the current meta-analysis found a lack of evidence to support different effects across the training status categories. Differences in results obtained in the present versus previous meta-analyses may be due to several reasons. Previous analyses have been less formal than those conducted herein, with authors identifying differences based primarily on point estimates. In contrast, participant training status was assessed in the present analysis with predictor variables for lower levels already included in the model and addition of the factor was assessed based on ability to improve model performance. The lack of data for highly trained participants, disproportionate inclusion of untrained participants, combined with short-duration (≤ 26 weeks) interventions, which are known limitations within S&C research [64], may have also influenced the results obtained and discordance with what is generally believed in the field. With advancements in technology and ability to collect valid and reliable high-frequency data over longer periods across all levels of sport and recreation, use of longitudinal data collected in the field provides opportunities to better investigate differences in dose-responses relative to participant training status.

In addition to training status, the current meta-analysis found a lack of evidence to support different effects between sexes. Although males exhibit greater levels of baseline strength and muscle mass [65], the current meta-analysis results are consistent with previous research that has failed to identify any difference in effects between sexes in improvements in strength or hypertrophy, with indications that varying levels of adaptations may be more related to relative strength [10, 65–67]. A previous meta-analysis conducted by de Villareal et al. [68], however, reported greater improvement in males following plyometric training

relative to females. The authors were unable to provide a strong rationale for the finding, and speculated that large differences in sample sizes between the sexes may have confounded results. Research has shown that stronger individuals are able to produce a greater rate of force development and power during time-restricted tasks [3], and so there is the potential that increased strength at baseline may be advantageous for plyometric training. Further research is required to identify potential differences in the dose–response relationship between sexes and more complex sport-specific outcomes. Female participants are largely under-represented in S&C research, with only 12% of the studies included here conducted with female-only groups. In addition, research suggests that only 39% of all published sport science data are collected with female participants [69]. To better address the question of potential differences in dose–response relationships, more female participant data are required.

Whilst this is the most comprehensive meta-analysis to date investigating dose–response relationships between resistance and resistance-dominant interventions, there are multiple limitations that should be considered when interpreting the findings. There are clear limitations in summarising different dose components from a training intervention based on, for example, the average number of sets, where this variable may change substantially depending on the periodisation or progression model. In addition, for variables such as overall training volume and intensity, there was a high degree of subjectivity and challenge in obtaining a single value, particularly when considering different training modes. The current meta-analysis was intended to uncover general relationships, but the extensive heterogeneity across the data set limits nuance and there are limitations in drawing strong inferences from pooling of indirect data. It is expected that there will be many instances where factors such as training frequency and volume strongly influence the effectiveness of an intervention; however, variables quantifying frequency and volume in the present analysis lacked predictive power across this large and heterogeneous data set. Despite these results and considering the limitations of the meta-analysis, practitioners are still recommended to implement periodized training interventions that include appropriate manipulations in intensity, volume and frequency over time in an attempt to maximise a given outcome. Overall, the results of the present meta-analysis suggest that practitioners should focus first on overall intensity and intensity of load with the appropriate target outcomes in mind.

5 Conclusion

The current meta-analysis is the most comprehensive to date to investigate dose–response relationships of resistance and resistance-dominant training with respect to a range

of commonly studied outcome domains in S&C research. The findings are that resistance exercise is effective over relatively short durations (~8 weeks) and extending a single intervention over longer periods is likely to result in further improvements. The expected magnitude of improvement appears to be predominantly influenced by intensity and the outcome domain measured. Performance of resistance training with a higher intensity as measured by a composite of the effort applied, the difficulty of the exercise, and maximising target biomechanical quantities result in greater improvements. When considering the magnitude of the load lifted as a % 1RM, the profile that creates the greatest improvements is dependent on the outcome domain. Improvements in strength are likely to be maximised with the heaviest loads, whereas vertical jump performance may be maximised with relatively light loads (~30% 1RM), and power with low to moderate loads (40–70% 1RM). Sprinting performance represents the most difficult outcome domain to improve with resistance and resistance-dominant training, and this may be influenced by lower specificity.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s40279-024-02006-3>.

Declarations

Conflicts of Interest/Competing Interests BJS serves on the scientific advisory board of Tonal Corporation, a manufacturer of fitness equipment. The other authors report no potential conflicts of interest.

Funding No funding was received to assist with the preparation of this article.

Author Contributions PS conceived the idea for this review. AM and PS designed the search protocol as part of a larger study. AM conducted the searches, screening and selected the studies for inclusion, with input from PS. PS designed and undertook the statistical analysis. AM and PS wrote the first draft of the manuscript with critical input from BS. All authors read and approved the final version.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Swinton PA, Burgess K, Hall A, Greig L, Psyllas J, Aspe R, Maughan P, Murphy A. Interpreting magnitude of change in strength and conditioning: Effect size selection, threshold values and Bayesian updating. *J Sports Sci.* 2022;40(18):2047–54. <https://doi.org/10.1080/02640414.2022.2128548>.
- Kraemer WJ, Ratamess NA, Flanagan SD, Shurley JP, Todd JS, Todd TC. Understanding the science of resistance training: An evolutionary perspective. *Sports Med.* 2017;47(12):2415–35. <https://doi.org/10.1007/s40279-017-0779-y>.
- Suchomel TJ, Nimphius S, Bellon CR, Stone MH. The importance of muscular strength: training considerations. *Sports Med.* 2018;48(4):765–85. <https://doi.org/10.1007/s40279-018-0862-z>.
- Murphy, A, Burgess, K, Hall, AJ, Aspe, RR, Swinton, PA. 2022. The effects of strength and conditioning interventions on sprinting performance in team sport athletes: a systematic review and meta-analysis. *J Strength Cond Res* [online], (accepted).
- Swinton, PA, Burges, K, Hall, A, Greig L, Psyllas J, Aspe R, Maughan P, Murphy A. 2021 A Bayesian approach to interpret intervention effectiveness in strength and conditioning: Part 1. A meta-analysis to derive context-specific thresholds. Pre-print available from SportRxiv. <https://doi.org/10.51224/SRXIV.9>.
- Bird SP, Tarpenning KM, Marino FE. Designing resistance training programmes to enhance muscular fitness. *Sports Med.* 2005;35(10):841–51. <https://doi.org/10.2165/00007256-200535100-00002>.
- Rhea MR, Alvar BA, Burkett LN, Ball SD. A meta-analysis to determine the dose response for strength development. *Med Sci Sports Exerc.* 2003;35(3):456–64. <https://doi.org/10.1249/01.MSS.0000053727.63505.D4>.
- Peterson MD, Rhea MR, Alvar BA. Maximizing strength development in athletes: a meta-analysis to determine the dose-response relationship. *J Strength Cond Res.* 2004;18(2):377–82. <https://doi.org/10.1519/R-12842.1>.
- Rhea MR, Alvar BA, Burkett LN. Single versus multiple sets for strength: a meta-analysis to address the controversy. *Res Q Exerc Sport.* 2002;73(4):485–8. <https://doi.org/10.1080/02701367.2002.10609050>.
- Schoenfeld BJ, Ogborn D, Krieger JW. Dose-response relationship between weekly resistance training volume and increases in muscle mass: a systematic review and meta-analysis. *J Sports Sci.* 2017;35(11):1073–82. <https://doi.org/10.1080/02640414.2016.1210197>.
- Ralston GW, Kilgore L, Wyatt FB, Baker JS. The effect of weekly set volume on strength gain: a meta-analysis. *Sports Med.* 2017;47(12):2585–601. <https://doi.org/10.1007/s40279-017-0762-7>.
- Grgic J, Schoenfeld BJ, Davies TB, Lazinica B, Krieger JW, Pedisic Z. Effect of resistance training frequency on gains in muscular strength: a systematic review and meta-analysis. *Sports Med.* 2018;48(5):1207–20. <https://doi.org/10.1007/s40279-018-0872-x>.
- Ralston GW, Kilgore L, Wyatt FB, Buchan D, Baker JS. Weekly training frequency effects on strength gain: a meta-analysis. *Sports Med Open.* 2018;4(1):1–24. <https://doi.org/10.1186/s40798-018-0149-9>.
- Schoenfeld BJ, Wilson JM, Lowery RP, Krieger JW. Muscular adaptations in low-versus high-load resistance training: a meta-analysis. *Eur J Sport Sci.* 2016;16(1):1. <https://doi.org/10.1080/17461391.2014.989922>.
- Schoenfeld BJ, Grgic J, Ogborn D, Krieger JW. Strength and hypertrophy adaptations between low-vs. high-load resistance training: a systematic review and meta-analysis. *J Strength Cond Res.* 2017;31(12):3508–23. <https://doi.org/10.1519/JSC.0000000000002200>.
- Schoenfeld BJ, Ogborn D, Krieger JW. Effects of resistance training frequency on measures of muscle hypertrophy: a systematic review and meta-analysis. *Sports Med.* 2016;46(11):1689–97. <https://doi.org/10.1007/s40279-016-0543-8>.
- Cuthbert M, Haff GG, Arent SM, Ripley N, McMahon JJ, Evans M, Comfort P. Effects of variations in resistance training

- frequency on strength development in well-trained populations and implications for in-season athlete training: a systematic review and meta-analysis. *Sports Med.* 2021;51(9):1967–82. <https://doi.org/10.1007/s40279-021-01460-7>.
18. Peterson MD, Rhea MR, Alvar BA. Applications of the dose-response for muscular strength development: a review of meta-analytic efficacy and reliability for designing training prescription. *J Strength Cond Res.* 2005;19(4):950–8. <https://doi.org/10.1519/R-16874.1>.
 19. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ.* 2021;372: n71. <https://doi.org/10.1136/bmj.n71>.
 20. Haff GG, Triplett NT, editors. *Essentials of strength training and conditioning.* 4th ed. Human Kinetics; 2015.
 21. Morris SB, DeShon RP. Combining effect size estimates in meta-analysis with repeated measures and independent-groups designs. *Psychol Methods.* 2002;7(1):105–25. <https://doi.org/10.1037/1082-989x.7.1.105>.
 22. Hedges LV. Estimation of effect size from a series of independent experiments. *Psychol Bull.* 1982;92(2):490–9. <https://doi.org/10.1037/0033-2909.92.2.490>.
 23. Vehtari A, Gelman A, Gabry J. Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Stat Comput.* 2017;27:1413–32. <https://doi.org/10.1007/s11222-016-9696-4>.
 24. R Core Team. R: A language and environment for statistical computing. 2021; R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>. Accessed 1 Aug 2022.
 25. Bürkner PC. brms: An R package for Bayesian multilevel models using Stan. *J Stat Softw.* 2017;80(1):1–28. <https://doi.org/10.18637/jss.v080.i01>.
 26. Vehtari A, Gabry J, Magnusson M, Yao Y, Bürkner P, Paananen T, Gelman A. “loo: Efficient leave-one-out cross-validation and WAIC for Bayesian models.” 2023; R package version 2.6.0.
 27. Verardi V, Vermandele C. Univariate and multivariate outlier identification for skewed or heavy-tailed distributions. *Stata J.* 2018;18(3):517–32. <https://doi.org/10.1177/1536867X1801800303>.
 28. Gelman A, Carlin JB, Stern HS, Rubin DB. *Bayesian data analysis.* Taylor & Francis; 2014.
 29. Short S, Tuttle M. The gap between research and clinical practice for injury prevention in elite sport: a clinical commentary. *Int J Sports Phys Ther.* 2020;15(6):1229–34. <https://doi.org/10.26603/ijsp20201229>.
 30. Steele J, Fisher JP, Giessing J, Androulakis-Korakakis P, Wolf M, Kroeske B, Reuters R. Long-term time-course of strength adaptation to minimal dose resistance training through retrospective longitudinal growth modeling. *Res Q Exerc Sport.* 2022. <https://doi.org/10.1080/02701367.2022.2070592>.
 31. Schoenfeld BJ, Grgic J, Van Every DW, Plotkin DL. Loading recommendations for muscle strength, hypertrophy, and local endurance: a re-examination of the repetition continuum. *Sports.* 2021;9(2):32. <https://doi.org/10.3390/sports9020032>.
 32. Swinton PA, Lloyd R, Keogh JW, Agouris I, Stewart AD. Regression models of sprint, vertical jump, and change of direction performance. *J Strength Cond Res.* 2014;28(7):1839–48. <https://doi.org/10.1519/JSC.0000000000000348>.
 33. Nuzzo JL, McBride JM, Cormie P, McCaulley GO. Relationship between countermovement jump performance and multijoint isometric and dynamic tests of strength. *J Strength Cond Res.* 2008;22(3):699–707. <https://doi.org/10.1519/JSC.0b013e31816d5eda>.
 34. Owen NJ, Watkins J, Kilduff LP, Bevan HR, Bennett MA. Development of a criterion method to determine peak mechanical power output in a countermovement jump. *J Strength Cond Res.* 2014;28(6):1552–8. <https://doi.org/10.1519/JSC.00000000000000311>.
 35. Barker LA, Harry JR, Mercer JA. Relationships between countermovement jump ground reaction forces and jump height, reactive strength index, and jump time. *J Strength Cond Res.* 2018;32(1):248–54. <https://doi.org/10.1519/JSC.00000000000002160>.
 36. Nicholson B, Dinsdale A, Jones B, Till K. The training of short distance sprint performance in football code athletes: a systematic review and meta-analysis. *Sports Med.* 2020;51:1179–207. <https://doi.org/10.1007/s40279-020-01372-y>.
 37. Morin JB, Edouard P, Samozino P. Technical ability of force application as a determinant factor of sprint performance. *Med Sci Sports Exerc.* 2011;43(9):1680–8. <https://doi.org/10.1249/MSS.0b013e318216ea37>.
 38. Morin JB, Bourdin M, Edouard P, Peyrot N, Samozino P, Lacour JR. Mechanical determinants of 100-m sprint running performance. *Eur J Appl Physiol.* 2012;112(11):3921–30. <https://doi.org/10.1249/MSS.0b013e318216ea37>.
 39. Colyer SL, Nagahara R, Takai Y, Salo AI. How sprinters accelerate beyond the velocity plateau of soccer players: Waveform analysis of ground reaction forces. *Scand J Med Sci Sports.* 2018;28(12):2527–35. <https://doi.org/10.1111/sms.13302>.
 40. Mann R, Murphy A. *The mechanics of sprinting and hurdling.* Long Beach: Ralph V. Mann; 2018.
 41. Suarez DG, Wagle JP, Cunanan AJ, Sausaman RW, Stone MH. Dynamic correspondence of resistance training to sport: a brief review. *Strength Cond J.* 2019;41(4):80–8. <https://doi.org/10.1519/SSC.0000000000000458>.
 42. Issurin VB. Training transfer: scientific background and insights for practical application. *Sports Med.* 2013;43(8):675–94. <https://doi.org/10.1007/s40279-013-0049-6>.
 43. Fitzpatrick DA, Cimadoro G, Cleather DJ. The magical horizontal force muscle? a preliminary study examining the “Force-Vector” theory. *Sports.* 2019;7(2):30. <https://doi.org/10.3390/sports702030>.
 44. Morin JB, Slawinski J, Dorel S, Couturier A, Samozino P, Brughelli M, Rabita G. Acceleration capability in elite sprinters and ground impulse: push more, brake less? *J Biomech.* 2015;48(12):3149–54. <https://doi.org/10.1016/j.jbiomech.2015.07.009>.
 45. Cronin J, Ogden T, Lawton T, Brughelli M. Does increasing maximal strength improve sprint running performance? *Strength Cond J.* 2007;29(3):86–95. [https://doi.org/10.1519/1533-4295\(2007\)29\[86:DIMSIS\]2.0.CO;2](https://doi.org/10.1519/1533-4295(2007)29[86:DIMSIS]2.0.CO;2).
 46. Harris NK, Cronin JB, Hopkins WG, Hansen KT. Squat jump training at maximal power loads vs. heavy loads: effect on sprint ability. *J Strength Cond Res.* 2008;22(6):1742–9. <https://doi.org/10.1519/JSC.0b013e318187458a>.
 47. Comfort P, Haigh A, Matthews MJ. Are changes in maximal squat strength during preseason training reflected in changes in sprint performance in rugby league players? *J Strength Cond Res.* 2012;26(3):772–6. <https://doi.org/10.1519/JSC.0b013e31822a5cbf>.
 48. Alcaraz PE, Carlos-Vivas J, Oponjuru BO, Martínez-Rodríguez A. The effectiveness of resisted sled training (RST) for sprint performance: a systematic review and meta-analysis. *Sports Med.* 2018;48(9):2143–65. <https://doi.org/10.1007/s40279-018-0947-8>.
 49. Harper DJ, McBurnie AJ, Santos TD, Eriksrud O, Evans M, Cohen DD, Rhodes D, Carling C, Kiely J. Biomechanical and neuromuscular performance requirements of horizontal deceleration: a review with implications for random intermittent multidirectional sports. *Sports Med.* 2022;52(10):2321–54. <https://doi.org/10.1007/s40279-022-01693-0>.
 50. Brughelli M, Cronin J, Levin G, Chaouachi A. Understanding change of direction ability in sport: a review of resistance training

- studies. *Sports Med.* 2008;38(12):1045–63. <https://doi.org/10.2165/00007256-200838120-00007>.
51. Sheppard JM, Young WB. Agility literature review: Classifications, training and testing. *J Sports Sci.* 2006;24(9):919–32. <https://doi.org/10.1080/02640410500457109>.
 52. Nimphius S, Callaghan SJ, Bezodis NE, Lockie RG. Change of direction and agility tests: challenging our current measures of performance. *Strength Cond J.* 2018;40(1):26–38. <https://doi.org/10.1519/SSC.000000000000309>.
 53. Thompson SW, Rogerson D, Ruddock A, Barnes A. The effectiveness of two methods of prescribing load on maximal strength development: a systematic review. *Sports Med.* 2020;50(5):919–38. <https://doi.org/10.1007/s40279-019-01241-3>.
 54. Nagata A, Doma K, Yamashita D, Hasegawa H, Mori S. The effect of augmented feedback type and frequency on velocity-based training-induced adaptation and retention. *J Strength Cond Res.* 2020;34(11):3110–7. <https://doi.org/10.1519/JSC.0000000000002514>.
 55. Ebben WP, Fauth ML, Garceau LR, Petushek EJ. Kinetic quantification of plyometric exercise intensity. *J Strength Cond Res.* 2011;25(12):3288–98. <https://doi.org/10.1519/JSC.0b013e31821656a3>.
 56. Schoenfeld BJ, Contreras B, Willardson JM, Fontana F, Tiriyaki-Sonmez G. Muscle activation during low- versus high-load resistance training in well-trained men. *Eur J Appl Physiol.* 2014;114:2491–7. <https://doi.org/10.1007/s00421-014-2976-9>.
 57. Rumpf MC, Lockie RG, Cronin JB, Jalilvand F. Effect of different sprint training methods on sprint performance over various distances: a brief review. *J Strength Cond Res.* 2016;30(6):1767–85. <https://doi.org/10.1519/JSC.0000000000001245>.
 58. Haugen T, McGhie D, Ettema G. Sprint running: From fundamental mechanics to practice—a review. *Eur J Appl Physiol.* 2019;119(6):1273–87. <https://doi.org/10.1007/s00421-019-04139-0>.
 59. Nicholson B, Dinsdale A, Jones B, Till K. The training of medium-to long-distance sprint performance in football code athletes: a systematic review and meta-analysis. *Sports Med.* 2022;52:257–86. <https://doi.org/10.1007/s40279-021-01552-4>.
 60. Kirby TJ, McBride JM, Haines TL, Dayne AM. Relative net vertical impulse determines jumping performance. *J Appl Biomech.* 2011;27(3):207–14. <https://doi.org/10.1123/jab.27.3.207>.
 61. Cormie P, McCaulley GO, McBride JM. Power versus strength-power jump squat training: influence on the load-power relationship. *Med Sci Sports Exerc.* 2007;39(6):996–1003. <https://doi.org/10.1097/mss.0b013e3180408e0c>.
 62. Soriano MA, Jiménez-Reyes P, Rhea MR, Marín PJ. The optimal load for maximal power production during lower-body resistance exercises: a meta-analysis. *Sports Med.* 2015;45:1191–205. <https://doi.org/10.1007/s40279-015-0341-8>.
 63. Krieger JW. Single versus multiple sets of resistance exercise: a meta-regression. *J Strength Cond Res.* 2009;23(6):1890–901. <https://doi.org/10.1519/JSC.0b013e3181b370be>.
 64. Stone MH, Collins D, Plisk S, Haff G, Stone ME. Training principles: evaluation of modes and methods of resistance training. *Strength Cond J.* 2000;22(3):65–76. <https://doi.org/10.1080/14763140208522788>.
 65. Hubal MJ, Gordish-Dressman HE, Thompson PD, Price TB, Hoffman EP, Angelopoulos TJ, Gordon PM, Moyna NM, Pescatello LS, Visich PS, Zoeller RF. Variability in muscle size and strength gain after unilateral resistance training. *Med Sci Sports Exerc.* 2005;37(6):964–72. <https://doi.org/10.1249/01.mss.0000170469.90461.5f>.
 66. Nimphius S, McBride JM, Rice PE, Goodman-Capps CL, Capps CR. Comparison of quadriceps and hamstring muscle activity during an isometric squat between strength-matched men and women. *J Sports Sci Med.* 2019;18(1):101–8.
 67. Nimphius S. Exercise and sport science failing by design in understanding female athletes. *Int J Sports Physiol Perform.* 2019;14(9):1157–8. <https://doi.org/10.1123/ijsp.2019-0703>.
 68. de Villarreal ES, Kellis E, Kraemer WJ, Izquierdo M. Determining variables of plyometric training for improving vertical jump height performance: a meta-analysis. *The J Strength Cond Res.* 2009;23(2):495–506. <https://doi.org/10.1519/JSC.0b013e318196b7c6>.
 69. Hagstrom AD, Marshall PW, Halaki M, Hackett DA. The effect of resistance training in women on dynamic strength and muscular hypertrophy: a systematic review with meta-analysis. *Sports Med.* 2020;50(6):1075–93. <https://doi.org/10.1007/s40279-019-01247-x>.

Authors and Affiliations

Paul Alan Swinton¹  · Brad J. Schoenfeld² · Andrew Murphy^{1,3}

✉ Paul Alan Swinton
p.swinton@rgu.ac.uk

¹ School of Health Sciences, Robert Gordon University, Garthdee Road, Aberdeen AB10 7QG, UK

² Department of Exercise Science and Recreation, CUNY Lehman College, Bronx, NY, USA

³ Greater Western Sydney Giants, Sydney, NSW, Australia

Supplementary File 1: Checklist of Preferred Reporting items for Systematic Review and Meta-Analysis

Section and Topic	Item #	Checklist item	Location where item is reported
TITLE			
Title	1	Identify the report as a systematic review.	Reported as a meta-analysis. Search is not exhaustive and does not include all relevant research.
ABSTRACT			
Abstract	2	See the PRISMA 2020 for Abstracts checklist.	
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of existing knowledge.	4-6
Objectives	4	Provide an explicit statement of the objective(s) or question(s) the review addresses.	6,7
METHODS			
Eligibility criteria	5	Specify the inclusion and exclusion criteria for the review and how studies were grouped for the syntheses.	8
Information sources	6	Specify all databases, registers, websites, organisations, reference lists and other sources searched or consulted to identify studies. Specify the date when each source was last searched or consulted.	7,8
Search strategy	7	Present the full search strategies for all databases, registers and websites, including any filters and limits used.	Not included.
Selection process	8	Specify the methods used to decide whether a study met the inclusion criteria of the review, including how many reviewers screened each record and each report retrieved, whether they worked independently, and if applicable, details of automation tools used in the process.	8
Data collection process	9	Specify the methods used to collect data from reports, including how many reviewers collected data from each report, whether they worked independently, any processes for obtaining or confirming data from study investigators, and if applicable, details of automation tools used in the process.	8
Data items	10a	List and define all outcomes for which data were sought. Specify whether all results that were compatible with each outcome domain in each study were sought (e.g. for all measures, time points, analyses), and if not, the methods used to decide which results to collect.	8,9
	10b	List and define all other variables for which data were sought (e.g. participant and intervention characteristics, funding sources). Describe any assumptions made about any missing or unclear information.	9
Study risk of bias assessment	11	Specify the methods used to assess risk of bias in the included studies, including details of the tool(s) used, how many reviewers assessed each study and whether they worked independently, and if applicable, details of automation tools used in the process.	Stated that risk of bias was not assessed on pg 8
Effect measures	12	Specify for each outcome the effect measure(s) (e.g. risk ratio, mean difference) used in the synthesis or presentation of	11

Section and Topic	Item #	Checklist item	Location where item is reported
		results.	
Synthesis methods	13a	Describe the processes used to decide which studies were eligible for each synthesis (e.g. tabulating the study intervention characteristics and comparing against the planned groups for each synthesis (item #5)).	11,12
	13b	Describe any methods required to prepare the data for presentation or synthesis, such as handling of missing summary statistics, or data conversions.	12
	13c	Describe any methods used to tabulate or visually display results of individual studies and syntheses.	12
	13d	Describe any methods used to synthesize results and provide a rationale for the choice(s). If meta-analysis was performed, describe the model(s), method(s) to identify the presence and extent of statistical heterogeneity, and software package(s) used.	12
	13e	Describe any methods used to explore possible causes of heterogeneity among study results (e.g. subgroup analysis, meta-regression).	13
	13f	Describe any sensitivity analyses conducted to assess robustness of the synthesized results.	13
Reporting bias assessment	14	Describe any methods used to assess risk of bias due to missing results in a synthesis (arising from reporting biases).	Stated that risk of bias was not assessed on pg 8
Certainty assessment	15	Describe any methods used to assess certainty (or confidence) in the body of evidence for an outcome.	Stated that no methods were used to assess certainty in the body of evidence for an outcome on pg 13
RESULTS			
Study selection	16a	Describe the results of the search and selection process, from the number of records identified in the search to the number of studies included in the review, ideally using a flow diagram.	15
	16b	Cite studies that might appear to meet the inclusion criteria, but which were excluded, and explain why they were excluded.	15
Study characteristics	17	Cite each included study and present its characteristics.	Citations for included studies presented in supplementary 3.
Risk of bias in studies	18	Present assessments of risk of bias for each included study.	Stated that risk of bias was not assessed on pg 8
Results of individual studies	19	For all outcomes, present, for each study: (a) summary statistics for each group (where appropriate) and (b) an effect estimate and its precision (e.g. confidence/credible interval), ideally using structured tables or plots.	15-21
Results of	20a	For each synthesis, briefly summarise the characteristics and risk of bias among contributing studies.	15-21

Section and Topic	Item #	Checklist item	Location where item is reported
syntheses	20b	Present results of all statistical syntheses conducted. If meta-analysis was done, present for each the summary estimate and its precision (e.g. confidence/credible interval) and measures of statistical heterogeneity. If comparing groups, describe the direction of the effect.	15-21
	20c	Present results of all investigations of possible causes of heterogeneity among study results.	15-21
	20d	Present results of all sensitivity analyses conducted to assess the robustness of the synthesized results.	16-20
Reporting biases	21	Present assessments of risk of bias due to missing results (arising from reporting biases) for each synthesis assessed.	Stated that risk of bias was not assessed on pg 8
Certainty of evidence	22	Present assessments of certainty (or confidence) in the body of evidence for each outcome assessed.	Stated that no methods were used to assess certainty in the body of evidence for an outcome on pg 13
DISCUSSION			
Discussion	23a	Provide a general interpretation of the results in the context of other evidence.	22-30
	23b	Discuss any limitations of the evidence included in the review.	30
	23c	Discuss any limitations of the review processes used.	30
	23d	Discuss implications of the results for practice, policy, and future research.	31
OTHER INFORMATION			
Registration and protocol	24a	Provide registration information for the review, including register name and registration number, or state that the review was not registered.	Stated that this is a follow-on review from a previous review.
	24b	Indicate where the review protocol can be accessed, or state that a protocol was not prepared.	No protocol was prepared.
	24c	Describe and explain any amendments to information provided at registration or in the protocol.	No protocol was prepared.
Support	25	Describe sources of financial or non-financial support for the review, and the role of the funders or sponsors in the review.	31
Competing interests	26	Declare any competing interests of review authors.	31
Availability of data, code and other materials	27	Report which of the following are publicly available and where they can be found: template data collection forms; data extracted from included studies; data used for all analyses; analytic code; any other materials used in the review.	Supplementary files include checklist, conversion chart, and included references.

Supplementary File 2: Table outlining %1RM estimation based on repetitions performed, adapted from Haff and Triplett [19].

Repetitions Performed	%1RM
1	100
2	95
3	93
4	90
5	87
6	85
7	83
8	80
9	77
10	75
11	70
12	67
15	65

Supplementary File 3: Further details of statistical approach

All meta-analyses were conducted using a nested four-level mixed effects meta-analytic model. The series of nestings included the individual study (level 4), the outcome (level 3), the measurement occasion (level 2) as many studies included more than just pre- and post-intervention assessments, and the within study sampling variance (level 1). A representation of the meta-analyses conducted includes:

$$\text{Level1: } d_{ijk} = \beta_{0ijk} + e_{ijk}, \quad e_{ijk} \sim N(0, \sigma_e^2)$$

$$\text{Level2: } \beta_{0ijk} = \eta_{0jk} + \beta_{2,1}x_{2,1ijk} + r_{ijk}, \quad r_{ijk} \sim N(0, \sigma_r^2)$$

$$\text{Level3: } \eta_{0jk} = \theta_{0k} + \beta_{3,1}x_{3,1jk} + \beta_{3,2}(x_{3,1jk} * x_{4,1k}) + u_{0jk}, \quad u_{0jk} \sim N(0, \sigma_u^2)$$

$$\text{Level4: } \theta_{0k} = \gamma_0 + \beta_{4,1}x_{4,1k} + s(x_{4,2k}) + v_{0k}, \quad v_{0k} \sim N(0, \sigma_v^2)$$

where d_{ijk} is the observed effect size at measurement occasion i ($i = 1, 2, \dots, I_{jk}$), from outcome j ($j = 1, 2, \dots, J_k$) and from study k ($k = 1, 2, \dots, K$). The indexing I_{jk} denotes that the number of measurement occasions may vary across outcomes and studies, and J_k denotes the number of outcomes may vary across studies. The random effects across the different levels ($v_{0k}, u_{0jk}, r_{ijk}, e_{ijk}$) were assumed to be independent. β terms represent regression coefficients for the predictor variables x included at levels 2 to 4. Cross-level interactions are denoted by $*$ and for some continuous predictors, smooth functions (simple basis functions) were used to model non-linear effects of the predictor and are denoted by $s(x)$.

Candidate models were fitted and compared based on predictive accuracy using the theoretical expected log pointwise predictive density (ELPD) for a new dataset that was estimated with leave-one-out cross validation (ELPD-LOO) [23]. The ELPD-LOO generates a standard error that describes the uncertainty in the predictive performance for unknown future data. Candidate

models were fit gradually increasing the number of predictors starting at level 2 and progressing to level 4. The ELPD-LOO difference between a new and previous model was calculated and addition of the predictor judged as an improvement and maintained in subsequent models if the value was at least two times the standard error.

Supplementary File 4: Reference list of included studies

1. Hoffman JR, Cooper J, Wendell M, Kang J. Comparison of Olympic vs. traditional power lifting training programs in football players. *Journal of Strength and Conditioning Research*. 2004; 18(1):129-35.
2. Balabinis CP, Psarakis CH, Moukas M, Vassiliou MP, Behrakis PK. Early phase changes by concurrent endurance and strength training. *Journal of Strength and Conditioning Research*. 2003; 17(2):393-401.
3. Carlson K, Magnusen M, Walters P. Effect of various training modalities on vertical jump. *Research in Sports Medicine*. 2009; 17(2):84-94.
4. Kyrolainen H, Avela J, McBride JM, Koskinen S, Andersen JL, Sipila S, et al. Effects of power training on muscle structure and neuromuscular performance. *Scandinavian Journal of Medicine and Science in Sports*. 2005; 15(1):58-64.
5. Whitehead MT, Scheett TP, McGuigan MR, Auckland NZ, Martin AV. A Comparison of the Effects of Short-Term Plyometric and Resistance Training on Lower Body Muscular Performance. *Journal of Strength and Conditioning Research*. 2018; 32(10):2743-2749
6. Zaras N, Spengos K, Methenitis S, Papadopoulos C, Karampatsos G, Georgiadis G, Stasinaki A, Manta P, Terzis G. Effects of strength vs. ballistic-power training on throwing performance. *Journal of sports science & medicine*. 2013; 12(1):130.
7. Argus CK, Gill ND, Keogh JWL, McGuigan MR, Hopkins WG. Effects of Two Contrast Training Programs on Jump Performance in Rugby Union Players During a Competition Phase. *International Journal of Sports Physiology & Performance*. 2012;7(1):68-75
8. Holmstrup ME, Jensen BT, Evans WS, Marshall EC. Eight Weeks of Kettlebell Swing Training Does not Improve Sprint Performance in Recreationally Active Females. *International Journal of Exercise Science*. 2016; 9(3):437-444.
9. Ronnestad BR, Kojedal O, Losnegard T, Kvamme B, Raastad T. Effect of heavy strength training on muscle thickness, strength, jump performance, and endurance performance in well-trained Nordic Combined athletes. *European journal of applied physiology*. 2012; 112(6):2341-2352.
10. Folland JP, Irish CS, Roberts JC, Tarr JE, Jones DA. Fatigue is not a necessary stimulus for strength gains during resistance training. *British journal of sports medicine*. 2002; 36(5):370-373.
11. Norrbrand L, Pozzo M, Tesch PA. Flywheel resistance training calls for greater eccentric muscle activation than weight training. *European journal of applied physiology*. 2010; 110(5):997-1005.
12. Sander A, Keiner M, Wirth K, Schmidtbleicher D. Influence of a 2-year strength training programme on power performance in elite youth soccer players. *European journal of sport science*. 2013; 13(5):445-51.
13. Gonzalez-Badillo JJ, Rodriguez-Rosell D, Sanchez-Medina L, Gorostiaga EM, Pareja-Blanco F. Maximal intended velocity training induces greater gains in bench press performance than deliberately slower half-velocity training. *European journal of sport science*. 2014; 14(8):772-781.
14. Cantrell GS, Schilling BK, Paquette MR, Murlasits Z. Maximal strength, power, and aerobic endurance adaptations to concurrent strength and sprint interval training. *European journal of applied physiology*. 2014; 114(4):763-71.
15. Sarabia JM, Fernandez-Fernandez J, Juan-Recio C, Hernández-Davó H, Urbán T, Moya M. Mechanical, hormonal and psychological effects of a non-failure short-term strength training program in young tennis players. *Journal of human kinetics*. 2015; 45:81-91.
16. Remaud A, Cornu C, Guevel A. Neuromuscular adaptations to 8-week strength training: Isotonic versus isokinetic mode. *European journal of applied physiology*. 2010; 108(1):59-69.
17. Harris GR, Stone MH, O'Bryant HS, Proulx CM, Johnson RL. Short-term performance effects of high power, high force, or combined weight-training methods. *Journal of Strength and Conditioning Research*. 2000; 14(1):14-20.

18. Brito J, Vasconcellos F, Oliveira J, Krusturup P, Rebelo A. Short-term performance effects of three different low-volume strength-training programmes in college male soccer players. *Journal of human kinetics*. 2014; 40:121-128.
19. Cholewa JM, Rossi FE, MacDonald C, Hewins A, Gallo S, Micenski A, et al. The effects of moderate- versus high-load resistance training on muscle growth, body composition, and performance in collegiate women. *Journal of Strength and Conditioning Research*. 2018; 32(6):1511-24.
20. James LP, Haff GG, Kelly VG, Connick M, Hoffman B, Beckman EM. The impact of strength level on adaptations to combined weightlifting, plyometric and ballistic training. *Scandinavian Journal of Medicine & Science in Sports*. 2018; 28(5):1494-505.
21. Coratella G, Milanese C, Schena F. Unilateral eccentric resistance training: A direct comparison between isokinetic and dynamic constant external resistance modalities. *European Journal of Sport Science*. 2015; 15(8):720-6.
22. Vantarakis A, Chatzinikolaou A, Avloniti A, Vezos N, Douroudos II, Draganidis D, Jamurtas AZ, Kambas A, Kalligeros S, Fatouros IG. A 2-month linear periodized resistance exercise training improved musculoskeletal fitness and specific conditioning of navy cadets. *Journal of Strength and Conditioning Research*. 2017; 31(5):1362-70.
23. Boyer, Brian T. A Comparison of the Effects of Three Strength Training Programs on Women. *Journal of Strength and Conditioning Research*. 1990; 4(3): 88-94.
24. Bartolomei S, Hoffman JR, Merni F, Stout JR. A comparison of traditional and block periodized strength training programs in trained athletes. *Journal of Strength and Conditioning Research*. 2014; 28(4):990-7.
25. Hall R. *A comparison of unilateral vs. bilateral leg strength training*. Eugene, Ore.; Microform Publications, University of Oregon; 1985TY: GENGEN; Accession Number: SPH175914; Author: Hall, R. ; Language: English; General Notes: Thesis (M.S.) - University of Arizona, 1983; includes bibliography. Available from: Microform Publications, International Institute for Sport and Human Performance, University of Oregon, Eugene, OR.; Description: 1 microfiche (41 fr.) : neg., ill.; 11 x 15 cm.; Publication Type: Microforms; Thesis or dissertation; Update Code: 19981201.
26. Cormie P, McGuigan MR, Newton RU. Adaptations in athletic performance after ballistic power versus strength training. *Medicine & Science in Sports & Exercise*. 2010; 42(8):1582-98.
27. Baker DG, Newton RU. Adaptations in upper-body maximal strength and power output resulting from long-term resistance training in experienced strength-power athletes. *Journal of Strength and Conditioning Research*. 2006; 1;20(3):541-6.
28. Stanforth PR, Painter TL, Wilmore JH. Alterations in Concentric Strength Consequent to Powercise and Universal Gym Circuit Training. *Journal of Strength and Conditioning Research*. 1992; 6(4):249-55.
29. Rhea MR, Kenn JG, Dermody BM. Alterations in speed of squat movement and the use of accommodated resistance among college athletes training for power. *Journal of Strength and Conditioning Research*. 2009; 23(9):2645-2650.
30. Wilmore J. Alterations in strength, body composition and anthropometric measurements consequent to a 10-week weight training program. *Medicine and Science in Sports*. 1974; 6(2):133-8.
31. Rhea MR, Peterson MD, Oliverson JR, Ayllon FN, Potenziano BJ. An examination of training on the VertiMax resisted jumping device for improvements in lower body power in highly trained college athletes. *Journal of Strength and Conditioning Research*. 2008; 22(3):735-740.
32. Brown BS, Gorman DR, DiBrezzo R, Fort I. Anaerobic power changes following short term, task specific, dynamic and static overload training. *Journal of Strength and Conditioning Research*. 1988; 2(2):35-8.
33. Newton RU, McEvoy KP. Baseball throwing velocity: A comparison of medicine ball training and weight training. *Journal of Strength and Conditioning Research*. 1994; 8(3):198-203.
34. Bartolomei S, Stout JR, Fukuda DH, Hoffman JR, Merni F. Block vs. weekly undulating periodized resistance training programs in women. *Journal of Strength and Conditioning Research*. 2015; 29(10):2679-87.

35. Abt JP, Oliver JM, Nagai T, Sell TC, Lovalekar MT, Beals K, et al. Block-Periodized Training Improves Physiological and Tactically Relevant Performance in Naval Special Warfare Operators. *Journal of Strength and Conditioning Research*. 2016; 30(1):39-52.
36. Siegler J, Gaskill S, Ruby B. Changes evaluated in soccer-specific power endurance either with or without a 10-week, in-season, intermittent, high-intensity training protocol. *Journal of Strength and Conditioning Research*. 2003; 17(2):379-387.
37. Frost DM, Bronson S, Cronin JB, Newton RU. Changes in Maximal Strength, Velocity, and Power After 8 Weeks of Training With Pneumatic or Free Weight Resistance. *Journal of Strength and Conditioning Research*. 2016; 30(4):934-944.
38. Lloyd RS, Radnor JM, Croix MB, Cronin JB, Oliver JL. Changes in sprint and jump performances after traditional, plyometric, and combined resistance training in male youth pre-and post-peak height velocity. *Journal of Strength and Conditioning Research*. 2016; 30(5):1239-47.
39. de Hoyo M, Gonzalo-Skok O, Sañudo B, Carrascal C, Plaza-Armas JR, Camacho-Candil F, Otero-Esquina C. Comparative effects of in-season full-back squat, resisted sprint training, and plyometric training on explosive performance in U-19 elite soccer players. *Journal of Strength and Conditioning Research*. 2016; 30(2):368-77.
40. Mihalik JP, Libby JJ, Battaglini CL, McMurray RG. Comparing short-term complex and compound training programs on vertical jump height and power output. *Journal of Strength and Conditioning Research*. 2008; 22(1):47-53.
41. Teo SY, Newton MJ, Newton RU, Dempsey AR, Fairchild TJ. Comparing the effectiveness of a short-term vertical jump vs. weightlifting program on athletic power development. *Journal of Strength and Conditioning Research*. 2016; 30(10):2741-8.
42. Hoffman JR, Ratamess NA, Klatt M, Faigenbaum AD, Ross RE, Tranchina NM, McCurley RC, Kang J, Kraemer WJ. Comparison between different off-season resistance training programs in Division III American college football players. *Journal of Strength and Conditioning Research*. 2009; 23(1):11-19.
43. Bartolomei S, Hoffman JR, Stout JR, Zini M, Stefanelli C, Merni F. Comparison of Block Versus Weekly Undulating Periodization Models on Endocrine and Strength Changes in Male Athletes. *Kinesiology*. 2016; 48(1):71-78.
44. Schmidtbleicher D, Wirth K. Comparison of different strength trainings methods for the development of power. [Accession Number: SPHS-1059438; Conference: International Symposium on Biomechanics in Sports (24th : 2006 : Salzburg, Austria)].
45. Rana SR, Chleboun GS, Gilders RM, Hagerman FC, Herman JR, Hikida RS, et al. Comparison of early phase adaptations for traditional strength and endurance, and low velocity resistance training programs in college-aged women. *Journal of Strength and Conditioning Research*. 2008; 22(1):119-127.
46. Harries SK, Lubans DR, Callister R. Comparison of resistance training progression models on maximal strength in sub-elite adolescent rugby union players. *Journal of Science and Medicine in Sport*. 2016; 19(2):163-9.
47. Bauer T, Thayer RE, Baras G. Comparison of training modalities for power development in the lower extremity. *Journal of Strength and Conditioning Research*. 1990; 4(4):115-21.
48. Aarskog R, Wisnes A, Wilhelmsen K, Skogen A, Bjordal JM. Comparison of Two Resistance Training Protocols, 6RM versus 12RM, to Increase the 1RM in Healthy Young Adults. A Single-Blind, Randomized Controlled Trial. *Physiotherapy Research International*. 2012; 17(3):179-186.
49. Hori N, Newton RU, Kawamori N, McGuigan MR, Andrews WA, Chapman DW, et al. Comparison of weighted jump squat training with and without eccentric braking. *Journal of Strength and Conditioning Research*. 2008; 22(1):54-65.
50. Robineau J, Lacombe M, Piscione J, Bigard X, Babault N. Concurrent Training in Rugby Sevens: Effects of High-Intensity Interval Exercises. *International journal of sports physiology and performance*. 2017; 12(3):336-344.

51. Hansen KT, Cronin JB, Pickering SL, Newton MJ. Does cluster loading enhance lower body power development in preseason preparation of elite rugby union players?. *Journal of Strength and Conditioning Research*. 2011; 25(8):2118-26.
52. Milanovic Z, Sporis G, Trajkovic N, Sekulic D, James N, Vuckovic G. Does SAQ training improve the speed and flexibility of young soccer players? A randomized controlled trial. *Human Movement Science*. 2014; 38:197-208.
53. Radaelli R, Fleck SJ, Leite T, Leite RD, Pinto RS, Fernandes L, Simão R. Dose-response of 1, 3, and 5 sets of resistance exercise on strength, local muscular endurance, and hypertrophy. *Journal of Strength and Conditioning Research*. 2015; 29(5):1349-58.
54. Atae J, Koozehchian MS, Kreider RB, Zuo L. Effectiveness of accommodation and constant resistance training on maximal strength and power in trained athletes. *PeerJ*. 2014; 2014(1) (pagination):Arte Number: e441. ate of Pubaton: 2014.
55. Kerksick CM, Wilborn CD, Campbell BI, Roberts MD, Rasmussen CJ, Greenwood M, Kreider RB. Early-phase adaptations to a split-body, linear periodization resistance training program in college-aged and middle-aged men. *Journal of Strength and Conditioning Research*. 2009; 23(3):962-71.
56. Wirth K, Keiner M, Hartmann H, Sander A, Mickel C. Effect of 8 weeks of free-weight and machine-based strength training on strength and power performance. *Journal of human kinetics*. 2016; 53:201.
57. Laurent C, Penzer F, Letroye B, Carpentier A, Baudry S, Duchateau J. Effect of a strength training method characterized by an incremental number of repetitions across sets and a very short rest interval. *Science and Sports*. 2016; 31(5):e115-e121.
58. Manolopoulos K, Gissis I, Galazoulas C, Manolopoulos E, Patikas D, Gollhofer A., et al. Effect of Combined Sensorimotor-Resistance Training on Strength, Balance, and Jumping Performance of Soccer Players. *Journal of Strength and Conditioning Research*. 2016; 30(1):53-59.
59. Koundourakis NE, Androulakis N, Spyridaki EC, Castanas E, Malliaraki N, Tsatsanis C, et al. Effect of different seasonal strength training protocols on circulating androgen levels and performance parameters in professional soccer players. *Hormones*. 2014; 13(1):104-118.
60. Coutts AJ, Murphy AJ, Dascombe BJ. Effect of direct supervision of a strength coach on measures of muscular strength and power in young rugby league players. *Journal of Strength and Conditioning Research*. 2004; 18(2):316-323.
61. Randell AD, Cronin JB, Keogh JW, Gill ND, Pedersen MC. Effect of instantaneous performance feedback during 6 weeks of velocity-based resistance training on sport-specific performance tests. *Journal of Strength and Conditioning Research*. 2011; 25(1):87-93.
62. Bartolomei S, Hoffman JR, Stout JR, Merni F. Effect of Lower-Body Resistance Training on Upper-Body Strength Adaptation in Trained Men. *Journal of Strength and Conditioning Research*. 2018; 32(1):13-18.
63. Pareja-Blanco F, Rodriguez-Rosell D, Sanchez-Medina L, Gorostiaga EM, Gonzalez-Badillo JJ. Effect of movement velocity during resistance training on neuromuscular performance. *International Journal of Sports Medicine*. 2014; 35(11):916-924.
64. Channell BT, Barfield JP. Effect of Olympic and traditional resistance training on vertical jump improvement in high school boys. *Journal of Strength and Conditioning Research*. 2008; 22(5):1522-1527.
65. Korak JA, Paquette MR, Brooks J, Fuller DK, Coons JM. Effect of rest-pause vs. traditional bench press training on muscle strength, electromyography, and lifting volume in randomized trial protocols. *European journal of applied physiology*. 2017; 117(9):1891-6.
66. Blazeovich AJ, Jenkins DG. Effect of the movement speed of resistance training exercise on sprint and strength performance in concurrently training elite junior sprinters. *Journal of Orthopaedic & Sports Physical Therapy*. 2002; 20(12):981-90.
67. Lamberth J, Hale B, Knight A, Boyd J, Luczak T. Effectiveness of a Six-Week Strength and Functional Training Program on Golf Performance. *International Journal of Golf Science*. 2013; 2(1):33-42.

68. Jimenez-Reyes P, Samozino P, Brughelli M, Morin JB. Effectiveness of an individualized training based on force-velocity profiling during jumping. *Frontiers in Physiology*. 2017; 7(JAN) (pagination):Arte Number: 677. ate of Pubaton: 2017.
69. Brown AC, Wells TJ, Schade ML, Smith DL, Fehling PC. Effects of Plyometric Training Versus Traditional Weight Training on Strength, Power, and Aesthetic Jumping Ability in Female Collegiate Dancers. *Journal of Dance Medicine & Science*. 2007; 11(2):38-44.
70. Paz-Franco A, Rey E, Barcala-Furelos R. Effects of 3 Different Resistance Training Frequencies on Jump, Sprint, and Repeated Sprint Ability Performances in Professional Futsal Players. *Journal of Strength and Conditioning Research*. 2017; 31(12):3343-3350.
71. Kim E, Dear A, Ferguson SL, Seo D, Bembem MG. Effects of 4 weeks of traditional resistance training vs. superslow strength training on early phase adaptations in strength, flexibility, and aerobic capacity in college-aged women. *Journal of Strength and Conditioning Research*. 2011; 25(11):3006-13.
72. Rodríguez-Rosell D, Franco-Márquez F, Pareja-Blanco F, Mora-Custodio R, Yáñez-García JM, González-Suárez JM, González-Badillo JJ. Effects of 6 weeks resistance training combined with plyometric and speed exercises on physical performance of pre-peak-height-velocity soccer players. *International journal of sports physiology and performance*. 2016; 11(2):240-6.
73. Hermassi S, Chelly MS, Tabka Z, Shephard RJ, Chamari K. Effects of 8-week in-season upper and lower limb heavy resistance training on the peak power, throwing velocity, and sprint performance of elite male handball players. *Journal of Strength and Conditioning Research*. 2011; 25(9):2424-2433.
74. Wong PL, Chamari K, Wisløff U. Effects of 12-week on-field combined strength and power training on physical performance among U-14 young soccer players. *Journal of Strength and Conditioning Research*. 2010; 24(3):644-52.
75. Harries SK, Lubans DR, Buxton A, MacDougall THJ, Callister R. Effects of 12-weeks resistance training on sprint and jump performance in competitive adolescent rugby union players. *Journal of Strength and Conditioning Research*. 2018; 32(10):2762-9.
76. Veliz RR, Requena B, Suarez-Arrones L, Newton RU, De Villarreal ES. Effects of 18-week in-season heavy-resistance and power training on throwing velocity, strength, jumping, and maximal sprint swim performance of elite male water polo players. *Journal of Strength and Conditioning Research*. 2014; 28(4):1007-14.
77. Manolopoulos E, Katis A, Manolopoulos K, Kalapotharakos V, Kellis E. Effects of a 10-week resistance exercise program on soccer kick biomechanics and muscle strength. *Journal of Strength and Conditioning Research*. 2013; 27(12):3391-3401.
78. Redondo JC, Alonso CJ, Sedano S, de Benito AM. Effects of a 12-week strength training program on experimented fencers' movement time. *Journal of Strength and Conditioning Research*. 2014; 28(12):3375-84.
79. Chelly MS, Fathloun M, Cherif N, Amar MB, Tabka Z, Van Praagh E. Effects of a back squat training program on leg power, jump, and sprint performances in junior soccer players. *Journal of Strength and Conditioning Research*. 2009; 23(8):2241-9.
80. Faigenbaum AD, McFarland JE, Keiper FB, Tevlin W, Ratamess NA, Kang J, Hoffman JR. Effects of a short-term plyometric and resistance training program on fitness performance in boys age 12 to 15 years. *Journal of sports science & medicine*. 2007; 6(4):519-525.
81. Contreras B, Vigotsky AD, Schoenfeld BJ, Beardsley C, McMaster DT, Reyneke JH, Cronin JB. Effects of a six-week hip thrust vs. front squat resistance training program on performance in adolescent males: a randomized controlled trial. *Journal of Strength and Conditioning Research*. 2017; 31(4):999-1008.
82. Inovero JG, Pagaduan JC. Effects of a Six-Week Strength Training and Upper Body Plyometrics in Male College Basketball Physical Education Students. *Sport Scientific & Practical Aspects*. 2015; 12(1):11-16.
83. Bourgeois FA, Gamble P, Gill ND, McGuigan MR. Effects of a six-week strength training programme on change of direction performance in youth team sport athletes. *Sports*. 2017; 5(4):83.

84. Vaara JP, Kokko J, Isoranta M, Kyrolainen H. Effects of Added Resistance Training on Physical Fitness, Body Composition, and Serum Hormone Concentrations During Eight Weeks of Special Military Training Period. *Journal of Strength and Conditioning Research*. 2015; 29:S168-S172.
85. Robbins DW, Young WB, Behm DG, Payne WR. Effects of agonist-antagonist complex resistance training on upper body strength and power development. *Journal of sports sciences*. 2009; 27(14):1617-1625.
86. Alvarez M, Sedano S, Cuadrado G, Redondo JC. Effects of an 18-week strength training program on low-handicap golfers' performance. *Journal of Strength and Conditioning Research*. 2012; 26(4):1110-1121.
87. Smart DJ, Gill ND. Effects of an off-season conditioning program on the physical characteristics of adolescent rugby union players. *Journal of Strength and Conditioning Research*. 2013; 27(3):708-17.
88. Newton RU, Kraemer WJ, Hakkinen K. Effects of ballistic training on preseason preparation of elite volleyball players. *Medicine and Science in Sports and Exercise*. 1999; 31(2):323-330.
89. Franco Márquez F, Rodríguez Rosell D, González Suárez JM, Pareja Blanco F, Mora Custodio R, Yáñez García JM, González Badillo JJ. Effects of combined resistance training and plyometrics on physical performance in young soccer players. *International Journal of Sports Medicine*. 2015.
90. Santos EJAM, Janeira MAAS. Effects of complex training on explosive strength in adolescent male basketball players. *Journal of Strength and Conditioning Research*. 2008; 22(3):903-9.
91. Hartmann H, Bob A, Wirth K, Schmidtbleicher D. Effects of different periodization models on rate of force development and power ability of the upper extremity. *Journal of Strength and Conditioning Research*. 2009; 23(7):1921-32.
92. Naclerio F, Faigenbaum AD, Larumbe-Zabala E, Perez-Bibao T, Kang J, Ratamess NA, et al. Effects of different resistance training volumes on strength and power in team sport athletes. *Journal of Strength and Conditioning Research*. 2013; 27(7):1832-1840.
93. Robinson JM, Stone MH, Johnson RL, Penland CM, Warren BJ, Lewis RD. Effects of different weight training exercise/rest intervals on strength, power, and high intensity exercise endurance. *Journal of Strength and Conditioning Research*. 1995; 9(4):216-21.
94. Wirth K, Keiner M, Szilvas E, Hartmann H, Sander A. Effects of eccentric strength training on different maximal strength and speed-strength parameters of the lower extremity. *Journal of Strength and Conditioning Research*. 2015; 29(7):1837-45.
95. Tomljanović M, Spasić M, Gabrilo G, Uljević O, Foretić N. Effects of Five Weeks of Functional Vs. Traditional Resistance Training on Anthropometric and Motor Performance Variables. / Efekti 5-Tjednog Funkcionalnoga i Tradicionalnogatreninga S Opterećenjem Na Antropometrijskekarakteristike i Motoričke Sposobnosti. *Kinesiology*. 2011; 43(2):145-154.
96. Gorostiaga EM, Izquierdo M, Iturralde P, Ruesta M, Ibanez J. Effects of heavy resistance training on maximal and explosive force production, endurance and serum hormones in adolescent handball players. *European journal of applied physiology and occupational physiology*. 1999; 80(5):485-493.
97. Janusevicius D, Snieckus A, Skurvydas A, Silinskas V, Trinkunas E, Cadefau JA, Kamandulis S. Effects of high velocity elastic band versus heavy resistance training on hamstring strength, activation, and sprint running performance. *Journal of sports science & medicine*. 2017; 16(2):239.
98. Pierce K, Rozenek R, Stone MH. Effects of high volume weight training on lactate, heart rate, and perceived exertion. *Journal of Strength and Conditioning Research*. 1993; 7(4):211-5.
99. Negra Y, Chaabene H, Hammami M, Hachana Y, Granacher U. Effects of high-velocity resistance training on athletic performance in prepuberal male soccer athletes. *Journal of Strength and Conditioning Research*. 2016; 30(12):3290-7.
100. Kubo K, Yata H, Kanehisa H, Fukunaga T. Effects of isometric squat training on the tendon stiffness and jump performance. *European journal of applied physiology*. 2006; 96(3):305-314.
101. Rodríguez-Rosell D, Torres-Torrel J, Franco-Márquez F, González-Suárez JM, González-Badillo JJ. Effects of light-load maximal lifting velocity weight training vs. combined weight training and plyometrics on sprint, vertical jump and strength performance in adult soccer players. *Journal of science and medicine in sport*. 2017; 20(7):695-9.

102. Wirtz N, Zinner C, Doermann U, Kleinoeder H, Mester J. Effects of Loaded Squat Exercise with and without Application of Superimposed EMS on Physical Performance. *Journal of Sports Science and Medicine*. 2016; 15(1):26-33.
103. Hammami M, Negra Y, Billaut F, Hermassi S, Shephard RJ, Chelly MS. Effects of Lower-Limb Strength Training on Agility, Repeated Sprinting With Changes of Direction, Leg Peak Power, and Neuromuscular Adaptations of Soccer Players. *Journal of Strength and Conditioning Research*. 2018; 32(1):37-47.
104. Moss BM, Refsnes PE, Abildgaard A, Nicolaysen K, Jensen J. Effects of maximal effort strength training with different loads on dynamic strength, cross-sectional area, load-power and load-velocity relationships. *European journal of applied physiology and occupational physiology*. 1997; 75(3):193-9.
105. Karsten B, Larumb-Zabala E, Kandemir G, Hazir T, Klose A, Naclerio F. The effects of a 6-week strength training on critical velocity, anaerobic running distance, 30-M sprint and Yo-Yo intermittent running test performances in male soccer players. *PLoS ONE*. 2016; 11(3) (pagination):Arte Number: e0151448. ate of Pubaton: Marh 2016.
106. Bruhn S, Kullmann N, Gollhofer A. The Effects of a Sensorimotor Training and a Strength Training on Postural Stabilisation, Maximum Isometric Contraction and Jump Performance. *International Journal of Sports Medicine*. 2004; 25(1):56-60.
107. Zisis P. The effects of an 8 weeks plyometric training program or an explosive strength training program on the Jump-and-Reach Height of male amateur soccer players. *Journal of Physical Education and Sport*. 2013; 13(4):594-600.
108. Fry AC, Kraemer WJ, Weseman CA, Conroy BP, Gordon SE, Hoffman JR, Maresh CM. The effects of an off-season strength and conditioning program on starters and non-starters in women's intercollegiate volleyball. *Journal of Strength and Conditioning Research*. 1991; 5(4):174-81.
109. Mangine GT, Ratamess NA, Hoffman JR, Faigenbaum AD, Kang J., Chilakos A. The effects of combined ballistic and heavy resistance training on maximal lower- and upper-body strength in recreationally trained men. *Journal of Strength and Conditioning Research*. 2008; 22(1):132-139.
110. Apanukul S, Suwannathada S, Intiraporn C. The Effects of Combined Weight and Pneumatic Training to Enhance Power Endurance in Tennis Players. *Journal of Exercise Physiology Online*. 2015; 18(2):8-16.
111. Anderson CE, Sforzo GA, Sigg JA. The effects of combining elastic and free weight resistance on strength and power in athletes. *Journal of strength and conditioning research*. 2008; 22(2):567-574.
112. Pritchard HJ, Fink PW, Stannard SR. The Effects of Concentric/eccentric Training Versus Concentric Only Training on Peak Power and Functional Muscle Performance. *Journal of Australian Strength and Conditioning*. 2015; 23(6):71-75.
113. Lockie RG, Murphy AJ, Schultz AB, Knight TJ, de Jonge XA. The effects of different speed training protocols on sprint acceleration kinematics and muscle strength and power in field sport athletes. *Journal of Strength and Conditioning Research*. 2012; 26(6):1539-50.
114. Dolezal SM, Frese DL, Llewellyn TL. The Effects of Eccentric, Velocity-Based Training on Strength and Power in Collegiate Athletes. *International Journal of Exercise Science*. 2016; 9(5):657-666.
115. Hoff J, Almasbakk B. The effects of maximum strength training on throwing velocity and muscle strength in female team-handball players. *Journal of Strength and Conditioning Research*. 1995; 9(4):255-258.
116. Liu C, Chen CS, Ho WH, Fule RJ, Chung PH, Shiang TY. The effects of passive leg press training on jumping performance, speed, and muscle power. *Journal of Strength and Conditioning Research*. 2013; 27(6):1479-1486.
117. Herrick AB, Stone WJ. The effects of periodization versus progressive resistance exercise on upper and lower body strength in women. *Journal of Strength and Conditioning Research*. 1996 May 1;10(2):72-6.
118. Santos EJAM, Janeira MAAS. The effects of resistance training on explosive strength indicators in adolescent basketball players. *Journal of Strength and Conditioning Research*. 2012; 26(10):2641-7.
119. Smith RA, Martin GJ, Szivak TK, Comstock BA, Dunn-Lewis C, Hooper DR, Flanagan SD, Looney DP, Volek JS, Maresh CM, Kraemer WJ. The effects of resistance training prioritization in

- NCAA Division I Football summer training. *Journal of Strength and Conditioning Research*. 2014; 28(1):14-22.
120. Hoffman JR, Kraemer WJ, Fry AC, Deschenes M, Kemp M. The effects of self-selection for frequency of training in a winter conditioning program for football. *Journal of Strength and Conditioning Research*. 1990; 4(3):76-82.
 121. Stalder MA, Noble BJ, Wilkinson JG. The effects of supplemental weight training for ballet dancers. *Journal of Strength and Conditioning Research*. 1990; 4(3):95-102.
 122. Cressey EM, West CA, Tiberio DP, Kraemer WJ, Maresh CM. The effects of ten weeks of lower-body unstable surface training on markers of athletic performance. *Journal of Strength and Conditioning Research*. 2007; 21(2):561-567.
 123. Wirth K, Keiner M, Szilvas E, Hartmann H, Sander A. Effects of eccentric strength training on different maximal strength and speed-strength parameters of the lower extremity. *Journal of Strength and Conditioning Research*. 2015; 29(7):1837-45.
 124. Sleivert GG, Backus RD, Wenger HA. The influence of a strength-sprint training sequence on multi-joint power output. *Medicine and science in sports and exercise*. 1995; 27(12):1655-1665.
 125. Mazzetti SA, Kraemer WJ, Volek JS, Duncan ND, Ratamess NA, Gomez AL, et al. The influence of direct supervision of resistance training on strength performance. *Medicine and science in sports and exercise*. 2000; 32(6):1175-1184.
 126. Bishop D, Jenkins DG. The influence of resistance training on the critical power function and time to fatigue at critical power. *Australian Journal of Science and Medicine in Sport*. 1996; 28(4):101-105.
 127. Bloomfield J, Blanksby BA, Ackland TR, Allison GT. The influence of strength training on overhead throwing velocity of elite water polo players. *Australian Journal of Science and Medicine in Sport*. 1990; 22(3):63-67.
 128. Petersen SR, Bagnall KM, Wenger HA, Reid DC, Castor WR, Quinney HA. The influence of velocity-specific resistance training on the in vivo torque-velocity relationship and the cross-sectional area of quadriceps femoris. *Journal of Orthopaedic & Sports Physical Therapy*. 1989; 10(11):456-462.
 129. Hennessy LC, Watson AW. The interference effects of training for strength and endurance simultaneously. *Journal of Strength and Conditioning Research*. 1994; 8(1):12-9.
 130. Wilson GJ, Newton RU, Murphy AJ, Humphries BJ. The optimal training load for the development of dynamic athletic performance. *Medicine and science in sports and exercise*. 1993; 25(11):1279-1286.
 131. Schiøtz MK, Potteiger JA, Huntsinger PG, Denmark LC. The short-term effects of periodized and constant-intensity training on body composition, strength, and performance. *Journal of Strength and Conditioning Research*. 1998; 12(3):173-8.
 132. Secomb JL, Farley OR, Nimphius S, Lundgren L, Tran TT, Sheppard JM. The training-specific adaptations resulting from resistance training, gymnastics and plyometric training, and non-training in adolescent athletes. *International Journal of Sports Science & Coaching*. 2017; 12(6):762-773.
 133. Sparkes R, Behm DG. Training adaptations associated with an 8-week instability resistance training program with recreationally active individuals. *Journal of Strength and Conditioning Research*. 2010; 24(7):1931-1941.
 134. Blazeovich AJ, Gill ND, Bronks R., Newton RU. Training-Specific Muscle Architecture Adaptation after 5-wk Training in Athletes. *Medicine and science in sports and exercise*. 2003; 35(12):2013-2022.
 135. Speirs DE, Bennett MA, Finn CV, Turner AP. Unilateral vs. bilateral squat training for strength, sprints, and agility in academy rugby players. *Journal of Strength and Conditioning Research*. 2016; 30(2):386-92.
 136. Rivière M, Louit L, Strokosch A, Seitz LB. Variable resistance training promotes greater strength and power adaptations than traditional resistance training in elite youth rugby league players. *Journal of Strength and Conditioning Research*. 2017; 31(4):947-55.
 137. Palmieri GA. Weight training and repetition speed. *Journal of Strength and Conditioning Research*. 1987; 1(2):36-8.

138. Augustsson J, Esko A, Thomee R, Svantesson U. Weight training of the thigh muscles using closed vs. Open kinetic chain exercises: A comparison of performance enhancement. *Journal of Orthopaedic and Sports Physical Therapy*. 1998; 27(1):3-8.
139. Vanderka M, Novosád A. Weighted Squat Training with and without Counter Movement for Strength and Power Development. / Vplyv TrÉningu Drepov so ZÁŤaŽou, Bez a S Protipohybom Na Zmeny Úrovne SilovÝch a RÝchlostno-SilovÝch Schopností. *Acta Facultatis Educationis Physicae Universitatis Comenianae*. 2012; 52(1):21-27.
140. Hakkinen K, Newton RU, Gordon SE, McCormick M, Volek JS, Nindl BC, et al. Changes in muscle morphology, electromyographic activity, and force production characteristics during progressive strength training in young and older men. *Journals of Gerontology Series A-Biological Sciences & Medical Sciences*. 1998; 53(6):B415-23.
141. Andersen LL, Andersen JL, Zebis MK, Aagaard P. Early and late rate of force development: differential adaptive responses to resistance training?. *Scandinavian Journal of Medicine & Science in Sports*. 2010; 20(1):e162-169.
142. Iglesias-Soler E, Mayo X, Rio-Rodriguez D, Carballeira E, Farinas J, Fernandez-Delolmo M. Inter-repetition rest training and traditional set configuration produce similar strength gains without cortical adaptations. *Journal of sports sciences*. 2016; 34(15):1473-1484.
143. MacDonald CJ, Lamont HS, Garner JC. A comparison of the effects of 6 weeks of traditional resistance training, plyometric training, and complex training on measures of strength and anthropometrics. *Journal of Strength and Conditioning Research*. 2012; 26(2):422-31.
144. Kraemer WJ, Nindl BC, Ratamess NA, Gotshalk LA, Volek JS, Fleck SJ, et al. Changes in Muscle Hypertrophy in Women with Periodized Resistance Training. *Medicine and science in sports and exercise*. 2004; 36(4):697-708.
145. Andersen LL, Andersen JL, Magnusson SP, Suetta C, Madsen JL, Christensen LR, et al. Changes in the human muscle force-velocity relationship in response to resistance training and subsequent detraining. *Journal of applied physiology*. 2005; 99(1):87-94.
146. Bruhn S, Kullmann N, Gollhofer A. Combinatory effects of high-intensity-strength training and sensorimotor training on muscle strength. *International Journal of Sports Medicine*. 2006; 27(5):401-406.
147. Vossen JF, Kramer JF, Burke DG, Vossen DP. Comparison of dynamic push-up training and plyometric push-up training on upper-body power and strength. *Journal of Strength & Conditioning Research*. 2000; 14(3):248-253.
148. Moore E.W.G., Hickey M.S., Reiser II RF. Comparison of two twelve week off-season combined training programs on entry level collegiate soccer players' performance. *Journal of Strength and Conditioning Research*. 2005; 19(4):791-798.
149. McCarthy JP, Agre JC, Graf BK, Pozniak MA, Vailas AC. Compatibility of adaptive responses with combining strength and endurance training. *Medicine and science in sports and exercise*. 1995; 27(3):429-436.
150. Izquierdo-Gabarren M, Gonzalez De Txabarri Exposito R, Garcia-pallares J, Sanchez-medina L, De Villarreal ESS, Izquierdo M. Concurrent endurance and strength training not to failure optimizes performance gains. *Medicine & Science in Sports & Exercise*. 2010; 42(6):1191-1199.
151. Loturco I, Ugrinowitsch C, Tricoli V, Pivetti B, Roschel H. Different loading schemes in power training during the preseason promote similar performance improvements in Brazilian elite soccer players. *Journal of Strength and Conditioning Research*. 2013; 27(7):1791-1797.
152. Izquierdo M, Ibanez J, Gonzalez-Badillo JJ, Hakkinen K, Ratamess NA, Kraemer WJ, et al. Differential effects of strength training leading to failure versus not to failure on hormonal responses, strength, and muscle power gains. *Journal of applied physiology*. 2006; 100(5):1647-1656.
153. Jones MT. Effect of compensatory acceleration training in combination with accommodating resistance on upper body strength in collegiate athletes. *Open Access Journal of Sports Medicine*. 2014; 5:183-189.
154. Potteiger JA, Williford Jr HN, Blessing DL, Smidt J. Effect of two training methods on improving baseball performance variables. *Journal of Strength and Conditioning Research*. 1992; 6(1):2-6.

155. Izquierdo M, Hakkinen K, Ibanez J, Kraemer WJ, Gorostiaga EM. Effects of combined resistance and cardiovascular training on strength, power, muscle cross-sectional area, and endurance markers in middle-aged men. *European journal of applied physiology*. 2005; 94(1-2):70-75.
156. Kopal R, Loturco I, Barroso R, Gil S, Cuniyochi R, Ugrinowitsch C, et al. Effects Of Different Combinations Of Strength, Power, And Plyometric Training On The Physical Performance Of Elite Young Soccer Players. *Journal of Strength and Conditioning Research*. 2017; 31(6):1468-76.
157. Aagaard P, Simonsen EB, Trolle M, Bangsbo J, Klausen K. Effects of different strength training regimes on moment and power generation during dynamic knee extensions. *European journal of applied physiology and occupational physiology*. 1994; 69(5):382-386.
158. Kraemer WJ, Hakkinen K, Newton RU, Nindl BC, Volek JS, McCormick M, et al. Effects of heavy-resistance training on hormonal response patterns in younger vs. older men. *Journal of applied physiology*. 1999; 87(3):982-992.
159. Aminaei M, Yazdani S, Amirseifadini M. Effects of plyometric and cluster resistance training on explosive power and maximum strength in karate players. *International Journal of Applied Exercise Physiology*. 2017; 6(2):34-44.
160. Hunter JP, Marshall RN. Effects of power and flexibility training on vertical jump technique. *Medicine and science in sports and exercise*. 2002; 34(3):478-486.
161. Hong A, Hong S, Shin Y. Effects of resistance training on muscle strength, endurance, and motor unit according to ciliary neurotrophic factor polymorphism in male college students. *Journal of Sports Science & Medicine*. 2014; 13(3):680-688.
162. Lamas L, Ugrinowitsch C, Rodacki A, Pereira G., Mattos E.C.T., Kohn A.F., et al. Effects of strength and power training on neuromuscular adaptations and jumping movement pattern and performance. *Journal of Strength and Conditioning Research*. 2012; 26(12):3335-3344.
163. Izquierdo M, Hakkinen K, Ibanez J, Garrues M, Anton A, Zuniga A, et al. Effects of strength training on muscle power and serum hormones in middle-aged and older men. *Journal of applied physiology*. 2001; 90(4):1497-1507.
164. de Hoyo M, Sañudo B, Carrasco L, Domínguez-Cobo S, Mateo-Cortes J, Cadenas-Sánchez MM, Nimphius S. Effects of traditional versus horizontal inertial flywheel power training on common sport-related tasks. *Journal of human kinetics*. 2015; 47:155-167.
165. Pareja-Blanco F, Rodriguez-Rosell D, Sanchez-Medina L, Sanchis-Moysi J, Dorado C, Mora-Custodio R, et al. Effects of velocity loss during resistance training on athletic performance, strength gains and muscle adaptations. *Scandinavian Journal of Medicine and Science in Sports*. 2017; 27(7):724-735.
166. Arazi H, Asadi A, Roohi S. Enhancing muscular performance in women: Compound versus complex, traditional resistance and plyometric training alone. *Journal of Musculoskeletal Research*. 2014; 17(2) (pagination):Arte Number: 1450007. ate of Pubaton: 22 Jun 2014.
167. De Villarreal ES, Requena B, Izquierdo M, Gonzalez-Badillo JJ. Enhancing sprint and strength performance: combined versus maximal power, traditional heavy-resistance and plyometric training. *Journal of science and medicine in sport*. 2013; 16(2):146-50.
168. Laird RH, Elmer DJ, Barberio MD, Salom LP, Lee KA, Pascoe DD. Evaluation of Performance Improvements After Either Resistance Training or Sprint Interval-Based Concurrent Training. *Journal of Strength and Conditioning Research*. 2016; 30(11):3057-3065.
169. Fink J, Kikuchi N, Yoshida S, Terada K, Nakazato K. Impact of high versus low fixed loads and non-linear training loads on muscle hypertrophy, strength and force development. *Springerplus*. 2016; 5(1):698.
170. Ignjatovic A., Radovanovic D., Stankovic R., Markovic Z., Kocic J. Influence of resistance training on cardiorespiratory endurance and muscle power and strength in young athletes. *Acta Physiologica Hungarica*. 2011; 98(3):305-312.
171. Kraemer WJ, Ratamess N, Fry AC, Triplett-Mcbride T, Koziris LP, Bauer JA, Lynch JM, Fleck SJ. Influence of resistance training volume and periodization on physiological and performance adaptations in collegiate women tennis players. *American Journal of Sports Medicine*. 2000; 28(5):626-33.

172. Talpey SW, Young WB, Saunders N. Is nine weeks of complex training effective for improving lower body strength, explosive muscle function, sprint and jumping performance? *International Journal of Sports Science and Coaching*. 2016; 11(5):736-745.
173. Berger R. Effect of varied weight training programs on strength. *Research Quarterly. American Association for Health, Physical Education and Recreation*. 1962; 33(2):168-81.
174. Dasteridis G, Pilianidis T, Mantzouranis N. The Effect of Different Strength Training Programs on Young Athletes' Sprint Performance. *Studies in Physical Culture & Tourism*. 2011; 18(2):141-147.
175. Whitley JD, Smith LE. Influence of three different training programs on strength and speed of a limb movement. *Research quarterly*. 1966; 37(1):132-142.
176. Jackson A, Jackson T, Hnatek J, West J. Strength development: Using functional isometrics in an isotonic strength training program. *Research quarterly for exercise and sport*. 1985; 56(3):234-237.
177. Kruszewski M, Kruszewski A, Kruszewski B. Changes in Relative and Absolute Force Measured during Powerlifting After a 4-Week Training using the Weightlifting Method and Stick Isometric (Mixed) Method. *Medicina Sportiva*. 2008; 12(2):41-45.
178. Clutch D, Wilton M, McGown C, Bryce GR. Effect of depth jumps and weight training on leg strength and vertical jump. *Research Quarterly for Exercise & Sport*. 1983; 54(1):5-10.
179. Bell GJ, Petersen SR, Quinney HA, Wenger HA. The effect of velocity-specific strength training on peak torque and anaerobic rowing power. *Journal of sports sciences*. 1989; 7(3):205-214.
180. Rutherford OM, Greig CA, Sargeant AJ, Jones DA. Strength training and power output: transference effects in the human quadriceps muscle. *Journal of sports sciences*. 1986; 4(2):101-107.
181. Doherty TJ, Campagna PD. The effects of periodized velocity-specific resistance training on maximal and sustained force production in women. *Journal of sports sciences*. 1993; 11(1):77-82.
182. Delecluse C, Van Coppenolle HE, Willems EU, Van Leemputte M, Diels R, Goris MA. Influence of high-resistance and high-velocity training on sprint performance. *Medicine and science in sports and exercise*. 1995; 27(8):1203-9.
183. O'Hagan FT, Sale DG, MacDougall JD, Garner SH. Comparative effectiveness of accommodating and weight resistance training modes. *Medicine and science in sports and exercise*. 1995; 27(8):1210-9.
184. Pesta D, Thaler A, Hoppel F, Macek C, Schocke M, Burtscher M. Effects of a 10-week conventional strength training program on lower leg muscle performance in adolescent boys compared to adults. *Journal of sports medicine and physical fitness*. 2014; 54(2):147-53.
185. Spinetti J, Figueiredo T, Assis M, Miranda H, Simão R. Comparison between traditional strength training and complex contrast training on repeated sprint ability and muscle architecture in elite soccer players. *Journal of sports medicine and physical fitness*. 2015; 56(11):1269-78.
186. Voigt M, Klausen K. Changes in muscle strength and speed of an unloaded movement after various training programmes. *European journal of applied physiology and occupational physiology*. 1990; 60(5):370-376.
187. McMaster D, Gill N, McGuigan M, Cronin J. Effects of Complex Strength and Ballistic Training on Maximum Strength, Sprint Ability and Force-Velocity-Power Profiles of Semi-Professional Rugby Union Players. *Journal of Australian Strength and Conditioning*. 2014; 22(1):17-30.
188. Petersen TS. *Effects of slide board training on the lateral movement of college-aged football players*. Eugene, Or.: Microform Publications, University of Oregon; 2000TY: GENGEN; Accession Number: SPHS-673351; Author: Petersen, T. S. ; Language: English; General Notes: Thesis (M.S.)University of Wisconsin, La Crosse, 2000; includes bibliographical references.; Description: 1 microfiche (97 fr.) : negative, ill. ; 11 x 15 cm.; Database Subset: ID; Publication Type: Microforms; Thesis or dissertation; Update Code: 20010401.
189. Ishoi L, Holmich P, Aagaard P, Thorborg K, Bandholm T, Serner A. Effects of the Nordic Hamstring exercise on sprint capacity in male football players: a randomized controlled trial. *Journal of sports sciences*. 2017; 1-10.
190. Housh T.J., Housh D.J., Weir J.P., Weir LL. Effects of unilateral concentric-only dynamic constant external resistance training. *International Journal of Sports Medicine*. 1996; 17(5):338-343.

191. Perez-Gomez J, Olmedillas H, Delgado-Guerra S, Royo IA, Vicente-Rodriguez G, Ortiz RA, et al. Effects of weight lifting training combined with plyometric exercises on physical fitness, body composition, and knee extension velocity during kicking in football. *Applied Physiology, Nutrition and Metabolism*. 2008; 33(3):501-510.
192. Brazell-Roberts JV, Thomas LE. Effects of weight training frequency on the self-concept of college females. *Journal of Strength and Conditioning Research*. 1989; 3(2):40-3.
193. Otto III WH, Coburn JW, Brown LE, Spiering BA. Effects of weightlifting vs. kettlebell training on vertical jump, strength, and body composition. *Journal of Strength and Conditioning Research*. 2012; 26(5):1199-202.
194. Tanimoto M, Sanada K, Yamamoto K, Kawano H, Gando Y, Tabata I, et al. Effects of whole-body low-intensity resistance training with slow movement and tonic force generation on muscular size and strength in young men. *Journal of Strength and Conditioning Research*. 2008; 22(6):1926-1938.
195. Moraes E, Fleck SJ, Dias MR, Simão R. Effects on strength, power, and flexibility in adolescents of nonperiodized vs. daily nonlinear periodized weight training. *Journal of Strength and Conditioning Research*. 2013; 27(12):3310-21.
196. Andersen V, Fimland MS, Kolnes MK, Saeterbakken AH. Elastic bands in combination with free weights in strength training: Neuromuscular effects. *Journal of Strength and Conditioning Research*. 2015; 29(10):2932-40.
197. Fyfe JJ, Bartlett JD, Hanson ED, Stepto NK, Bishop DJ. Endurance training intensity does not mediate interference to maximal lower-body strength gain during short-term concurrent training. *Frontiers in Physiology*. 2016; 7(NOV) (pagination):Arte Number: 487. ate of Pubaton: 03 No 2016.
198. De Villarreal ESS, Izquierdo M, Gonzalez-Badillo JJ. Enhancing jump performance after combined vs. maximal power, heavy-resistance, and plyometric training alone. *Journal of Strength and Conditioning Research*. 2011; 25(12):3274-3281.
199. De Villarreal ES, Suarez-Arrones L, Requena B, Haff GG, Veliz RR. Enhancing performance in professional water polo players: dryland training, in-water training, and combined training. *Journal of Strength and Conditioning Research*. 2015; 29(4):1089-1097.
200. Lyttle AD, Wilson GJ, Ostrowski KJ. Enhancing performance: maximal power versus combined weights and plyometrics training. *Journal of Strength and Conditioning Research*. 1996; 10(3):173-179.
201. Fatouros IG, Jamurtas AZ, Leontsini D, Taxildaris K, Aggelousis N, Kostopoulos N, Buckenmeyer P. Evaluation of plyometric exercise training, weight training, and their combination on vertical jumping performance and leg strength. *Journal of Strength and Conditioning Research*. 2000; 14(4):470-6.
202. Jenkins ND, Housh TJ, Buckner SL, Bergstrom HC, Smith CM, Cochrane KC, Hill EC, Miramonti AA, Schmidt RJ, Johnson GO, Cramer JT. Four weeks of high-versus low-load resistance training to failure on the rate of torque development, electromechanical delay, and contractile twitch properties. *Journal of musculoskeletal & neuronal interactions*. 2016; 16(2):135-144.
203. Maroto-Izquierdo S, García-López D, de Paz JA. Functional and muscle-size effects of flywheel resistance training with eccentric-overload in professional handball players. *Journal of human kinetics*. 2017; 60:133-143.
204. O'Shea K, O'Shea JP. Functional isometrics weight training: Its effects on dynamic and static strength. *Journal of Strength and Conditioning Research*. 1989; 3(2):30-33.
205. Giorgi A, Wilson GJ, Weatherby RP, Murphy AJ. Functional isometric weight training: its effects on the development of muscular function and the endocrine system over an 8-week training period. *Journal of Strength and Conditioning Research*. 1998; 12(1):18-25.
206. Fimland MS, Helgerud J, Gruber M, Leivseth G, Hoff J. Functional maximal strength training induces neural transfer to single-joint tasks. *European journal of applied physiology*. 2009; 107(1):21-29.
207. Oliver JM, Jagim AR, Sanchez AC, Mardock MA, Kelly KA, Meredith HJ, et al. Greater gains in strength and power with intraset rest intervals in hypertrophic training. *Journal of Strength and Conditioning Research*. 2013; 27(11):3116-3131.
208. Loturco I, Pereira LA, Kobal R, Zanetti V, Gil S, Kitamura K, Abad CC, Nakamura FY. Half-squat or jump squat training under optimum power load conditions to counteract power and speed

- decrements in Brazilian elite soccer players during the preseason. *Journal of sports sciences*. 2015; 33(12):1283-92.
209. Ayers JL, DeBeliso M, Sevene TG, Adams KJ. Hang cleans and hang snatches produce similar improvements in female collegiate athletes. *Biology of Sport*. 2016; 33(3):251-256.
210. Dæhlin TE, Haugen OC, Haugerud S, Hollan I, Raastad T, Rønnestad BR. Improvement of ice hockey players' on-ice sprint with combined plyometric and strength training. *International journal of sports physiology and performance*. 2017; 12(7):893-900.
211. Ahmed TA. Improving musculoskeletal fitness and the performance enhancement of basketball skills through neuromuscular training program. *Journal of Human Sport and Exercise*. 2015;10(3):795-804.
212. Neils CM, Udermann BE, Brice GA, Winchester JB, McGuigan MR. Influence of Contraction Velocity in Untrained Individuals Over the Initial Early Phase of Resistance Training. *Journal of Strength and Conditioning Research*. 2005; 19(4):883-887.
213. Franchini E, Branco BM, Agostinho MF, Calmet M, Candau R. Influence of linear and undulating strength periodization on physical fitness, physiological, and performance responses to simulated judo matches. *Journal of Strength and Conditioning Research*. 2015; 29(2):358-67.
214. Hartmann H, Wirth K, Klusemann M, Dalic J, Matuschek C, Schmidtbleicher D. Influence of squatting depth on jumping performance. *Journal of Strength and Conditioning Research*. 2012; 26(12):3243-3261.
215. Otero-Esquina C, de HL, Gonzalo-Skok Ó, Domínguez-Cobo S, Sánchez H. Is strength-training frequency a key factor to develop performance adaptations in young elite soccer players? *European Journal of Sport Science*. 2017; 17(10):1241-1251.
216. Shalfawi SAI, Haugen T, Jakobsen TA, Enoksen E, Tønnessen E. The effect of combined resisted agility and repeated sprint training vs. strength training on female elite soccer players. *Journal of Strength and Conditioning Research*. 2013; 27(11):2966-72.
217. Tasi YJ, Liu GC, Chen CY, Huang C. The effect of different plyometric-squat training on taekwondo power development in the lower extremity. [Accession Number: SPHS-162162; Author: Tasi, Y.J. Author: Liu, G.C. Author: Chen, C.Y. Author: Huang, C. ; Corporate Author: International Society of Biomechanics in Sports; Conference: International Symposium on Biomechanics in Sports (17th : 1999 : Perth, Western Australia).; No. of Pages: 4; Language: English; Parent Item: SPHS-163221; References: 9; Database Subset: ID; Publication Type: Book Analytic; Update Code: 20000301; SIRC Article No.: S-162162]. Reproduced ; 1999.
218. Lopes CR, Aoki MS, Crisp AH, de Mattos RS, Lins MA, da Mota GR, Schoenfeld BJ, Marchetti PH. The effect of different resistance training load schemes on strength and body composition in trained men. *Journal of human kinetics*. 2017; 58:177-186.
219. Hawkins SB, Doyle TL, McGuigan MR. The effect of different training programs on eccentric energy utilization in college-aged males. *Journal of Strength and Conditioning Research*. 2009; 23(7):1996-2002.
220. Lee A, Craig BW, Lucas J, Pohlman R, Stelling H. The effect of endurance training, weight training and a combination of endurance and weight training upon the blood lipid profile of young male subjects. *Journal of Strength and Conditioning Research*. 1990; 4(3):68-75.
221. Alemdaroglu U, Dundar U, Koklu Y, Asci A, Findikoglu G. The effect of exercise order incorporating plyometric and resistance training on isokinetic leg strength and vertical jump performance: A comparative study. *Isokinetics and Exercise Science*. 2013; 21(3):211-217.
222. Hermassi S, Chelly MS, Fathloun M, Shephard RJ. The effect of heavy-vs. moderate-load training on the development of strength, power, and throwing ball velocity in male handball players. *Journal of Strength and Conditioning Research*. 2010; 24(9):2408-18.
223. Moir G, Sanders R, Button C, Glaister M. The effect of periodized resistance training on accelerative sprint performance. *Sports Biomechanics*. 2007; 6(3):285-300.
224. Arazi H, Khanmohammadi A, Asadi A, Haff GG. The effect of resistance training set configuration on strength, power, and hormonal adaptation in female volleyball players. *Applied physiology, nutrition, and metabolism*. 2018;43(2):154-64.

225. Wenzel RR, Perfetto EM. The effect of speed versus non-speed training in power development. *Journal of Applied Sport Science Research*. 1992; 6(2):82-87.
226. Hammami M, Negra Y, Shephard RJ, Chelly MS. The effect of standard strength vs. contrast strength training on the development of sprint, agility, repeated change of direction, and jump in junior male soccer players. *Journal of Strength and Conditioning Research*. 2017; 31(4):901-12.
227. Beattie K, Carson BP, Lyons M, Rossiter A, Kenny IC. The Effect of Strength Training on Performance Indicators in Distance Runners. *Journal of Strength and Conditioning Research*. 2017; 31(1):9-23.
228. Jakobsen MD, Sundstrup E, Randers MB, Kjær M, Andersen LL, Krstrup P, Aagaard P. The effect of strength training, recreational soccer and running exercise on stretch–shortening cycle muscle performance during countermovement jumping. *Human movement science*. 2012; 31(4):970-86.
229. Young WB, Bilby GE. The effect of voluntary effort to influence speed of contraction on strength, muscular power, and hypertrophy development. *Journal of Strength & Conditioning Research*. 1993; 7(3):172-178.
230. Ostrowski KJ, Wilson GJ, Weatherby R, Murphy PW, Lyttle AD. The effect of weight training volume on hormonal output and muscular size and function. *Journal of Strength and Conditioning Research*. 1997; 11:148-54.
231. Holcomb WR, Lander JE, Rutland RM, Wilson GD. The effectiveness of a modified plyometric program on power and the vertical jump. *Journal of Strength and Conditioning Research*. 1996; 10(2):89-92.
232. Rhea MR, Peterson MD, Lunt KT, Ayllon FN. The effectiveness of resisted jump training on the VertiMax in high school athletes. *Journal of Strength and Conditioning Research*. 2008; 22(3):731-734.
233. Shoepe TC, Ramirez DA, Rovetti RJ, Kohler DR, Almstedt HC. The Effects of 24 weeks of Resistance Training with Simultaneous Elastic and Free Weight Loading on Muscular Performance of Novice Lifters. *Journal of Human Kinetics*. 2011; 29:93-106.
234. Alcaraz PE, Elvira JLL, Palao JM. Kinematic, strength, and stiffness adaptations after a short-term sled towing training in athletes. *Scandinavian Journal of Medicine and Science in Sports*. 2014; 24(2):279-290.
235. Caruso JF, Coday MA, Ramsey CA, Griswold SH, Polanski DW, Drummond JL, et al. Leg and calf press training modes and their impact on jump performance adaptations. *Journal of strength and conditioning research / National Strength & Conditioning Association*. 2008; 22(3):766-772.
236. Torres-Torrel J, Rodriguez-Rosell D, Gonzalez-Badillo JJ. Light-load maximal lifting velocity full squat training program improves important physical and skill characteristics in futsal players. *Journal of sports sciences*. 2017; 35(10):967-975.
237. Keiner M, Sander A, Wirth K, Schmidtbleicher D. Long-term strength training effects on change-of-direction sprint performance. *Journal of Strength and Conditioning Research*. 2014; 28(1):223-31.
238. Hong J, Smith JD, Ross CN, Lee S. Low volume progressive single set of resistance training is as effective as high volume multiple sets of resistance protocol on muscle strength and power. *International Journal of Applied Sports Sciences*. 2015; 27(1):33-42.
239. Kikuchi N, Nakazato K. Low-load bench press and push-up induce similar muscle hypertrophy and strength gain. *Journal of Exercise Science & Fitness*. 2017; 15(1):37-42.
240. Marx JO, Ratamess NA, Nindl BC, Gotshalk LA, Volek JS, Dohi K, et al. Low-volume circuit versus high-volume periodized resistance training in women. *Medicine and science in sports and exercise*. 2001; 33(4):635-643.
241. Tillin N, Folland J. Maximal and explosive strength training elicit distinct neuromuscular adaptations, specific to the training stimulus. *European journal of applied physiology*. 2014; 114(2):365-374.
242. Cherif M, Chtourou H, Souissi N, Aouidet A, Chamari K. Maximal power training induced different improvement in throwing velocity and muscle strength according to playing positions in elite male handball players. *Biology of Sport*. 2016; 33(4):393-398.

243. Coyne JO, Tran TT, Secomb JL, Lundgren LE, Farley OR, Newton RU, et al. Maximal Strength Training Improves Surfboard Sprint and Endurance Paddling Performance in Competitive and Recreational Surfers. *Journal of Strength and Conditioning Research*. 2017; 31(1):244-253.
244. Heggelund J, Fimland MS, Helgerud J, Hoff J. Maximal strength training improves work economy, rate of force development and maximal strength more than conventional strength training. *European journal of applied physiology*. 2013; 113(6):1565-73.
245. Smilios I, Sotiropoulos K, Christou M, Douda H, Spaias A, Tokmakidis SP. Maximum power training load determination and its effects on load-power relationship, maximum strength, and vertical jump performance. *Journal of Strength and Conditioning Research*. 2013; 27(5):1223-1233.
246. Newton RU, Hakkinen K, Hakkinen A, McCormick M, Volek J, Kraemer WJ. Mixed-methods resistance training increases power and strength of young and older men. *Medicine and science in sports and exercise*. 2002; 34(8):1367-1375.
247. Zourdos MC, Jo E, Khamoui AV, Lee SR, Park BS, Ormsbee MJ, et al. Modified Daily Undulating Periodization Model Produces Greater Performance Than a Traditional Configuration in Powerlifters. *Journal of Strength and Conditioning Research*. 2016; 30(3):784-791.
248. Vissing K, Brink M, Lønbro S, Sørensen H, Overgaard K, Danborg K, Mortensen J, Elstrøm OL, Rosenhøj N, Ringgaard S, Andersen JL. Muscle adaptations to plyometric vs. resistance training in untrained young men. *Journal of Strength and Conditioning Research*. 2008; 22(6):1799-810.
249. Toumi H, Best TM, Martin A, Poumarat G. Muscle plasticity after weight and combined (weight + jump) training. *Medicine and Science in Sports and Exercise*. 2004; 36(9):1580-1588.
250. Stasinaki AN, Gloumis G, Spengos K, Blazeovich AJ, Zaras N., Georgiadis G., et al. Muscle Strength, Power, and Morphologic Adaptations After 6 Weeks of Compound vs. Complex Training in Healthy Men. *Journal of Strength and Conditioning Research*. 2015; 29(9):2559-2569.
251. Campos GER, Luecke TJ, Wendeln HK, Toma K, Hagerman FC, Murray TF, et al. Muscular adaptations in response to three different resistance-training regimens: Specificity of repetition maximum training zones. *European journal of applied physiology*. 2002; 88(1-2):50-60.
252. Vikne H, Refsnes PE, Ekmark M, Medbo JI, Gundersen V, Gundersen K. Muscular performance after concentric and eccentric exercise in trained men. *Medicine and science in sports and exercise*. 2006; 38(10):1770-1781.
253. Judge L, Moreau C, Burke J. Neural adaptations with sport-specific resistance training in highly skilled athletes. *Journal of Sports Sciences*. 2003; 21(5):419-27.
254. Häkkinen K, Alen M, Kraemer WJ, Gorostiaga E, Izquierdo M, Rusko H, Mikkola J, Häkkinen A, Valkeinen H, Kaarakainen E, Romu S. Neuromuscular adaptations during concurrent strength and endurance training versus strength training. *European journal of applied physiology*. 2003; 89(1):42-52.
255. Mikkola J, Rusko H, Izquierdo M, Gorostiaga EM, Hakkinen K. Neuromuscular and cardiovascular adaptations during concurrent strength and endurance training in untrained men. *International Journal of Sports Medicine*. 2012; 33(9):702-710.
256. Romero-Arenas S, Ruiz R, Vera-Ibanez A, Colomer-Poveda D, Guadalupe-Grau A, Marquez G. Neuromuscular and Cardiovascular Adaptations in Response to High-Intensity Interval Power Training. *Journal of Strength and Conditioning Research*. 2018; 32(1):130-138.
257. Ullrich B, Pelzer T, Oliveira S, Pfeiffer M. Neuromuscular responses to short-term resistance training with traditional and daily undulating periodization in adolescent elite judoka. *Journal of Strength and Conditioning Research*. 2016; 30(8):2083-99.
258. Myer GD, Ford KR, Palumbo JP, Hewett TE. Neuromuscular training improves performance and lower-extremity biomechanics in female athletes. *Journal of Strength and Conditioning Research*. 2005; 19(1):51-60.
259. Arabatzis F, Kellis E. Olympic weightlifting training causes different knee muscle-coactivation adaptations compared with traditional weight training. *Journal of Strength and Conditioning Research*. 2012; 26(8):2192-2201.
260. Jolley RI, Goodwin JE, Cleather DJ. Peak Power Output in the Bench Pull Is Maximized After Four Weeks of Specific Power Training. *Journal of Strength and Conditioning Research*. 2016; 30(4):966-972.

261. Baker D, Wilson G, Carlyon R. Periodization: the effect on strength of manipulating volume and intensity. *Journal of Strength and Conditioning Research (Allen Press Publishing Services Inc.)*. 1994; 8(4):235-242.
262. Groves BR, Gayle RC. Physiological changes in male basketball players in year-round strength training. *Journal of Strength and Conditioning Research*. 1993; 7(1):30-33.
263. Kraemer WJ, Hakkinen K, Triplett-McBride NT, Fry AC, Koziris LP, Ratamess NA, et al. Physiological changes with periodized resistance training in women tennis players. *Medicine and science in sports and exercise*. 2003; 35(1):157-168.
264. Cormie P, McCaulley GO, McBride JM. Power versus strength-power jump squat training: influence on the load-power relationship. *Medicine and science in sports and exercise*. 2007; 39(6):996-1003.
265. Voelzke M, Stutzig N, Thorhauer HA, Granacher U. Promoting lower extremity strength in elite volleyball players: Effects of two combined training methods. *Journal of Science and Medicine in Sport*. 2012; 15(5):457-462.
266. Munn J, Herbert RD, Hancock MJ, Gandevia SC. Resistance training for strength: effect of number of sets and contraction speed. *Medicine and science in sports and exercise*. 2005; 37(9):1622-1626.
267. Mangine GT, Hoffman JR, Wang R, Gonzalez AM, Townsend JR, Wells AJ, Jajtner AR, Beyer KS, Boone CH, Miramonti AA, LaMonica MB. Resistance training intensity and volume affect changes in rate of force development in resistance-trained men. *European journal of applied physiology*. 2016; 116(11):2367-74.
268. Muehlbauer T, Gollhofer A, Granacher U. Sex-related effects in strength training during adolescence: A pilot study. *Perceptual and motor skills*. 2012; 115(3):953-968.
269. Winwood PW, Buckley JJ. Short Term Effects of Resistance Training Modalities on Performance Measures in Male Adolescents. *Journal of Strength and Conditioning Research*. 2019; 33(3):641-50.
270. Herrero AJ, Martin J, Martin T, Abadia O, Fernandez B, GarciaLopez D. Short-term effect of strength training with and without superimposed electrical stimulation on muscle strength and anaerobic performance. A randomized controlled trial. Part I. *Journal of Strength and Conditioning Research*. 2010; 24(6):1609-1615.
271. Alves JVM, Rebelo AN, Abrantes C, Sampaio J. Short-term effects of complex and contrast training in soccer players' vertical jump, sprint, and agility abilities. *Journal of Strength and Conditioning Research*. 2010; 24(4):936-941.
272. Cavaco B, Sousa N, Dos Reis VM, Garrido N, Saavedra F, Mendes R, Vilaça-Alves J. Short-term effects of complex training on agility with the ball, speed, efficiency of crossing and shooting in youth soccer players. *Journal of human kinetics*. 2014; 43:105.
273. Ronnestad BR, Kvamme NH, Sunde A, Raastad T. Short-term effects of strength and plyometric training on sprint and jump performance in professional soccer players. *Journal of Strength and Conditioning Research*. 2008; 22(3):773-80.
274. Tricoli V, Lamas L, Carnevale R, Ugrinowitsch C. Short-term effects on lower-body functional power development: weightlifting vs. vertical jump training programs. *Journal of Strength and Conditioning Research*. 2005; 19(2):433-7.
275. Sanborn K, Boros R, Hrubby J, Schilling B, O'Bryant HS, Johnson RL, Hoke T, Stone ME, Stone MH. Short-term performance effects of weight training with multiple sets not to failure vs. a single set to failure in women. *Journal of Strength and Conditioning Research*. 2000; 14(3):328-31.
276. Kijowksi KN, Capps CR, Goodman CL, Erickson TM, Knorr DP, Triplett NT, Awelewa OO, McBride JM. Short-term resistance and plyometric training improves eccentric phase kinetics in jumping. *Journal of Strength and Conditioning Research*. 2015; 29(8):2186-96.
277. Farup J, Sørensen H, Kjølhede T. Similar changes in muscle fiber phenotype with differentiated consequences for rate of force development: endurance versus resistance training. *Human movement science*. 2014; 34:109-19.
278. Alcaraz PE, Perez-Gomez J, Chavarrias M, Blazeovich AJ. Similarity in adaptations to high-resistance circuit vs. traditional strength training in resistance-trained men. *Journal of Strength and Conditioning Research*. 2011; 25(9):2519-2527.

279. Palmer T, Uhl TL, Howell D, Hewett TE, Viele K, Mattacola CG. Sport-specific training targeting the proximal segments and throwing velocity in collegiate throwing athletes. *Journal of Athletic Training*. 2015; 50(6):567-77.
280. Knox A, Sculthorpe N, Baker JS, Grace F. Strength adaptation to squat exercise is different between Caucasian and South Asian novice exercisers. *Research in Sports Medicine*. 2017; 25(3):373-83.
281. Marshall P.W.M., McEwen M., Robbins DW. Strength and neuromuscular adaptation following one, four, and eight sets of high intensity resistance exercise in trained males. *European journal of applied physiology*. 2011; 111(12):3007-3016.
282. Painter KB, Haff GG, Ramsey MW, McBride J, Triplett T., Sands WA, et al. Strength gains: Block versus daily undulating periodization weight training among track and field athletes. *International Journal of Sports Physiology and Performance*. 2012; 7(2):161-169.
283. Delecluse C, Roelants M, Verschueren S. Strength increase after whole-body vibration compared with resistance training. *Medicine & Science in Sports & Exercise*. 2003; 35(6):1033-1041.
284. Manolopoulos E, Papadopoulos C, Salonikidis K, Katartzi E, Poluha S. Strength training effects on physical conditioning and instep kick kinematics in young amateur soccer players during preseason. *Perceptual and motor skills*. 2004; 99(2):701-710.
285. Folland JP, Hawker K, Leach B, Little T, Jones DA. Strength training: Isometric training at a range of joint angles versus dynamic training. *Journal of sports sciences*. 2005; 23(8):817-824.
286. Stone WJ, Coulter SP. Strength/endurance effects from three resistance training protocols with women. *Journal of Strength and Conditioning Research*. 1994; 8(4):231-4.
287. Blakey JB, Southard D. The combined effects of weight training and plyometrics on dynamic leg strength and leg power. *Journal of Applied Sport Science Research*. 1987; 1(1):14-16.
288. Parnow A, Derakhshandeh S, Hosseini A. The Effect of 4-week Difference Training Methods on Some Fitness Variables in Youth Handball Players. *International Journal of Applied Exercise Physiology*. 2016; 5(3).
289. Mills JD, Taunton JE, Mills WA. The effect of a 10-week training regimen on lumbo-pelvic stability and athletic performance in female athletes: A randomized-controlled trial. *Physical Therapy in Sport*. 2005; 6(2):60-66.
290. Kotzamanidis C, Chatzopoulos DI, Michailidis C, Papaiakevou G, Patikas DI. The effect of a combined high-intensity strength and speed training program on the running and jumping ability of soccer players. *Journal of Strength and Conditioning Research*. 2005; 19(2):369-75.
291. Kubo K, Morimoto M, Komuro T, Yata H, Tsunoda N, Kanehisa H, Fukunaga T. Effects of plyometric and weight training on muscle-tendon complex and jump performance. *Medicine and science in sports and exercise*. 2007; 39(10):1801-10.
292. Damasceno MV, Lima-Silva AE, Pasqua LA, Tricoli V, Duarte M, Bishop DJ, et al. Effects of resistance training on neuromuscular characteristics and pacing during 10-km running time trial. *European journal of applied physiology*. 2015; 115(7):1513-1522.
293. Christou M, Smilios I, Sotiropoulos K, Volaklis K, Pilianidis T, Tokmakidis SP. Effects of resistance training on the physical capacities of adolescent soccer players. *Journal of Strength and Conditioning Research*. 2006; 20(4):783-791.
294. Glowacki SP, Martin SE, Maurer A, Baek W, Green JS, Crouse SF. Effects of resistance, endurance, and concurrent exercise on training outcomes in men. *Medicine and science in sports and exercise*. 2004; 36(12):2119-2127.
295. Kramer JB, Stone MH, O'Bryant HS, Conley MS, Johnson RL, Nieman DC, Honeycutt DR, Hoke TP. Effects of single vs. multiple sets of weight training: impact of volume, intensity, and variation. *Journal of Strength and Conditioning Research*. 1997; 11:143-7.