

AUTHOR(S):

TITLE:

YEAR:

Publisher citation:

OpenAIR citation:

Publisher copyright statement:

This is the _____ version of an article originally published by _____
in _____
(ISSN _____; eISSN _____).

OpenAIR takedown statement:

Section 6 of the "Repository policy for OpenAIR @ RGU" (available from <http://www.rgu.ac.uk/staff-and-current-students/library/library-policies/repository-policies>) provides guidance on the criteria under which RGU will consider withdrawing material from OpenAIR. If you believe that this item is subject to any of these criteria, or for any other reason should not be held on OpenAIR, then please contact openair-help@rgu.ac.uk with the details of the item and the nature of your complaint.

This publication is distributed under a CC _____ license.


RESEARCH ARTICLE

Perspectives of healthcare professionals in Qatar on causes of medication errors: A mixed methods study of safety culture

Derek Stewart^{1*}, Binny Thomas^{1,2}, Katie MacLure¹, Abdulrouf Pallivalapila², Wessam El Kassem², Ahmed Awaisu³, James S. McLay⁴, Kerry Wilbur³, Kyle Wilby³, Cristin Ryan⁵, Andrea Dijkstra⁶, Rajvir Singh⁶, Moza Al Hail²

1 School of Pharmacy and Life Sciences, Robert Gordon University, Aberdeen, United Kingdom, **2** Women's Hospital, Hamad Medical Corporation, Doha, Qatar, **3** College of Pharmacy, Qatar University, Doha, Qatar, **4** Institute of Medical Sciences, University of Aberdeen, Aberdeen, United Kingdom, **5** School of Pharmacy, Royal College of Surgeons in Ireland, Dublin, Ireland, Dublin 2, **6** Hamad Medical Corporation, Doha, Qatar

* d.stewart@rgu.ac.uk



Abstract

OPEN ACCESS

Citation: Stewart D, Thomas B, MacLure K, Pallivalapila A, El Kassem W, Awaisu A, et al. (2018) Perspectives of healthcare professionals in Qatar on causes of medication errors: A mixed methods study of safety culture. PLoS ONE 13(9): e0204801. <https://doi.org/10.1371/journal.pone.0204801>

Editor: Susan Hrisos, Newcastle University, UNITED KINGDOM

Received: June 30, 2017

Accepted: September 15, 2018

Published: September 28, 2018

Copyright: © 2018 Stewart et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All quantitative data derived from the cross-sectional survey are given in the paper. For the focus group discussions, data are qualitative in nature and cannot be made publicly available due to ethical as they contain potentially identifiable information. Researchers who meet the criteria for access to confidential data may contact the Robert Gordon University Research Ethics Sub-Committee, Research-Integrity@rgu.ac.uk.

Background

There is a lack of robust, rigorous mixed methods studies of patient safety culture generally and notably those which incorporate behavioural theories of change. The study aimed to quantify and explain key aspects of patient safety culture which were of most concern to healthcare professionals in Qatar.

Methods

A sequential explanatory mixed methods design of a cross-sectional survey followed by focus groups in Hamad Medical Corporation, Qatar. All doctors, nurses and pharmacists were invited to complete the Hospital Survey on Patient Safety Culture (HSOPS). Respondents expressing interest in focus group participation were sampled purposively, and discussions based on survey findings using the Theoretical Domains Framework (TDF) to explain behavioural determinants.

Results

One thousand, six hundred and four questionnaires were received (67.9% nurses, 13.3% doctors, 12.9% pharmacists). HSOPS composites with the lowest levels of positive responses were non-punitive response to errors (24.0% positive) and staffing (36.2%). Specific TDF determinants potentially associated with these composites were social/professional role and identity, emotions, and environmental context and resources. Thematic analysis identified issues of doctors relying on pharmacists to correct their errors and being reluctant to alter the prescribing of fellow doctors. There was a lack of recognition of nurses' roles and frequent policy non-adherence. Stress, workload and lack of staff at key times were perceived to be major contributors to errors.

Funding: This publication was made possible by NPRP grant NPRP 7-388-3-095 from Qatar National Research Fund (a member of Qatar Foundation). The statements made herein are solely the responsibility of the authors.

Competing interests: The authors have declared that no competing interests exist.

Conclusions

This study has quantified areas of concern relating to patient safety culture in Qatar and suggested important behavioural determinants. Rather than focusing on changing behaviour at the individual practitioner level, action may be required at the organisational strategic level to review policies, structures (including resource allocation and distribution) and processes which aim to promote patient safety culture.

Introduction

Promoting patient safety in healthcare settings is a global challenge, with an estimated one in ten patients being harmed whilst receiving care [1]. In an effort to raise awareness of key concepts and strategies in patient safety, the World Health Organization (WHO) published 'Medication Without Harm, WHO Global Patient Safety Challenge' in March 2017 [2,3]. The challenge calls for action to reduce patient harm which occurs as a result of unsafe medication practices and medication errors [2,3]. The goal is to 'gain worldwide commitment and action to reduce severe, avoidable medication-related harm by 50% in the next five years, specifically by addressing harm resulting from errors or unsafe practices due to weaknesses in health systems'. Accumulation of evidence confirms that healthcare professionals often prescribe, dispense and administer medication in ways and circumstances that may increase the risk of patient harm [4–8].

Whilst it is noted that the magnitude and nature of medication harm may differ between countries, globally the cost associated with medication errors has been estimated at US\$ 42 billion annually [2,3]. The most commonly cited and accepted definition of the term 'medication error' is that of the National Coordinating Council for Medication Error Reporting and Prevention (NCCMERP) in the United States (US), 'any preventable event that may cause or lead to inappropriate medication use or patient harm, while the medication is in the control of the health care professional, patient, or consumer' [9]. Most research literature focuses on errors relating to prescribing, administration and dispensing, with evidence that causation is often complex and multifactorial. Systematic reviews have focused on causes of medication errors in different patient populations and settings [10–13]. Common to all systematic reviews is the relatively poor research methodologies reported in most of the primary literature, a lack of behavioural theory and organisational culture in study design. Furthermore, very few studies employed a mixed methods approach to allow quantification and in-depth description and explanation of contributory factors.

Behavioural theories may be used to provide explanation hence providing a robust and rigorous foundation for development of behaviour change interventions. The United Kingdom (UK) Medical Research Council (MRC) framework, 'Developing and implementing complex interventions' highlights the importance of considering theory, noting that interventions grounded in theory are more likely to be effective than those developed empirically or pragmatically' [14]. The Theoretical Domains Framework (TDF) is being used increasingly within health-related research to provide insight into influences on behaviour. TDF derives from 33 psychological theories and 128 theoretical constructs organised into 14 domains of behavioural determinants, as described in Table 1 [15].

It is apparent that there is also a need to focus attention on organisational safety culture. The 'Study Group on Human Factors' defines organisational safety culture as, 'the product of individual and group values, attitudes, perceptions, competencies, and patterns of behaviour

Table 1. Description of TDF domains (adapted from Cain et al. [15]).

TDF Domains	Description
Knowledge	An awareness of the existence of something
Skills	An ability or proficiency acquired through practice
Social/Professional Role and Identity	A coherent set of behaviours and displayed personal qualities of an individual in a social or work setting
Beliefs about Capabilities	Acceptance of the truth, reality, or validity about an ability, talent, or facility that a person can put to constructive use
Optimism	The confidence that things will happen for the best or that desired goals will be attained
Beliefs about Consequences	Acceptance of the truth, reality, or validity about outcomes of a behaviour in a given situation
Reinforcement	Increasing the probability of a response by arranging a dependent relationship, or contingency, between the response and a given stimulus
Intentions	A conscious decision to perform a behaviour or a resolve to act in a certain way
Goals	Mental representations of outcomes or end states that an individual wants to achieve
Memory, Attention and Decision Processes	The ability to retain information, focus selectively on aspects of the environment and choose between two or more alternatives
Environmental Context and Resources	Any circumstance of a person's situation or environment that discourages or encourages the development of skills and abilities, independence, social competence, and adaptive behaviour
Social Influences	Those interpersonal processes that can cause individuals to change their thoughts, feelings, or behaviours
Emotion	A complex reaction pattern, involving experiential, behavioural, and physiological elements, by which the individual attempts to deal with a personally significant matter or event
Behavioural Regulation or measured actions	Anything aimed at managing or changing objectively observed

<https://doi.org/10.1371/journal.pone.0204801.t001>

that determine the commitment to, and the style and proficiency of, an organization's health and safety management [16]. While two systematic reviews have explored interventions to promote safety culture in hospitals in general and acute hospitals specifically, medication safety was not a feature of any primary research [17,18].

In an attempt to promote and standardise the measurement of organisational safety culture, the US Agency for Healthcare Research and Quality (AHRQ) and Medical Errors Workgroup of the Quality Interagency Coordination Task Force (QuIC) sponsored the development of the Hospital Survey on Patient Safety Culture (HSOPS) [19]. Items are clustered within 12 composites as presented in Table 2.

Research on medication errors within the Middle East has historically been reported to be of poor quality [20]. Recently, Elmontsri et al. conducted a systematic review to explore the status of patient safety culture in Arab countries based on the findings of the HSOPS [21]. Data from 18 studies across seven countries (excluding Qatar) were included, identifying that composites relating to non-punitive response to error to be infrequently practised in their organisation, that staffing levels were often inadequate and that communication needed to be more open. The authors concluded that further research is warranted to provide explanation of these findings and to identify potential interventions to enhance culture and patient safety.

The aim of the present study was to quantify and explain key aspects of patient safety culture which were of most concern to health professionals in Qatar.

Table 2. HSOPS composites and definitions [19].

Composite	Definition: The extent to which. . .
Communication openness	staff freely speak up if they see something that may negatively affect a patient and feel free to question those with more authority
Feedback and communication about error	staff are informed about errors that happen, are given feedback about changes implemented, and discuss ways to prevent errors
Frequency of events reported	mistakes of the following types are reported: mistakes caught and corrected before affecting the patient; mistakes with no potential to harm the patient; and mistakes that could harm the patient but do not
Handoffs and transitions	important patient care information is transferred across hospital units and during shift changes
Management support for patient safety	hospital management provides a work climate that promotes patient safety and shows that patient safety top priority
Non-punitive response to error	staff feel that their mistakes and event reports are not held against them and that mistakes are not kept in their personnel file
Organisational learning—continuous improvement	mistakes have led to positive changes and changes evaluated for effectiveness
Overall perceptions of patient safety	procedures and systems are good at preventing errors and there is a lack of patient safety problems
Staffing	there are enough staff to handle the workload and work hours are appropriate to provide the best care for patients
Supervisor/manager expectations and actions promoting patient safety	supervisors/managers consider staff suggestions for improving patient safety, praise staff for following patient safety procedures, and do not overlook patient safety problems
Teamwork across units	hospital units cooperate and coordinate with one another to provide the best care for patients
Teamwork within units	staff support each other, treat each other with respect, and work together as a team

<https://doi.org/10.1371/journal.pone.0204801.t002>

Methods

Study design

A sequential explanatory mixed methods design was employed, with a cross-sectional survey followed by focus groups in samples of questionnaire respondents to provide further depth and explanation of survey findings [22,23].

Setting

The research was conducted within Hamad Medical Corporation (HMC), the main provider of secondary and tertiary healthcare in Qatar.

Cross-sectional survey

The first phase of the research was a cross-sectional survey.

Questionnaire development

The questionnaire was adapted from AHRQ HSOPS with items presented as 5-point Likert type scales; personal and practice demographic items were added. The common language of care delivery at HMC is English thus translation into other languages (e.g. Arabic) was not required. The questionnaire was piloted in a convenience sample of 100 healthcare professionals. Test-retest reliability was assessed in pilot respondents by requesting that the questionnaire be completed on a second occasion within an interval of two weeks. A high level of test-retest

reliability was achieved ($p < 0.001$, Cohen's kappa) for all Likert statements. The final questionnaire was formatted in Snap 10 Professional (software for web and email questionnaire design, publication, data entry and analysis). On completion of the questionnaire, respondents were invited to participate in focus groups to discuss responses in more detail.

Recruitment

All doctors, nurses and pharmacists working within HMC were eligible to participate, with no exclusions. Three hundred and sixty responses were required to give a margin of error of 5% with 95% confidence intervals [24]. Online participation was encouraged via HMC web alerts and promotional posters; paper-based questionnaires were distributed to all doctors, nurses and pharmacists. Data were collected from mid-January 2016 to mid-April 2016.

Data analysis

Anonymised online submissions were imported into Snap before direct export to SPSS version 21.0 and data cleaned prior to analysis. Paper-based questionnaire data were entered manually using the survey link. Descriptive statistics were used to describe respondents' demographic and professional characteristics as well as their responses to individual HSOPS safety culture items. For each composite, a score was determined in line with the recommendations of the AHRQ task force [19]. This involved firstly calculating the percentage of positive responses to each item. A positive response may reflect agreement with a positively phrased statement, or may reflect disagreement with a negatively phrased statement. Therefore, for positively worded items the percentage of positive responses is the proportion of 'agree' and 'strongly agree' responses to the item, and for negatively worded items the percentage of positive responses is the proportion of disagree and strongly disagree responses to the item. Higher percentages (%) therefore reflect more positive responses to HSOPS safety culture items as defined in Table 2. The composite score was then expressed as the mean of the positive responses. For those composites with overall positive responses of $< 50\%$, Chi-square was used to determine any statistically significant associations with demographic variables.

Focus groups

To clarify, explore and explain issues identified in the survey phase, a qualitative approach was employed.

Sampling and recruitment

Respondents of the survey who expressed interest in participating in the focus groups were sampled purposively to represent a range of professions, hospitals and number of years of experience.

Data generation

The focus group topic guide was developed following analysis of the questionnaire data, with reference to the TDF [15]. Initial discussions were based around views and experiences on safety culture in relation to causes of medication errors. This was followed by targeted discussions around patient cases of errors illustrating errors in prescribing, dispensing and administration in diverse patient groups which led to significant harm. TDF domains were used as prompts in relation to potential causes of errors. Focus groups were moderated by two experienced qualitative researchers, with informed consent obtained from each participant at the outset. Discussions were audio-recorded (with permission), transcribed verbatim and checked

for transcribing reliability. A clear audit trail was maintained which documented details of data gathering to promote dependability [25]. Sampling and recruitment continued to the point of data saturation, when no new themes emerged from data analysis [26]. Focus groups were conducted between mid-May 2016 and mid-June 2016.

Data analysis

Data analysis followed the Framework Approach, using TDF domains deductively for to generate a coding framework [27]. Two researchers independently read each focus group transcript repeatedly to ensure familiarity, then coded text to one or more TDF domains. Any disagreements were resolved by discussion which involved a third researcher if needed.

Ethics

The study received ethical approval from Hamad Medical Corporation, Medical Research Center Qatar, Qatar University Institutional Review Board and Robert Gordon University Research Ethics Sub-Committee. Return of the questionnaire was taken as an indication of informed consent; written informed consent was obtained from all focus group participants.

Results

Cross-sectional survey

Respondents' demographics and professional characteristics. One thousand, six hundred and four completed questionnaires were received, with most (67.9%) from nurses followed by doctors (13.3%) and pharmacists (12.9%). Around three quarters (70.9%) were female, <40 years (76.0%) and almost half (48.1%) with more than 10 years of experience as healthcare professionals. Respondents had varying involvement with medicines-related processes as follows: prescribing medicines (15.1%); administering (61.1%); preparation and dispensing (25.9%); and monitoring (42.0%) (Table 3).

Patient safety culture items. Positive responses to the HSOPS composites and items are given in Table 4. Composites with the lowest levels of mean positive responses were: non-punitive response to errors (24.0%); staffing (36.2%); communication openness (50.5%); handoffs and transitions (53.1%); and supervisor/manager expectations and actions promoting patient safety (56.5%). Composites with the highest levels of positive responses were: organisational learning–continuous improvement (85.5%); team working within unit (82.1%); and management support for patient safety (75.4%). For the two composites with mean positive responses of <50%, Chi-square was used to determine the associations between percentage positive responses and demographics/professional characteristics.

Non-punitive response to errors—all individual items contributing to this HSOPS composite attracted a low level of positive response, this was particularly the case for items relating to staff concerns over errors being kept in their personnel files (26.2%), and the perception that errors counted against them (14.6%). There were highly statistically significant associations with mean composite agreement and gender (females most positive, X^2 (1, $N = 1547$) = 8.23, $p < 0.005$), age (older most positive, X^2 (4, $N = 1555$) = 11.62, $p < 0.05$) and experience as a healthcare professional (the most experienced being most positive, X^2 (5, $N = 1536$) = 18.42, $p < 0.005$).

Staffing—while all responses attracted a low level of positive response, this was particularly the case for work pressures and speed of work (23.5%). There were highly statistically significant associations with mean composite agreement and healthcare professions (doctors most positive and pharmacists least, X^2 (2, $N = 1494$) = 42.06, $p < 0.001$), age (youngest least and

Table 3. Respondents' demographic and professional characteristics (N = 1604).

Characteristic	Percentage	Frequency, n
<i>Current role in the hospital</i>		
Clinical nurse educator	0.7	12
Clinical pharmacist	2.8	45
Consultant physician	5.4	86
Head/Charge/Specialist nurse	17.1	275
Nurse	50.0	802
Pharmacist	8.9	143
Pharmacy Director/Supervisor/Specialist	1.2	19
Resident Physician	3.5	56
Specialist Physician	4.5	72
Other	5.0	80
Missing	0.9	14
<i>Age (years)</i>		
≤29	24.2	392
30–39	41.8	670
40–49	21.5	345
50–59	9.5	153
≥60	1.6	25
Missing	1.7	19
<i>Gender</i>		
Male	27.6	442
Female	70.9	1137
Missing	1.6	25
<i>Country of receiving entry-to-practice degree</i>		
India	42.7	685
Philippines	17.6	283
Egypt	9.3	149
Qatar	9.2	148
Jordan	4.8	77
Other	14.5	231
Missing	1.9	31
<i>Experience as healthcare professional in hospital (years)</i>		
<1	1.6	25
1–5	19.1	306
6–10	29.4	471
11–15	21.4	343
16–20	12.0	193
>20	14.7	235
Missing	1.9	31
<i>Experience as healthcare professional in Qatar (years)</i>		
<1	8.5	136
1–5	40.3	647
6–10	21.8	350
11–15	16.5	264
16–20	5.1	82
>20	6.7	108
Missing	1.1	17

(Continued)

Table 3. (Continued)

Characteristic	Percentage	Frequency, n
<i>Hours worked in a typical week</i>		
<20	1.3	21
20–39	10.6	170
40–59	82.7	1326
≥60	3.0	48
Missing	2.4	39
<i>In your role you typically have direct interaction or contact with patients</i>		
Yes	85.6	1373
No	9.0	145
Missing	5.4	86
<i>Your primary roles in the medicines process are (multiple options could be chosen)</i>		
Prescribing	15.1	243
Administering	61.1	980
Preparation and Dispensing	25.9	415
Monitoring	42.0	673
Missing	3.1	49

<https://doi.org/10.1371/journal.pone.0204801.t003>

oldest most positive, $X^2(4, N = 1564) = 28.89, p < 0.001$) and experience as a health professional (positive responses increasing with experience, $X^2(1, N = 1550) = 42.06, p < 0.001$).

For those ten composites with higher mean agreement, several items had less than half responding positively. There were issues around: supervisors/ managers overlooking recurring patient safety problems (31.9% positive); that it was due to chance that serious errors did not occur (36.0%); problems occurring when exchanging information across hospital units (42.9%); staff being able to ask questions if things did not seem right (44.0%); that at particular pressure points supervisors/ managers wanted staff to work faster, even if this required short-cuts to be taken (46.1%); and staff feeling able to question those in positions of authority (46.6%).

More detailed data on the responses to individual items within each composite are given in [S1 File](#).

Focus groups

Demographics of participants. Two hundred and ninety-five survey respondents (18.4%) expressed interest in participating in focus groups. Nine focus groups were conducted (duration 45–60 minutes), at which point data saturation was deemed to have been achieved. Fifty-four individuals from different disciplines participated, with just under half ($n = 26, 48.1\%$) being nurses, followed by 18 (33.3%) pharmacists and 10 (18.5%) doctors. Most were highly experienced with only 11 (20.4%) having <5 years of experience. During the focus groups, there was wide-ranging discussion across the spectrum of medication errors of prescribing, administration and dispensing.

Behavioural determinants associated with errors. Themes and subthemes relating to safety culture identified during focus group discussions are mapped to TDF behavioural determinants, with illustrative quotes provided for each.

A. Social/professional role and identity (a coherent set of behaviours and displayed personal qualities of an individual in a social or work setting)

Table 4. Positive responses to HSOPS items and composites (N = 1604).

Statements	% positive response (100% representing the highest positive response to each statement)
Non-punitive response to errors , overall positive response = 24.0%**	
*Staff feel like errors count against them	26.2 (disagreed)
*When an error is reported, it feels like the person is being reported, not the problem	31.1 (disagreed)
*Staff worry that errors they make are kept in their personnel file	14.6 (disagreed)
Staffing , overall positive response = 36.2%	
We have enough staff to handle the workload	54.7 (agreed)
*We use more locum staff than is best for patient care	30.5 (disagreed)
*We work under pressure trying to do too much, too quickly	23.5 (disagreed)
Communication openness , overall positive response = 50.5%	
Staff will speak up freely if they see something that may negatively affect patient care	60.9 (agreed)
Staff feel free to question the decisions or actions of those with more authority	46.6 (agreed)
*In this unit, staff are afraid to ask questions when something does not seem right	44.0 (disagreed)
Handoffs and transitions , overall positive response = 53.1%	
*Things get missed when transferring patients from one unit to another	53.7 (disagreed)
*Important patient care information is often lost during shift changes	60.8 (disagreed)
*Problems often occur in the exchange of information across hospital units	42.9 (disagreed)
*Shift changes are problematic for patients in this hospital	55.1 (disagreed)
Supervisor/manager expectations and actions promoting patient safety , overall positive response = 56.5%	
My supervisor/ manager says a good word when he/she sees a job done according to established patient safety procedures	73.0 (agreed)
My supervisor/ manger seriously considers staff suggestions for improving patient safety	74.9 (agreed)
*Whenever pressure builds up, my supervisor/ manager wants us to work faster, even if it means taking shortcuts	46.1 (disagreed)
*My supervisor/ manager overlooks patient safety problems that happen again and again	31.9 (disagreed)
Frequency of events reported , overall positive response = 58.1%	
When an error is made, but is noticed and corrected before affecting the patient, how often is this reported?	53.5 (agreed)
When an error is made, but has no potential to harm the patient, how often is this reported?	56.9 (agreed)
When an error is made that could potentially harm the patient but does not, how often is this reported?	63.8 (agreed)
Overall perceptions of patient safety , overall positive response = 59.1%	
Patient safety is never sacrificed to get more work done	70.6 (agreed)
Our procedures and systems are good at preventing errors from happening	78.7 (agreed)
*It is just by chance that more serious mistakes don't happen around here	36.0 (disagreed)
*We have patient safety problems in this unit	51.3 (disagreed)
Feedback and communication about error , overall positive response = 61.9%	
We are given feedback about changes put into place based on error reports	55.8 (agreed)

(Continued)

Table 4. (Continued)

Statements	% positive response (100% representing the highest positive response to each statement)
We are informed about medication errors in this unit	62.0 (agreed)
In this unit, we discuss ways to prevent medication errors from happening again	68.0 (agreed)
Teamwork across units , overall positive response = 67.7%	
There is good cooperation among hospital units that need to work together	72.9 (agreed)
Hospital units work well together to provide the best care for patients	82.8 (agreed)
*Hospital units do not coordinate well with each other	57.5 (disagreed)
*It is often unpleasant to work with staff from other hospital units	57.5 (disagreed)
Management support for patient safety , overall positive response = 75.4%	
Hospital management provides a work environment that promotes patient safety	87.0 (agreed)
The actions of hospital management show that patient safety is a top priority	84.2 (agreed)
Hospital management seems interested in patient safety only after an error happens	54.9 (agreed)
Teamwork within units , overall positive response = 82.1%	
People support one another in this unit	81.1 (agreed)
When a lot of work needs to be done quickly, we work as a team to get the work done	83.4 (agreed)
In this unit, people treat each other with respect	81.9 (agreed)
Organisational learning—continuous improvement , Overall positive response = 85.8%	
We are actively doing things to improve patient safety	90.2 (agreed)
After we make changes to improve patient safety, we evaluate their effectiveness	81.3 (agreed)

*Reverse scored negatively worded statement

** Calculated from the mean items within each composite

<https://doi.org/10.1371/journal.pone.0204801.t004>

1. Doctors reliance on pharmacists to correct errors. During discussion, it emerged that there were instances where doctors would rely on pharmacists to correct their prescribing errors and this led to complacency around prescribing,

‘Yes. Most of the physicians make a medication error and wait for the pharmacist to correct it.’
(Focus Group [FG] 5 Pharmacist 4)

2. Doctors reluctance to alter other doctors’ prescribing. During one focus group, there was concern that doctors were unwilling to alter prescriptions written by other doctors, particularly for doctors from other specialities. The doctors considered this to be the responsibility of the original prescriber, even if a prescribing error had been made and initial prescriber was unavailable,

‘This will happen when you’re in the Ob-Gyn [obstetrics and gynaecology] setup. If one physician came from Hamad from other. . . from cardiac or other site, if they write any prescription, if you call the Ob-Gyn doctor here, the on duty doctor, she will never agree to change because

she will say it's an order from the consultant from cardiology or neurology.' (FG7 Pharmacist 4)

3. Lack of recognition of the role of nurses. Some of the nurses described that they were often omitted from discussions around patient care and decision making, even when present on ward rounds or meetings. There were instances where discussions took place in a different language,

'Even I'm noting that during the rounds, team decisions, the nurses are not informed. Sometimes they [the doctors] are discussing in Arabic. The nurse, she cannot understand their plan and what is the decision.' (FG3 Nurse 1)

4. Policy non-adherence. Health professionals not adhering to various policies was considered a cause of medication errors,

'Not abiding the. . . complying with the policies' (FG2 Doctor 2)

'There are seven or eight points that the pharmacist should check. If the pharmacist, for example, dispensed the wrong medication it means that he didn't follow the policy.' (FG5 Pharmacist 4)

B. Emotions (a complex reaction pattern, involving experiential, behavioural, and physiological elements, by which the individual attempts to deal with a personally significant matter or event)

1. Stress leading to medication errors. Stress and high pressure situations were described in all focus groups as influences on medication errors. While workload was a common factor leading to stress, patients themselves could also put undue pressure and hence stress of health professionals,

'And I think that probably the stresses of the work [lead to errors].' (FG1 Doctor 2)

'And parents are too tense than they are. . . even the parents they are too much angry. Yeah, they will scold the staff then like that time they will get pressure.' (FG7 Nurse 3)

C. Environmental Context and Resources (any circumstance of a person's situation or environment that discourages or encourages the development of skills and abilities, independence, social competence, and adaptive behaviour)

Much of the discussion centred on aspects of environmental context and resources as key influences leading to medication errors. These were discussed by all participants in all focus groups. There were several key themes within this domain.

1. Workload issues leading to medication errors. Workload issues were discussed by doctors, nurses and pharmacists. Doctors believed one of the reasons for errors to happen was the heavy workload that they had.

'Too many patients. Labour ward is full, you know, too many patients for the residents to see.' (FG1 Doctor 2)

'Yeah, I'm working in emergency. So what I feel is it's too much. . . sometime it is too busy and doctors are giving too much orders. . . they cannot to cope with the situation.' (FG1 Nurse 1)

One pharmacist noted that the excessive workload for the doctor can lead to errors occurring and that this workload also put pressure on other health professionals which could compound errors.

‘There are two problems here, a load on the physician that can lead to many mistakes and a load on the pharmacist because he needs to dispense medication for this patient and at the same time answer the questions of physician, nurses.’ (FG5 Pharmacist 4)

One of the nurses also explained that the main cause of errors committed by junior medical staff was workload rather than lack of knowledge.

‘And this is why the medication errors are also increasing, so it’s not always related to the knowledge of the resident. And if the resident is overloaded because he has to document for all the patients and see all the patients and he is receiving calls from other units as well’ (FG3 Nurse 3)

2. Lack of staff at key times. Closely related to workload issues was a critical lack of staff at key times such as weekends and evening which could compromise patient safety.

‘On the whole days of the week, there is complete staff, complete number of physicians. In weekend, well, only one physician is doing the whole work.’ (FG4 Doctor 2)

‘Especially the areas like emergency, less staff. They will be get. . . too tense by the patients and they just want to do the things for faster. so it will make errors. (FG2 Nurse 1)

3. System-related issues. Discussion also centred on key issues related to the systems in operation in various wards and departments. There was particular concern over the implementation of Cerner (electronic health record system for hospitals, health care providers, clinics) from doctors, nurses and pharmacists.

‘The electronic system is not robust, and I mean, the hardware is not good enough.’ (FG2 Doctor 1)

‘We have now to concentrate on the mistakes or medication errors happening by the prescribing system.’ (FG5 Pharmacist 2)

One senior doctor commented that following implementation of Cerner, fewer checks were being performed compared to the previous paper-based system.

‘Before it was like, when you have the hard copy of medication profile, someone is checking and countersigning. Now in the system, it [checking] is not there as far as I know.’ (FG1 Doctor 2)

Themes and subthemes for those behavioural determinants less related to safety culture are summarised in [Table 5](#).

Discussion

Key findings

Our study of the causes of medication errors in Qatar highlighted that the key composites of patient safety culture which merit attention are: non-punitive response to errors; staffing;

Table 5. A summary of TDF domains and themes (less related to culture) relating to causes of medication errors.

TDF Domain	Subtheme	Illustrative quotes
Knowledge	1. Lack of medication related knowledge	'So coming to the nursing knowledge regarding the dose. I will never believe they have that much knowledge about the doses...' (FG1 Doctor 1)
	2. Knowledge is limited to a particular speciality/area	'If we're dealing with the general hospital, medicine department they have good orientation regarding medication, but if you go to ortho [orthopaedics] or surgery, really their knowledge about medication is very low.' (FG5 Pharmacist 3)
	3. Lack of knowledge attributed to staff induction	'Proper induction, you know, they should have proper induction regarding the medication, the medications that are used, how you do the checking and things like that. Nothing is done.' (FG1 Doctor 2)
	4. Need for continuing professional development to reduce medication errors	'There is too much error in this area, they can provide another or a new continuous education for this field. It's very important and this can prevent such error.' (FG7 Nurse 1)
Skills	1. Suboptimal medication related skills	'We need to think about the administration. I have seen plenty of times the paper on which they [nurses] have written the calculation and it's wrong, actually most of the time.' (FG4 Pharmacist 1)
Beliefs about Capabilities	1. Lack of medication related competence	'But you think it's... it's... it's valid to let the nurses check the dose before administering? No, I don't think it's possible. For me, I feel it's impossible for them to check the correct dose.' (FG1 Doctor 1)
	2. Overconfidence leading to medication errors	'Overconfidence with some particular medicines like I have been with this medicine for many years and I know by heart' (FG1 Pharmacist 2)
Goals	1. Promoting patient safety	'But you know, serious errors are part of the package, you know. As we save lives, we are not ensuring... I mean, we should expect that we cannot have zero even serious errors because we are human beings'. (FG5 Pharmacist 1)

<https://doi.org/10.1371/journal.pone.0204801.t005>

communication openness; handoffs and transitions; and supervisor/manager expectations and actions promoting patient safety. During focus group discussions, specific TDF determinants suggested as being potentially associated with these composites were: social/professional role and identity; emotions; and environmental context and resources. Thematic analysis identified issues of doctors relying on pharmacists to correct their errors and being reluctant to alter the prescribing of fellow doctors. There was a lack of recognition of nurses' roles and frequent policy non-adherence. Stress was perceived to be a major contributor to errors, as was excessive workload and lack of staff at key times.

Strengths and weaknesses

The mixed methods design is a key study strength providing quantification of results followed by in-depth explanation. Further strengths are the use of the validated HSOPS tool and embedding psychological behaviour change theory (TDF) within qualitative data generation and analysis [15,19]. There are, however, several limitations hence findings should be interpreted with caution. Self-reported questionnaire responses could not be validated and may have been impacted by response and social desirability biases [22]. While responses were received from healthcare professionals in all HMC hospitals, these may have been skewed towards females and nurses hence there are potential issues of lack of generalisability within Qatar and beyond. Similarly, qualitative findings may not be transferable to other healthcare professionals, settings and countries.

Interpretation

This mixed methods study has contributed to the expressed need for robustness and rigour in patient safety research within the Middle East [20]. Furthermore, it aligns to the WHO 'Global Patient Safety Challenge' calling for action to reduce severe, avoidable medication-related harm by 50% in the next five years [2,3]. Whilst the HSOPS questionnaire has been used

within the Middle East [21], this is the first study to publish Qatari data. There are, however, similarities between the Qatari data and those reported by Elmontsri et al. [21], with the lowest agreement scores (and hence of most concern) relating to the composites of non-punitive response to errors; staffing; communication openness; handoffs and transitions; and supervisor/ manager expectations and actions promoting patient safety. Within the two composites of lowest scores (non-punitive response to errors and staffing) there were issues with staff perceiving that errors counted against them and that details of errors committed were kept in their personnel files. This appeared to be an issue for male, younger and less experienced healthcare professionals. Staffing was the other key composite with very low agreement scores, particularly in relation to work pressures and speed of work, with similar statistically significant associations as for the non-punitive response to errors. There may be some merit in initially prioritising any intervention towards these specific groupings.

One limitation of the published studies using the HSOPS is the lack of qualitative research to provide in-depth explanation of the results [21]. The use of behavioural theory within the focus groups in this study identified key determinants which could facilitate intervention development. TDF has been incorporated within intervention developments for smoking cessation, physical activity, hand hygiene, acute low back pain and schizophrenia [28]. To date only one other published study has applied TDF to explore potential causes of medication errors, focusing on prescribing errors in a sample of junior doctors in Scotland [29]. There are some similarities with the findings of this study, most notably within the domains of knowledge and skills, particularly the general lack of medication-related knowledge. While pharmacists can provide support, and indeed doctors were found to rely on pharmacists to correct errors, the HSOPS data and the focus groups identified issues around staff complement and workload, particularly at key times.

TDF domains of social/ professional role and identity, emotions and environmental context and resources are related to organisational safety culture, as defined by 'Study Group on Human Factors' [16]. Concerns were expressed around nurses perceiving that their professional role was not recognised leading to poor communication compromising patient safety. This is also reflected in the HSOPS score of ~ 50% agreement for communication openness. There were instances of doctors relying on pharmacists to correct their prescribing errors and, at times, would not alter the prescribing of others, even when errors could potentially lead to patient harm. Themes of environmental context and resources also emerged in the discussions around workload as a leading cause of errors, with lack of staff at key pressure times of evening and weekends. Furthermore, the electronic prescribing and records system was considered to have introduced potential for error. While such systems have been shown to enhance patient safety, others have also highlighted the risky human factors and user-centred design issues that have been encountered [13].

Stress was the main theme which emerged in the TDF emotions domain as a determinant of error, arising due to workload, work pressures and the influence of patients. Issues of workload were also identified in the HSOPS data around staff numbers to handle the workload, working under pressure to do too much, too quickly.

These TDF I determinants which were highlighted as potential contributors to medication errors can be used during the development of behaviour change interventions, defined as 'coordinated sets of activities designed to change specified behaviour patterns'. These are often complex, consisting of interacting components known as 'behaviour change techniques' (BCTs), 'observable and replicable components designed to change behaviour' [30]. Michie et al. developed a cross-disciplinary taxonomy of evidence based BCTs [31], mapped to specific TDF domains [32]. Whilst knowledge and skills can be impacted through education and training [31,32], altering aspects of social/ professional role and identity and environmental context

and resources are more complex. Indeed, the work of Michie et al. [31,32] did not identify any evidence-based BCTs which mapped reliably to social/professional role and identity. Those for environmental context and resources relating mainly to restructuring the physical environment and providing prompts and cues for safer practice, which in this case would focus on the electronic medication systems [31,32]. Rather than focusing on changing behaviour at the individual practitioner level, action may be required at the organisational strategic level to review policies, structures (including resource allocation and distribution) and processes which aim to promote patient safety culture and minimise harm. Qualitative research focusing on understanding the perspectives of key strategic decision-makers in relation to promoting all aspects of medication safety is warranted.

Conclusion

This mixed methods study has provided further confirmation of key areas of concern relating to patient safety culture in Qatar. Non-punitive response to errors and staffing had the lowest levels of agreement, followed by communication openness, handoffs and transitions, and supervisor/manager expectations and actions. The qualitative component provided further detail of specific TDF determinants highlighting issues of social/professional role and identity, emotions, and environmental context and resources. Further attention on these issues at strategic and policy levels is required.

Supporting information

S1 File. Responses to each of the HSOPS composites.

(DOCX)

S2 File. Study questionnaire.

(DOCX)

Acknowledgments

The authors wish to acknowledge the contributions of all questionnaire respondents and focus group participants, as well as support departments at Hamad Medical Corporation, Doha, Qatar.

Author Contributions

Conceptualization: Derek Stewart, Katie MacLure, Abdulrouf Pallivalapila, Ahmed Awaisu, James S. McLay, Kerry Wilbur, Kyle Wilby, Cristin Ryan, Rajvir Singh, Moza Al Hail.

Data curation: Derek Stewart, Binny Thomas, Katie MacLure, Abdulrouf Pallivalapila, Wesam El Kassem, Ahmed Awaisu, James S. McLay, Kerry Wilbur, Kyle Wilby, Cristin Ryan, Andrea Dijkstra, Rajvir Singh, Moza Al Hail.

Formal analysis: Derek Stewart, Binny Thomas, Katie MacLure, Abdulrouf Pallivalapila, Ahmed Awaisu, Kerry Wilbur, Kyle Wilby, Cristin Ryan, Andrea Dijkstra, Rajvir Singh.

Funding acquisition: Derek Stewart, Abdulrouf Pallivalapila, Ahmed Awaisu, James S. McLay, Kerry Wilbur, Kyle Wilby, Cristin Ryan, Rajvir Singh.

Methodology: Derek Stewart, Binny Thomas, Katie MacLure, Abdulrouf Pallivalapila, Wesam El Kassem, Ahmed Awaisu, James S. McLay, Kerry Wilbur, Kyle Wilby, Cristin Ryan, Andrea Dijkstra, Rajvir Singh, Moza Al Hail.

Supervision: Derek Stewart, Rajvir Singh.

Writing – original draft: Derek Stewart.

Writing – review & editing: Binny Thomas, Katie MacLure, Abdulrouf Pallivalapila, Wessam El Kassem, Ahmed Awaisu, James S. McLay, Kerry Wilbur, Kyle Wilby, Cristin Ryan, Andrea Dijkstra, Rajvir Singh, Moza Al Hail.

References

1. Jha AK, Larizgoitia I, Audera-Lopez C, Prasopa-Plaizier N, Waters H, Bates DW. The global burden of unsafe medical care: analytic modelling of observational studies. *BMJ Qual Saf* 2013; 22:809–15. <https://doi.org/10.1136/bmjqs-2012-001748> PMID: 24048616
2. World Health Organization. Medication Without Harm, WHO Global Patient Safety Challenge. Geneva: World Health Organization, 2017.
3. Donaldson LJ, Kelley ET, Dhingra-Kumar N, Kienny MP, Sheikh A. Medication Without Harm: WHO's Third Global Patient Safety Challenge. *Lancet* 2017; 389:1680–81. [https://doi.org/10.1016/S0140-6736\(17\)31047-4](https://doi.org/10.1016/S0140-6736(17)31047-4) PMID: 28463129
4. Lewis PJ, Dorman T, Taylor D, Tully MP, Wass V, Ashcroft DM. Prevalence, incidence and nature of prescribing errors in hospital inpatients. *Drug Saf* 2009; 32:379–89. <https://doi.org/10.2165/00002018-200932050-00002> PMID: 19419233
5. Ross S, Bond C, Rothnie H, Thomas S, Macleod MJ. What is the scale of prescribing errors committed by junior doctors? A systematic review. *Brit J Clin Pharmacol* 2009; 67:629–40.
6. James KL, Barlow D, McArtney R, Hiom S, Roberts D, Whittlesea C. Incidence, type and causes of dispensing errors: a review of the literature. *Int J Pharm Pract* 2009; 17(1):9–30. PMID: 20218026
7. Franklin BD, Reynolds M, Shebl NA, Burnett S, Jacklin A. Prescribing errors in hospital inpatients: a three-centre study of their prevalence, types and causes. *Postgrad Med J* 2011; 87:739–45. <https://doi.org/10.1136/pgmj.2011.117879> PMID: 21757461
8. Keers RN, Williams SD, Cooke J, Ashcroft DM. Prevalence and nature of medication administration errors in health care settings: a systematic review of direct observational evidence. *Ann Pharmacother* 2013; 47:237–56. <https://doi.org/10.1345/aph.1R147> PMID: 23386063
9. National Coordinating Council for Medication Error Reporting and Prevention. Available at <http://www.nccmerp.org> [accessed June 2017].
10. Tully MP, Ashcroft DM, Dorman T, Lewis PJ, Taylor D, Wass V. The causes of and factors associated with prescribing errors in hospital inpatients. *Drug Saf* 2009; 32:819–36. <https://doi.org/10.2165/11316560-000000000-00000> PMID: 19722726
11. Sears K, Ross-White A, Godfrey CM. The incidence, prevalence and contributing factors associated with the occurrence of medication errors for children and adults in the community setting: A systematic review. *JBHI Database Syst Rev Implement* 2012; 10:2350–464.
12. Keers RN, Williams SD, Cooke J, Ashcroft DM. Causes of medication administration errors in hospitals: a systematic review of quantitative and qualitative evidence. *Drug Saf* 2013; 36(11):1045–67. <https://doi.org/10.1007/s40264-013-0090-2> PMID: 23975331
13. Brown CL, Mulcaster HL, Triffitt KL, Sittig DF, Ash JS, Reygate K et al. A systematic review of the types and causes of prescribing errors generated from using computerized provider order entry systems in primary and secondary care. *J Am Med Inform* 2016;ocw119.
14. Craig P, Dieppe P, Macintyre S, Michie S, Nazareth I, Petticrew M. Developing and evaluating complex interventions: the new Medical Research Council guidance. *BMJ* 2008; 337:a1665.
15. Cane J, O'Connor D, Michie S. Validation of the theoretical domains framework for use in behaviour change and implementation research. *Implement Sci* 2012; 7:37. <https://doi.org/10.1186/1748-5908-7-37> PMID: 22530986
16. Study Group on Human Factors. Organising for safety: third report of the ACSNI (Advisory Committee on the Safety of Nuclear Installations). England: HSE Books, 1993.
17. Morello RT, Lowthian JA, Barker AL, McGinnes R, Dunt D, Brand C. Strategies for improving patient safety culture in hospitals: a systematic review. *BMJ Qual Saf* 2013; 22:11–18. <https://doi.org/10.1136/bmjqs-2011-000582> PMID: 22849965
18. Weaver SJ, Lubomksi LH, Wilson RF, Pfoh ER, Martinez KA, Dy SM. Promoting a culture of safety as a patient safety strategy: a systematic review. *Ann Intern Med* 2013; 158:369–374. <https://doi.org/10.7326/0003-4819-158-5-201303051-00002> PMID: 23460092

19. Sorra J, Gray L, Streagle S, Famolaro T, Yount N, Behm J. AHRQ Hospital Survey on Patient Safety Culture: User's Guide. (Prepared by Westat, under Contract No. HHS290201300003C). AHRQ Publication No. 15-0049-EF (Replaces 04-0041). Rockville, MD: Agency for Healthcare Research and Quality. Available at <http://www.ahrq.gov/professionals/quality-patient-safety/patientsafetyculture/hospital/index.html> [accessed June 2017].
20. Alsulami Z, Conroy S, Choonara I. Medication errors in the Middle East countries: a systematic review of the literature. *Eur J Clin Pharmacol* 2013; 69:995–1008. <https://doi.org/10.1007/s00228-012-1435-y> PMID: 23090705
21. Elmontsri M, Almashrafi A, Banarsee R, Majeed A. Status of patient safety culture in Arab countries: a systematic review. *BMJ Open* 2017; 7:e013487. <https://doi.org/10.1136/bmjopen-2016-013487> PMID: 28237956
22. Creswell J, Clark V. Choosing a mixed method design. Designing and conducting mixed method research: SAGE Inc., 2006.
23. Pluye P, Hong QN. Combining the power of stories and the power of numbers: mixed methods research and mixed studies reviews. *Annu Rev Public Health* 2014; 35:29–45. <https://doi.org/10.1146/annurev-publhealth-032013-182440> PMID: 24188053
24. Barlett JE, Kotrlik JW, Higgins CC. Organizational research: Determining appropriate sample size in survey research. *Inf Technol Learn Perform J* 2001; 19:43.
25. Shenton AK. Strategies for ensuring trustworthiness in qualitative research projects. *Educ Inform* 2004; 22:63–67.
26. Fusch PI, Ness LR. Are we there yet? Data saturation in qualitative research. *Qual Rep* 2015; 20:1408.
27. Ritchie J, Spencer L. Qualitative data analysis for applied policy research. *The Qualitative Researcher's Companion* 2002:305–329.
28. Francis JJ, O'Connor D, Curran J. Theories of behaviour change synthesised into a set of theoretical groupings: introducing a thematic series on the theoretical domains framework. *Implement Sci* 2012; 7:35. <https://doi.org/10.1186/1748-5908-7-35> PMID: 22531601
29. Duncan EM, Francis JJ, Johnston M, Davey P, Maxwell S, McKay GA, et al. Learning curves, taking instructions, and patient safety: using a theoretical domains framework in an interview study to investigate prescribing errors among trainee doctors. *Implement Sci* 2012; 7(1):86.
30. Michie S, van Stralen MM, West R. The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implement Sci* 2011; 6:42. <https://doi.org/10.1186/1748-5908-6-42> PMID: 21513547
31. Michie S, Richardson M, Johnston M, Abraham C, Francis J, Hardeman W, et al. The behavior change technique taxonomy (v1) of 93 hierarchically clustered techniques: building an international consensus for the reporting of behavior change interventions. *Ann Behav Med* 2013; 46:81–95. <https://doi.org/10.1007/s12160-013-9486-6> PMID: 23512568
32. Michie S, Wood CE, Johnston M, Abraham C, Francis J, Hardeman W. Behaviour change techniques: the development and evaluation of a taxonomic method for reporting and describing behaviour change interventions (a suite of five studies involving consensus methods, randomised controlled trials and analysis of qualitative data). *Health Technol Assess* 2015; 19.

SUPPLEMENTARY FILE

[illegible]

Table 2. Supervisor/manager expectations and actions promoting patient safety, responses to HSOPS items (N=1604)

Statements	Strongly Agree	Agree	Unsure	Disagree	Strongly Disagree	Missing	% positive response (100% the highest positive response to each statement)
	% (n)	% (n)	% (n)	% (n)	% (n)	% (n)	
My supervisor/manager says a good word when he/she sees a job done according to established patient safety procedures	21.4 (344)	51.6 (827)	13.5 (216)	8.5 (136)	3.4 (54)	1.7 (27)	73.0
My supervisor/manager seriously considers staff suggestions for improving patient safety	20.7 (332)	54.2 (870)	14.0 (224)	6.4 (102)	2.6 (42)	2.1 (34)	74.9
*Whenever pressure builds up, my supervisor/manager wants us to work faster, even if it means taking shortcuts	6.3 (101)	22.9 (367)	22.3 (357)	37.5 (601)	8.6 (138)	2.5 (40)	46.1
*My supervisor/manager overlooks patient safety problems that happen again and again	11.7 (188)	39.8 (638)	13.7 (219)	22.6 (363)	9.3 (149)	2.9 (47)	31.9
Overall positive response = 56.5%							

*negatively worded so reverse scored

Table 3. Organisational learning - continuous improvement, responses to HSOPS items (N=1604)

[illegible]

Table 4. Management support for patient safety, responses to HSOPS items (N=1604)

[illegible]

Table 5. Overall perceptions of patient safety, responses to HSOPS items (N=1604)

[illegible]

*negatively worded so reverse scored

Table 6. Feedback and communication about error, responses to HSOPS items (N=1604)

[illegible]

Table 7. Communication openness, responses to HSOPS items (N=1604)

Statements	Always	Most of the time	Some times	Rarely	Never	Missing	% positive response (100% the highest positive response to each statement)
	% (n)	% (n)	% (n)	% (n)	% (n)	% (n)	
Staff will speak up freely if they see something that may negatively affect patient care	21.6 (347)	39.3 (631)	25.6 (410)	8.2 (132)	2.6 (42)	2.6 (42)	60.9
Staff feel free to question the decisions or actions of those with more authority	14.7 (236)	31.9 (511)	28.9 (463)	15.5 (248)	6.0 (96)	3.1 (50)	46.6
*In this unit, staff are afraid to ask questions when something does not seem right	7.4 (118)	17.5 (280)	28.1 (450)	25.1 (402)	18.9 (303)	3.2 (51)	44.0
Overall positive response = 50.5%							

*negatively worded so reverse scored

Table 8. Frequency of events reported, responses to HSOPS items (N=1604)

[illegible]

Table 9. Teamwork across units, responses to HSOPS items (N=1604)

Statements	Strongly Agree	Agree	Unsure	Disagree	Strongly Disagree	Missing	% positive response (100% the highest positive response to each statement)
	% (n)	% (n)	% (n)	% (n)	% (n)	% (n)	
There is good cooperation among hospital units that need to work together	15.7 (252)	57.2 (918)	15.4 (247)	6.5 (104)	1.9 (30)	3.3 (53)	72.9
Hospital units work well together to provide the best care for patients	31.7 (509)	51.1 (820)	9.4 (151)	2.7 (44)	1.2 (19)	3.8 (61)	82.8
*Hospital units do not coordinate well with each other	5.4 (86)	18.1 (291)	15.5 (249)	45.6 (732)	11.9 (191)	3.4 (55)	57.5
*It is often unpleasant to work with staff from other hospital units	3.2 (51)	16.6 (266)	19.5 (313)	46.4 (745)	11.1 (178)	3.2 (51)	57.5
Overall positive response = 67.7%							

*negatively worded so reverse scored

Table 10. Staffing, responses to HSOPS items (N=1604)

Statements	Strongly Agree % (n)	Agree % (n)	Unsure % (n)	Disagree % (n)	Strongly Disagree % (n)	Missing % (n)	% positive response (100% the highest positive response to each statement)
We have enough staff to handle the workload	13.2 (211)	41.5 (665)	14.7 (235)	21.8 (349)	7.5 (120)	1.5 (24)	54.7
*We use more locum staff than is best for patient care	9.0 (145)	29.5 (473)	26.2 (421)	22.4 (360)	8.1 (130)	4.7 (75)	30.5
*We work under pressure trying to do too much, too quickly	17.5 (280)	40.0 (641)	17.2 (276)	20.2 (324)	3.3 (53)	1.9 (30)	23.5
Overall positive response = 36.2%							

*negatively worded so reverse scored

Chi-square analysis

1. Association between age and positive responses

	20-29 years	30-39 years	40-49 years	50-59 years	60-69 years
Positive responses	359	733	411	140	20
Negative responses	790	1220	603	305	54

Three items hence each individual represented by three counts

Youngest least and oldest most positive, $X^2 (4, N=1564)=28.89, p<0.001$.

2. Association between years of experience and positive responses

	<1 year	1-5 years	5-10 years	11-15 years	16-20 years	>20 years
Positive responses	16	292	459	416	240	292
Negative responses	50	601	911	596	329	391

Three items hence each individual represented by three counts

Positive responses increase with experience, $X^2 (1, N=1550)=42.06, p<0.001$.

3. Association between profession and positive responses

	Doctors	Nurses	Pharmacists
Positive responses	258	1236	169
Negative responses	366	1966	433

Three items hence each individual represented by three counts

Doctors most positive and pharmacists least, $X^2 (2, N=1494)=42.06, p<0.001$.

Table 11. Handoffs and transitions, responses to HSOPS items (N=1604)

[illegible]

Table 12. Non-punitive response to errors, responses to HSOPS items (N=1604)

[illegible]

Chi-square analysis

1. Association between age and positive responses

	20-29 years	30-39 years	40-49 years	50-59 years	60-69 years
Positive responses	276	447	273	126	24
Negative responses	882	1518	738	330	51

Three items hence each individual represented by three counts

Older most positive, $X^2 (4, N=1555)=11.62, p<0.05$.

2. Association between years of experience and positive responses

	<1 year	1-5 years	5-10 years	11-15 years	16-20 years	>20 years
Positive responses	13	201	299	269	146	201
Negative responses	46	688	1084	742	423	493

Three items hence each individual represented by three counts

Most experienced being most positive, $X^2 (5, N=1536)= 18.42, p<0.005$.

3. Association between gender and positive responses

	Female	Male
Positive responses	855	279
Negative responses	2490	1017

Three items hence each individual represented by three counts

Females most positive, $X^2 (1, N=1547)=8.23, p<0.005$.

We will start with some information about you which will help with the analysis of the data.

SECTION 1: Demographics

1. Are you:

☐ a. male

☐ b. female

2. Your age in years is:

☐ a. under 20

☐ c. 30 - 39

☐ e. 50 - 59

☐ b. 20 - 29

☐ d. 40 - 49

☐ f. 60 - 69

☐ g. over 69

3. Your country of origin is:

☐ a. Qatar

☐ d. India

☐ h. Philippines

☐ b. other GCC

☐ e. Jordan

☐ i. Sudan

☐ c. Egypt

☐ f. Lebanon

☐ j. UK

☐ g. Palestine

☐ k. Other, please specify:

4. Your country of receiving entry-to-practice degree was:

☐ a. Qatar

☐ d. India

☐ h. Philippines

☐ b. other GCC

☐ e. Jordan

☐ i. Sudan

☐ c. Egypt

☐ f. Lebanon

☐ j. UK

☐ g. Palestine

☐ k. Other, please specify:

5. Your country of receiving highest academic degree was:

☐ a. Qatar

☐ d. India

☐ h. Philippines

☐ b. other GCC

☐ e. Jordan

☐ i. Sudan

☐ c. Egypt

☐ f. Lebanon

☐ j. UK

☐ g. Palestine

☐ k. Other, please specify:

6. Your highest academic degree is:

☐ a. BSc (Nursing)

☐ e. MSc

☐ i. PhD

☐ b. BSc (Pharm)

☐ f. MBBS or MBChB

☐ j. Additional professional qualification, please specify:

☐ c. MPharm

☐ g. MD

☐ d. PharmD

☐ h. HND/Diploma

7. How long have you worked in this hospital?

- | | | |
|------------------------------------------|--------------------------------------------|----------------------------------------------|
| <input type="checkbox"/> a. < 1 year | <input type="checkbox"/> c. 6 to 10 years | <input type="checkbox"/> e. 16 to 20 years |
| <input type="checkbox"/> b. 1 to 5 years | <input type="checkbox"/> d. 11 to 15 years | <input type="checkbox"/> f. 21 years or more |

8. How long have you worked in your current hospital work area/unit?

- | | | |
|------------------------------------------|--------------------------------------------|----------------------------------------------|
| <input type="checkbox"/> a. < 1 year | <input type="checkbox"/> c. 6 to 10 years | <input type="checkbox"/> e. 16 to 20 years |
| <input type="checkbox"/> b. 1 to 5 years | <input type="checkbox"/> d. 11 to 15 years | <input type="checkbox"/> f. 21 years or more |

9. Typically, how many hours per week do you work in this hospital?

- | | |
|-----------------------------------------------------|-----------------------------------------------------|
| <input type="checkbox"/> a. < 20 hours per week | <input type="checkbox"/> c. 40 to 59 hours per week |
| <input type="checkbox"/> b. 20 to 39 hours per week | <input type="checkbox"/> d. > 60 hours per week |

10. What is your role in this hospital? Select ONE answer that best describes your role

- | | | |
|----------------------------------------------------|-----------------------------------------------------|----------------------------------------------------------|
| <input type="checkbox"/> a. Consultant Physician | <input type="checkbox"/> d. Clinical Nurse Educator | <input type="checkbox"/> g. Clinical Pharmacy Specialist |
| <input type="checkbox"/> b. Specialist Physician | <input type="checkbox"/> e. Specialist Nurse | <input type="checkbox"/> h. Clinical Pharmacist |
| <input type="checkbox"/> c. Resident Physician | <input type="checkbox"/> f. Nurse | <input type="checkbox"/> i. Pharmacist |
| <input type="checkbox"/> j. Other, please specify: | | |

11. In your role, do you typically have direct interaction or contact with patients?

- | | |
|-----------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------|
| <input type="checkbox"/> a. YES, I typically have direct interaction or contact with patients | <input type="checkbox"/> b. NO, I typically do NOT have direct interaction or contact with patients |
|-----------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------|

12. What are your primary roles in the medicines process? (Tick all that apply)

- | | | |
|------------------------------------------------------|-------------------------------------------|----------------------------------------------------|
| <input type="checkbox"/> a. prescribing | <input type="checkbox"/> c. administering | <input type="checkbox"/> e. Other, please specify: |
| <input type="checkbox"/> b. preparation & dispensing | <input type="checkbox"/> d. monitoring | |

13. How long have you worked in your current role?

- | | | |
|------------------------------------------|--------------------------------------------|----------------------------------------------|
| <input type="checkbox"/> a. < 1 year | <input type="checkbox"/> c. 6 to 10 years | <input type="checkbox"/> e. 16 to 20 years |
| <input type="checkbox"/> b. 1 to 5 years | <input type="checkbox"/> d. 11 to 15 years | <input type="checkbox"/> f. 21 years or more |

14. How long have you practiced your profession in Qatar?

- | | | |
|------------------------------------------|--------------------------------------------|----------------------------------------------|
| <input type="checkbox"/> a. < 1 year | <input type="checkbox"/> c. 6 to 10 years | <input type="checkbox"/> e. 16 to 20 years |
| <input type="checkbox"/> b. 1 to 5 years | <input type="checkbox"/> d. 11 to 15 years | <input type="checkbox"/> f. 21 years or more |

Thank you. The next section is about your attitude and beliefs about medication errors.

SECTION 2:

Hospital Survey on Patient Safety

Reprinted/translated with permission from the Agency for Healthcare Research and Quality (an Agency of the United States Department of Health and Human Services); Rockville, Maryland USA

Internet Citation: International Use of the Surveys on Patient Safety Culture. December 2014. Agency for Healthcare Research and Quality, Rockville, MD. <http://www.ahrq.gov/professionals/quality-patient-safety/patientsafetyculture/pscintusers.html>

Please allow 30 minutes to complete this survey. If you do not wish to answer a question, or if a question does not apply to you, you may leave your answer blank.

SECTION A: Your Work Area/Unit

In this survey, think of your “unit” as the work area, department, or clinical area of the hospital where you spend *most* of your work time or provide *most* of your clinical services.

What is your primary work area or unit in this hospital? Select ONE answer.

- | | | |
|--------------------------------------------------------------------------|------------------------------------------------------------|-------------------------------------------------------------------------|
| <input type="checkbox"/> a. More than one hospital unit/No specific unit | | |
| <input type="checkbox"/> b. Medicine (non-surgical) | <input type="checkbox"/> g. Intensive care unit (any type) | <input type="checkbox"/> i. Radiology |
| <input type="checkbox"/> c. Surgery | <input type="checkbox"/> h. Psychiatry/mental health | <input type="checkbox"/> m. Anesthesiology |
| <input type="checkbox"/> d. Obstetrics & Gynecology | <input type="checkbox"/> i. Rehabilitation | <input type="checkbox"/> n. Other, please specify: |
| <input type="checkbox"/> e. Pediatrics | <input type="checkbox"/> j. Pharmacy services | <div style="border: 1px solid black; height: 20px; width: 100%;"></div> |
| <input type="checkbox"/> f. Emergency department | <input type="checkbox"/> k. Laboratory | |

Please indicate your agreement or disagreement with the following statements about your work area/unit

Think about your hospital work area/unit...	Strongly Disagree ▼	Disagree ▼	Neither ▼	Agree ▼	Strongly Agree ▼
1. People support one another in this unit	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
2. We have enough staff to handle the workload.....	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
3. When a lot of work needs to be done quickly, we work together as a team to get the work done	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
4. In this unit, people treat each other with respect	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
5. Staff in this unit work longer hours than is best for patient care	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
6. We are actively doing things to improve patient safety	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
7. We use more agency/ temporary staff than is best for patient care	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
8. Staff feel like their errors count against them	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
9. Errors have led to remedial action and positive changes here.....	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5

SECTION A: Your Work Area/Unit (continued)

Think about your hospital work area/unit...	Strongly Disagree ▼	Disagree ▼	Neither ▼	Agree ▼	Strongly Agree ▼
10. It is just by chance that more serious mistakes don't happen around here	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
11. When one area in this unit gets really busy, others help out	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
12. When an error is reported, it feels like the person is being reported, not the problem	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
13. After we make changes to improve patient safety, we evaluate their effectiveness	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
14. We work under pressure trying to do too much, too quickly	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
15. Patient safety is never sacrificed to get more work done	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
16. Staff worry that errors they make are kept in their personnel file	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
17. We have patient safety problems in this unit	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
18. Our procedures and systems are good at preventing errors from happening	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅

SECTION B: Your Supervisor/Manager

Please indicate your agreement or disagreement with the following statements about your immediate supervisor/manager or person to whom you directly report.

	Strongly Disagree ▼	Disagree ▼	Neither ▼	Agree ▼	Strongly Agree ▼
1. My supervisor/manager says a good word when he/she sees a job done according to established patient safety procedures	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
2. My supervisor/manager seriously considers staff suggestions for improving patient safety	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
3. Whenever pressure builds up, my supervisor/manager wants us to work faster, even if it means taking shortcuts	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
4. My supervisor/manager overlooks patient safety problems that happen again and again	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅

SECTION C: Communications

How often do the following things happen in your work area/unit?

Think about your hospital work area/unit...	Never ▼	Rarely ▼	Some- times ▼	Most of the time ▼	Always ▼
1. We are given feedback about changes put into place based on error reports	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
2. Staff will speak up freely if they see something that may negatively affect patient care	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
3. We are informed about errors in this unit	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
4. Staff feel free to question the decisions or actions of those with more authority	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅

SECTION C: Communications (continued)

	Never ▼	Rarely ▼	Some- times ▼	Most of the time ▼	Always ▼
5. In this unit, we discuss ways to prevent errors from happening again ...	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
6. Staff are afraid to ask questions when something does not seem right .	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅

SECTION D: Frequency of Errors Reported

In your hospital work area/unit, when the following errors happen, *how often are they reported?*

	Never ▼	Rarely ▼	Some- times ▼	Most of the time ▼	Always ▼
1. When an error is made, but is <u>noticed and corrected before affecting the patient</u> , how often is this reported?	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
2. When an error is made, but has <u>no potential to harm the patient</u> , how often is this reported?	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
3. When an error is made that <u>could potentially harm the patient but does not</u> , how often is this reported?	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅

SECTION E: Patient Safety Grade

Please give your work area/unit in this hospital an overall grade on patient safety.

<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
A	B	C	D	E
Excellent	Very Good	Acceptable	Poor	Failing

SECTION F: Your Hospital

Please indicate your agreement or disagreement with the following statements about your hospital.

Think about your hospital...	Strongly Disagree ▼	Disagree ▼	Neither ▼	Agree ▼	Strongly Agree ▼
1. Hospital management provides a work environment that promotes patient safety.....	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
2. Hospital units do not coordinate well with each other.....	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
3. Things get missed when transferring patients from one unit to another	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
4. There is good cooperation among hospital units that need to work together	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
5. Important patient care information is often lost during shift changes	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
6. It is often unpleasant to work with staff from other hospital units	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
7. Problems often occur in the exchange of information across hospital units	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
8. The actions of hospital management show that patient safety is a top priority	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅

SECTION F: Your Hospital (continued)

	Strongly Disagree ▼	Disagree ▼	Neither ▼	Agree ▼	Strongly Agree ▼
9. Hospital management seems interested in patient safety only after an error happens.....	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
10. Hospital units work well together to provide the best care for patients.....	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
11. Shift changes are problematic for patients in this hospital.....	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅